each seat of disease beginning in the skin as the first did. The excision of one bone is useless; five or six have been excised from one foot; but the disease has recurred in those left. The man at present in hospital with this is rather old (48) to have it if being usually found in young men. In this case, the disease has not penetrated beyond the skin. The patient ruptured the gastrocnemius of the same side a year ago; the ends were not approximated; and the result—the tendo Achillis and muscle not acting on the heel—is that unusual weight has been thrown upon the diseased heel in walking. A week later, M. Nélaton spoke of the case, remarking upon the peculiar character of an opening—in the centre of a circular, depressed, black portion of skin, of the size of a shilling—which had formed, and looked “as if it had been cleanly punched out”: there was not any puriform discharge. A stick of chloride of zinc was applied to the opening. A fortnight subsequently, the disease was in much the same state, but the edges of the opening were blue. I regret that my observation of the case then ceased.

I have italicised those portions of Nélaton's description in which it appears to me that you differ with him. That description was taken during a clinical lecture, and forms part of a paper which I read before the Southampton Medical Society on the 3rd September 1867.

I am glad you have given your attention to this disease, fully concurring with you that it has not been sufficiently recognised. My interest in the case now mentioned was partly due to the disease being a new one to me at the time. I then sought in vain in several British works on surgery for information or notice of the disease.

As my object in now writing has been to extend the knowledge of a locally intractable malady, I hope you will have no objection to my sending a copy of this to the BRITISH MEDICAL JOURNAL.

I am, sir, your obedient servant,

STANLEY L. HAYNES, M.D.

Henry Hancock, Esq., F.R.C.S.

ON THE VALUE OF TINCTURE OF CANTHARIDES IN SOME FORMS OF PYELITIS.

By EDWARD MACKEY, M.B.LOND., ETC.,

Joint Professor of Materia Medica in Queen's College, Birmingham.

To Dr. Greenfield belongs the credit of having first drawn attention to the above subject; and his *Treatise on the Safe Internal Use of Cantharides*, published in London in 1706, contains some valuable observations, now not generally known. In our recent literature, good accounts of the disease may be found in Prout (*Diseases of the Stomach and Urinary Organs*, 3rd edit., p. 347) and in Roberts (*Renal Diseases*, p. 383); but neither there nor in any of a large number of authorities whom I have consulted do I find any notice of this remedy for it, except in Stillé (*Materia Medica*, 3rd edit., vol. i, p. 381).

The two following cases occurred to me some years ago, but I delayed their publication, in the hope of being able to support them by additional facts; but the disease is comparatively rare, and I relate them now, in the hope that a more extended trial may be given to the remedy, when the attention of the general body of the profession is redirected to it.

CASE I.—Agnes H., aged 34; married early; no children; by profession a musician. About the age of ten occurred the earliest symptoms she can remember; viz., smarting pain in the urethra, with frequent calls to pass urine, and violent pain in the right loin, darting downwards. After some days of suffering, she recovered for the time, but felt the same occasionally, though to no great extent, till the year 1862, when, being engaged to sing at Bristol, she caught cold. Then she had most severe smarting at the fourchette; pain up the urethra, made intense by the passage of urine, which became full of pus; and pain in the right loin. (Mr. Lang attended her; and, in addition to other treatment, removed from the meatus what seems to have been a vascular tumour, which had been cauterised once before.)

In the Bristol Hospital, under Dr. Fripp, she remained for four months, and by the end of that time was free from pain; the urine was clear, but she could not retain it longer than an hour.

At the end of the year 1862, she was in Leeds, when the above symptoms returned; and, in the course of the attack, she passed much dark blood *per urethram* for about a week. Later on, she was admitted into the Infirmary, when Mr. Teale sounded her for stone, without finding any; and she passed under the care of Dr. Hardwick, then of Mr. Wheelhouse. In addition to the other symptoms, paraplegia came on, and continued for two or three months. The part of her treatment that is most impressed on her memory is the application of

three leeches to the fundament every other day for some time. It was nearly twelve months before she was able to return to her home in Birmingham (1864). Afterwards, she continued to suffer more or less, and came under the care of a large number of practitioners in this town.

In July 1865, I first saw her as a dispensary patient. Her symptoms had been much aggravated during the last few weeks; and she lay in bed, with the knees drawn up. She had rigors at times, and thirst; there was much mental depression. She had constipation, but was able to take a fair amount of food, and there was not much emaciation. The menses were regular, always inducing aggravation of the nephritic symptoms. Her greatest torment was a constant call to micturition, occurring literally every ten or fifteen minutes, day and night. She had pain down the right leg, and often felt a darting pain along the course of the right ureter; then a sudden flow; then a severe smarting, “as of an ulcer at the neck of the bladder;” then relief. Once or twice every day, she had paroxysms of violent pain, referred to the right kidney and “passages”. A specimen of urine, taken from twenty-four hours' quantity of about two pints, was pale, slightly acid, of specific gravity 1015, with a precipitate of pure pus to the extent of two to four ounces. On some days, there were passed numbers of soft whitish rolls, narrow, about a quarter of an inch long. She described them as “skins”; and, under the microscope, they were found to consist of tessellated or polygonal epithelium. There would be, at times, much fullness of the renal region when these did not pass; and relief and subsidence of the fullness would follow the excretion of a large quantity. As to her treatment, she had already taken a great number of remedies without much relief. I saw prescriptions for iodide of potassium, copalba, alkaline diuretics, and anodynes in every shape. She found temporary ease from drinking infusion of althæa with acacia, and from hot baths and poultices; but, after inquiring into the effect of all the medicines she could remember, I found that she attributed good effects to a mixture containing one drop of tincture of cantharides, to be taken three times in the day. It had been prescribed for her by my predecessor and friend, Mr. G. Elkington, at the suggestion of our mutual friend, Dr. Dyce Brown; but circumstances had not permitted her to continue its use for more than six weeks. I began it immediately in sugared water; but her pains were so intense, that I gave her also chlorodyne once or twice a day (this she had had for many months before). She took a colocynth and henbane pill once or twice in the week, and continued the althæa and poultices, but had nothing else as remedy. Soon I increased the dose to two drops of the tincture, and so gradually to fifteen, which she took thrice daily, well diluted. At the end of a fortnight, she was decidedly easier, and lay often as much as twenty minutes without a call to micturate. This frequency was the first symptom to be relieved; the discharge of pus continued, and the pain was bad. In September, the bladder was sounded, without detecting any stone.

We persevered with the use of tincture of cantharides, in doses varying from two to twenty-five minims, with occasional intermissions. After a time, I found that it agreed best when administered with small doses of magnesia—the sulphate and carbonate. She was able to omit her stronger purgative; and, at the end of a few months, two and a half grains of soap and opium pill, once a day, replaced her chlorodyne.

In the course of two months, she was able to go down stairs; but the urgency of her symptoms returned whenever the remedy was omitted for a few days. She herself had great confidence in its power to relieve her, and saw the necessity for perseverance in its use, which she continued for upwards of two years, under my direction. Before the end of that time, however, she was able, when wearing an urinal, to do her work, and to take journeys almost as long as before her illness. The dose which she took at this time was often as much as forty drops three times in the day; and this without any ill effects whatever.

In September 1867, the urine was straw-coloured, generally clear, though showing a deposit after extra exertion. She could retain it for two hours, and could omit her remedy occasionally without any relapse. No “skins” had been passed for a long time. She seemed, in fact, fairly well, and took an engagement to sing at Leicester: this she accomplished successfully, and then started on her usual professional tour. For ten months she was actively engaged, and only required her medicine (the prescription for which she always carried with her, as a treasure) for a week or so, during a slight relapse which occurred at Sheffield. In March 1868, she carried through a harassing lawsuit in Manchester, and seemed none the worse for it. In July 1868, she was returning to Birmingham, when the train left the metals, and she was thrown violently backwards and forwards in a second-class carriage, striking especially her back. She felt the concussion much. Vomiting came on soon afterwards; and, next day, blood passed with the urine in red streaks and in small dark clots, and the urine was blood-red. She suffered also from general symptoms of illness. Her new medical

attendants prescribed at first, at her request, her old remedy, but without relief, though its use was continued for six weeks. The passing of blood ceased at the end of the first week; but the urine continued to be scanty and offensive, and with much stringy deposit, differing thus from its condition when cantharides was useful to her. In September, it showed crystals of triple phosphate; and her medicine was altered for the tincture of perchloride of iron with dilute hydrochloric acid, ten minims of each, but without benefit. She took also an effervescent alkaline mixture to allay the constant nausea. In November, she was seen by Mr. S—, her present medical attendant from the dispensary; and he informs me that she was then unable to sit up even for a few minutes, but that marked alleviation of the pain followed the use of mild diuretics. She was able to move about again in the course of two or three weeks, and has gradually improved up to a certain point. She took at first the dilute acids with juniper; then the acetate of potash with juniper, in decoction of pareira; and continued also her effervescent saline, and opiate and aperient pills. In February, she began the bitartrate of potash with scopolium. At present (April), she is able to go out a little, and can retain urine for half an hour, sometimes for an hour. The amount passed is about two pints in the twenty-four hours; it is slightly acid, of specific gravity 1008, with much purulentropy deposit, no blood. The old "obstructive" pain, followed by passage of "skins", has occurred four times since the accident. She now suffers more aching throbbing pain than she formerly had over the left kidney and in the left groin, and has also pain over the position of the right ureter and in the "passages". This unfortunate present condition does not invalidate the evidence of relief afforded previously, but only suggests that some alteration has taken place either in the position of the calculi or in the kind of irritation existing; and proves, what is perfectly rational, that cantharides, though good in some forms of the disease, is not in all. Good, perhaps, it may be, in chronic irritation—not good in acute.

CASE II.—Jane E., aged 28, married, when a child had pains in the back and right side. She remembers that, at the age of sixteen, pus was noticed in her urine; and then she had frequent and painful micturition. During such attacks, she had vomiting, pain in the back, and general illness. In the intervals, she was able to continue her work as a servant. Later, menstruation came on, and occurred slightly every two or three months. At various times, she came under the care of many medical men, to some of whom the disease appeared to be "rheumatism", to others "unknown"; but both Dr. Wade and Mr. Schofield considered that it was "abscess of the kidney, probably from stone."

At the age of twenty-two, she married, and had a miscarriage, then a premature confinement, then two living children. In January 1866, she had "inflammation of the bowels", and had been failing rapidly since. On September 5th, 1866, I saw her for the first time. She was much emaciated; her face was sallow, and bore an expression of constant agony. She had "nervous irritability", headache, vomiting, and cough (no special lung-sounds), numbness of the right leg, and intense pain over the right kidney, worse at times, when the urine was almost suppressed. These paroxysms had come on lately, at intervals of only a fortnight. The urine was usually offensive, and contained much pus and some blood. I ordered her, at first, bismuth and magnesia, with a little morphia; quiet and nourishment, and baths. When the stomach-symptoms were relieved, I added (September 15th) two minims of tincture of cantharides to the mixture. On October 6th, she was better, and increased her dose to three minims, congratulating herself that her bad attack had been deferred longer than usual. On October 8th, however, it began with intense prostration, pyrexia, and complete suppression of urine, with frightful pain. I stopped the above remedies, and gave opium with alkalis in effervescence, enemata, and poultices. On October 11th, she was exhausted, with sordes on the teeth, brown dry tongue, glazed eye, almost dying, when urine began to flow, and with it pus, and many curious lumps, varying from the size of a pea to that of a hazel-nut, rolled up, soft, whitish outside, brown within, much like toadstools. Many of the lumps contained gritty matter and small dark irregular masses, probably carbonate and phosphate of lime. On October 25th, she was able safely to resume three minims of tincture of cantharides thrice daily, in combination with quinine and a few minims of chlorodyne; and, under this treatment, she recovered more rapidly than usual. On November 10th, she had been up ever since, attending to her house-work. She had continued her last medicine, and considered herself better in health than for many years. She had still frequency of micturition, pain over the kidney, and numbness of the legs, though in less degree.

She remained under my care for a month or two longer, without another bad attack. Then she was obliged to remove to some distance, and I did not see her alive again. I heard that she frequently

expressed to her family her wish for more of the medicine last mentioned.

She died in April 1867, in one of her attacks; and Mr. Schofield, who made an examination of the body, kindly invited me to be present. The ordinary results of former peritonitis were observable. The left kidney was atrophied, and was little else than a sac containing a pulsataceous mass of pus and mineral matter. The right kidney was larger, and contained several calculi of phosphate and carbonate of lime, possibly of oxalate. The ureters were completely obstructed.

In estimating the value of cantharides in these two cases, I do not omit consideration of the possible good effects of other medicines taken during part or all of the same period—magnesia in the first case, chlorodyne (containing morphia, cannabis, and capsicum) in both; only, I would record that observation convinced me that tincture of cantharides was the principal, if not the only, agent which gave marked relief to the renal symptoms at certain periods of their progress.

CASE OF LOCKED TWINS.

By MARTEN PERRY, M.D., Evesham.

M. S., primipara, aged 28, menstruated last at the beginning of September, and expected her confinement at the end of May or beginning of June 1869. During the pregnancy she had enjoyed her usual health till the first week in March, when I was called in to attend her for severe muscular pains on the right side of the chest from incessant coughing. She was quickly relieved, and well again in a week or less. Slight labour-pains set in about three or four o'clock in the afternoon of Friday, March 19th, but did not become severe till about 2 P.M. on the following day. The membranes gave way spontaneously about four o'clock, and a midwife was then summoned. She examined her patient about five o'clock, and states that she found a foot presenting; but, considering herself competent to manage the case, did not send for assistance until she had in vain tried her skill at traction.

I first saw the case about 9.30 P.M., and ascertained that a breech was already born. There was no pulsation in the cord; the legs and body were cold and livid; and the child's head was looking towards the pubes of the mother. I proceeded to draw down first one arm and then the other, without encountering anything unusual, and, as the head did not even then follow, I introduced the forefinger of my right hand with a view of placing it in the mouth of the foetus and thus expediting delivery. Instead, however, of meeting with a mouth, I found, to my great surprise, an eye. I then sought and readily detected its fellow, and above them the nose and mouth. Having withdrawn my right hand, I introduced the left at the back of the neck and fairly passed it up between the neck of one child and the face of another, this face looking also towards the pubes of the mother. My next step was to try whether I could press back above the brim the second head; but, not succeeding in this, and being satisfied that, though the pains continued very severe, no progress of moment had been made since the arms were brought down, I concluded that nature unassisted would be unable to effect delivery without great risk to the mother, and determined to have the opinion of another medical man before undertaking what I foresaw would be necessary to enable the children to be born. I thereupon gave the patient fifty drops of tincture of opium, and dispatched the husband to Evesham for assistance.

My friend, Mr. A. H. Martin, rode over at once, and was with me in about an hour. During the interval the pains had continued violent, but though the head of the second child had descended to the perineum, it was too evident that something must shortly be done. Mr. Martin agreed with me as to the steps to be taken in this emergency, and, at 0.15 A.M., seized the shoulders and forcibly drew them down. I then retracted the integuments of the chest and shoulders as much as possible, and (while he protected the maternal structure) divided them low down with a bistoury. The third and fourth vertebrae having been already dislocated by traction, there was a large flap to cover the bones, and protect the vagina from injury. The pains then subsided for some time, but on their return the second head began to advance. We then pressed upwards the disarticulated head, and retaining it during one pain as high as possible, the head of the second child came down more readily, distended the perineum, and was speedily expelled with its face still towards the pubes. Mr. Martin then turned its face towards the right hip (upwards), and, at the next pain, the *detruncated head* descended at the back of the neck, distending the perineum to an immense extent, and was at once expelled between the thorax of the second child and the coccyx of the mother. The body of No. 2 speedily followed, still-born, and about ten minutes afterwards the two placentae. These were perfectly distinct, and connected only by the membranes. The peri-