A LECTURE

ON

HÆMOPHTYSIS: ITS CAUSES, RESULTS,
AND TREATMENT.

By GEORGE JOHNSON, M.D., F.R.C.P.,

Physician to King's College Hospital; Professor of Medicine in King's College,
London; etc.

SINCE the time of Laennec, it has been very generally assumed that hæmoptysis, when not associated with valvular disease of the heart, or vicarious of the catamenial discharge, is almost always a result and an indication of tubercular disease of the lung. I have long known and taught that this doctrine, thus broadly stated, involves a large amount of error. I have seen a considerable number of cases of hæmoptysis in which there has been no evidence of structural disease within the chest, either at the time or for months and even years afterwards. Spitting of blood is a symptom which at the best is sufficiently alarming; there is no need to aggravate its terrors by the erroneous assumption that it is almost invariably associated with serious organic disease either of the heart or the lungs.

Let a man who has once spat blood apply to an insurance-office, and the probability is that he will be rejected, or required to pay a large increase of premium; yet it may be that the blood-spitting was as much the result of a harmless accident as if the nose had been the source of the bleeding. The lung is by far the most vascular organ in the body. In addition to its own nutrient bronchial vessels, the whole of the blood from every other organ passes through the pulmonary capillaries. These two systems of vessels in the lung, in consequence of their relation to each other, to the heart, to the movements of the chest, and to the function of respiration, are more liable to sudden strain and pressure than the blood-vessels of any other organ; and it would be indeed marvellous, if bleeding did not frequently occur from their accidental rupture, without previous disease.

Now let us pass in review the chief known causes of hæmoptysis; and by hæmoptysis in this lecture I mean hæmorrhage from the air-passages, excluding cases of bleeding from the gums or fauces.

First, then, tubercular disease of the lung is a frequent cause of hæmoptysis. Certain of the pulmonary capillaries are compressed by a tubercular deposit; the surrounding vessels are subjected to increased strain; they consequently give way, and hæmorrhage occurs. In a more advanced stage of the disease, when softening of the deposit is in progress, blood-vessels may be opened by ulceration. In some instances, considerable branches of the pulmonary artery in tuberculous lungs become aneurismal; and the rupture of such an aneurism may be a source of sudden, copious, and even fatal hæmorrhage.

Disease on the left side of the heart is a well-known cause of hæmoptysis. Suppose an incompetent mitral valve: the left ventricle drives a portion of blood forcibly backwards into the lungs, while the right ventricle continues to propel the blood onwards. The pulmonary capillaries, strained by the increased pressure upon their walls, give way, and the result is hæmoptysis and pulmonary apoplexy.

Disease on the right side of the heart is comparatively rare; but, when it does exist, it not unfrequently causes bronchial hæmoptysis. In the *Medical Times and Gazette* of February 13th, 1858, I published the case of a woman whose sole organic disease was great dilatation of the tricuspid orifice. The whole systemic venous system was much engorged, in consequence of the reflux of blood through the tricuspid orifice. The bronchial veins and capillaries were partakers in this engorgement, and the result was rupture of bronchial capillaries and hæmoptysis. In cases of congenital narrowing of the pulmonary artery, hæmoptysis has been a frequent symptom. This, again, is a result of a retrograde engorgement of the bronchial veins and capillaries. These patients, too, have frequently become phthisical. I shall presently refer to the occurrence of phthisis as a *consequence* of pulmonary hæmorrhage.

Bronchial hæmoptysis, resulting from disease on the right side of the heart, will help us to understand the hæmoptysis of apnoea. Thus hæmoptysis, to a variable extent, is an occasional accident in cases of spasmodic asthma. What is the explanation of this? Bronchial spasm limits the supply of air to the pulmonary capillaries. In consequence, the minute pulmonary arteries, by their contraction, check the onward movement of the blood. The systemic arteries are comparatively empty,

while the whole systemic venous system is distended. The bronchial veins and capillaries share in this distension; and hence bronchial mucous exudation, and occasionally bronchial hæmorrhage.

Active inflammatory congestion of the pulmonary and bronchial vessels is a frequent source of hæmorrhage. In most cases of pneumonia, the expectoration is more or less blood-tinged. The mucous expectoration of bronchitis is not unfrequently mixed with blood. In both pneumonia and bronchitis, profuse hæmoptysis is an occasional occurrence, quite unconnected with tubercular disease.

About three years ago, I attended one of our former house-surgeons with bronchitis. He had profuse hæmoptysis for several days in succession. I felt very anxious about him; but there were no positive physical signs of tubercular disease; and I always hoped, and expressed hope. He completely recovered, and is now actively at work. I have seen several cases of the same kind.

Plastic bronchitis is a rare disease. I have met with it only once. When it does occur, hæmoptysis is a frequent symptom.

In cases of primary cancer of the lung, the sputa are more or less blood-tinged, and not unfrequently assume the appearance of red currant-jelly.

In several cases of hæmoptysis that have come under my observation, emphysema of the lung has been the only structural change that I could discover. In most of these cases, the exciting cause of the hæmoptysis has been either an attack of bronchitis or over-exertion. In one case, a fat lady with emphysematous lungs, while suffering from an attack of catarrh, had profuse hæmoptysis after walking up stairs. She died some years afterwards, of Bright's disease; but had no symptom of any other pulmonary disease than emphysema, with occasional catarrh.

A former clinical clerk, a powerful muscular man, who has overstrained his lungs and rendered them emphysematous by excessive gymnastic exertion, has twice had hæmoptysis after playing at football. He remains in good health.

Without doubt, excessive exertion may cause rupture of pulmonary vessels and hæmoptysis in persons whose lungs and heart are perfectly sound; but it is obvious that vesicular emphysema is a predisposing cause of hæmoptysis from over-exertion. Vesicular emphysema is associated with more or less obliteration of pulmonary capillaries, and with hypertrophy of the right ventricle consequent on the impeded circulation through the lungs. Therefore, when the heart's action is excited by active exercise, the pulmonary capillaries are injected with great force by the strong right ventricle, and the result may be rupture of vessels and hæmorrhage.

In a considerable number of recorded cases, the irritation of the air-passages by a foreign body introduced through the larynx has been attended with bronchial hæmoptysis.

It is obvious that a blow on the chest, either with or without fracture of the ribs, and a consequent wound of the lung, may rupture vessels, and cause hæmoptysis. So an aneurism of the aorta, or of one of its primary branches, may open into the air-passages, and cause hæmoptysis.

In all the cases of hæmoptysis to which I have referred, the hæmorrhage results from various forms of structural disease or injury, either in the lungs, or heart, or blood-vessels; but pulmonary hæmorrhage may occur unassociated with any appreciable structural change within the chest. In some of these cases, deterioration of blood, and consequent weakening of the walls of the vessels, are the probable causes of the hæmorrhage. Thus hæmoptysis is one of the forms of hæmorrhage which is not unfrequently associated with scurvy, with purpura, and with uræmia. I have repeatedly seen both profuse hæmoptysis and epistaxis as results of an habitual excess of alcohol and a deficiency of nutritious food.

Hæmoptysis is occasionally vicarious of the menstrual discharge. In cases of this kind, we must guard against deception. I once had a patient whose blood-spitting was supposed to have this origin. On examination, however, I discovered the source of the bleeding in numerous scratches which she had made with a needle in the roof of her mouth. I believe that this case is a type of some other supposed cases of vicarious hæmoptysis.

Lastly, there are cases in which there is no apparent cause for the hæmoptysis; and, to conceal our ignorance, we speak of the hæmorrhage as resulting from a hæmorrhagic diathesis. The following case, recently in the Hospital, is an example of this (vol. xxxix, p. 75).

John Herring, aged 37, a blacksmith, was admitted Dec. 18th, on account of hæmoptysis, which began ten days before his admission. He said that, for a fortnight before he began to spit blood, he had suffered from pain in the right side of the chest. The hæmoptysis began on the 8th December, and has continued daily until his admission. One day, the 11th December, he brought up as much as three-quarters of a pint. The blood, he said, had varied in colour, sometimes being dark, at other times bright red. He said that twelve years ago he had

pleurisy; that five or six weeks after this he first spat blood, and that he had since had frequent attacks of hæmoptysis. He had some times passed nine months without an attack; at other times he had had two attacks within a week. The blood-spitting had generally continued from twenty-four to thirty hours. If it continued longer, he usually stopped it by a dose of elixir of vitriol. Three years ago, he had inflammation of the lung, and was laid up for six months, but during this illness he had no hæmoptysis. He said that he had always been temperate. His father died of old age (upwards of 80); his mother from "change of life, and something wrong in her head." He has three brothers and a sister in good health; none have died; his grandparents all lived to old age. On admission, he was a fairly nourished man; but his face and lips were pallid. He complained of a troublesome cough, and he expectorated a quantity of dark yellowish-brown blood-tinged material, having a fetid odour; his breath, too, was fetid, and there was some pain over the right side. Tongue coated, appetite bad, bowels costive, urine normal; the skin was moist; at night, sometimes profuse perspiration; pulse, 66; temperature, 99.8. Percussion gave natural resonance over the whole front of the chest; behind, there was some dullness on percussion on the right side, from the spine of the scapula to the base of the lung; over the dull space there was rather large crepitation, with diffused blowing expiration; the vocal resonance and vibration were unchanged over the dull space; elsewhere the respiratory sounds were normal, both front and back; the heart's sounds and action were normal. He was ordered to take ten grains of gallic acid every three hours; to inhale the vapour of turpentine and boiling water night and morning. On the 22nd December, the expectoration had ceased to be blood-tinged; it was of a dark-greenish colour, and still had an offensive odour. Temperature, 98.5; the dullness and crepitation over the lower lobe of the right lung were diminished; he was ordered to inhale creasote instead of turpentine. He continued steadily to improve; the expectoration gradually ceased; and he left the Hospital on the 15th January. The day before his discharge, it was noted that there was normal resonance over the right lower lobe; inspiration was vesicular and free from crepitation. The only abnormal sound was somewhat prolonged expiration over the right back.

Now here is a case in which a frequently recurring hæmoptysis, extending over a period of twelve years, is pretty certainly not associated with organic disease of the lung either as a cause or a consequence. When he came in, the physical signs indicated that the lower lobe of the right lung was partially consolidated, probably by blood, which had been driven into the extreme bronchi and air-cells, where it appears to have decomposed and become fetid. This blood was gradually expectorated; the lung has recovered its normal condition; and the man has been restored to his usual state of health.

Although I have long known that hæmoptysis not unfrequently occurs unconnected with phthisis, it is only recently that I have learnt that pulmonary hæmorrhage is an occasional exciting cause of phthisis. Laennec taught us to believe that, when profuse hæmoptysis is quickly followed by the symptoms of a rapid phthisis, tubercles were latent in the lung and caused the hæmorrhage. Niemeyer has lately revived the doctrine, which was accepted before the time of Laennec, that a copious pulmonary hæmorrhage, in a person previously healthy, may be an exciting cause of phthisis. (See Niemeyer's *Text-Book of Practical Medicine*, translated by Drs. Humphreys and Hackley of New York.) Niemeyer's doctrine is, that a portion of the blood, being driven down to the ultimate bronchi and air-cells, acts as a foreign body, and sets up inflammatory and degenerative changes in the lungs, from which the patient may slowly recover, or which may result in cheesy deposits, excavations, and fatal phthisis.

In the second volume of the *Clinical Society's Transactions*, there are two interesting papers; one, by Dr. Christian Bäumler, on "Cases of Hæmoptysis followed by Inflammatory Changes in the Lungs"; another, by Dr. Hermann Weber, on "Hæmoptysis as a cause of Inflammatory Processes and Phthisis." These papers are confirmatory of Niemeyer's doctrine; and I am sure that in the main the doctrine is true, for it has rendered intelligible to me some facts in my past experience which before I could not comprehend.

On the 10th May, 1867, I first saw a married lady, aged about 25, who, a month before, during a violent fit of coughing, had brought up a large quantity of blood. Since the hæmoptysis, she had been weak and ill, but not confined to bed. I found all physical signs normal at the upper and front part of the chest; but, over the right base at the back, there were some dullness on percussion, and rather loose and large bubbling rhonchi. The family history indicated no tendency to phthisis. The history and the physical signs were not those of tubercular disease, of ordinary pneumonia, or of ordinary bronchitis. I have no doubt that, consequent on the pulmonary hæmorrhage, blood had been driven into the lower lobe of the right lung, and had there excited inflammatory

changes. I saw this lady occasionally for some months, during which she gained strength, but the physical signs remained unchanged. I have heard quite recently that her health has continued to improve; but her medical attendant in the country says there is still dullness and crepitation over the right base, and still absence of abnormal physical signs over the upper and front part of the chest. There is reason to hope that ultimately she may recover from the accidental results of the pulmonary hæmorrhage, though the long continuance of the physical signs gives the case a somewhat serious aspect.

A few months ago, I saw a case in which the fatal consequences of pulmonary hæmorrhage were of an unusual character. A lady, about 40 years of age, healthy, but of a somewhat delicate frame, during the early part of September, had a troublesome dry cough, and one day she began to cough up a large quantity of florid blood. So rapid was the hæmorrhage, that at one time she was nearly suffocated, and the active bleeding continued for several hours. I first saw her at Eastbourne, on the 17th September, a week after the onset of the bleeding. She was then expectorating small quantities of semi-solid dark blood, which had evidently been for a considerable time out of the vessels. Percussion and auscultation over the front of the chest gave quite normal results; and, as the chest-walls were thin, the air could be heard entering the lung with great clearness. Over the right lower lobe at the back, there was marked dullness on percussion, and, on auscultation, rather fine crepitation, with diffused blowing expiration. The inference was, that the right lower lobe had been partially consolidated by blood driven into its tubes and vesicles. Percussion and respiratory sounds over the left back were quite normal. The pulse and breathing were somewhat quickened, but there was no febrile excitement. A few days afterwards, I heard that the chest-symptoms were better, but that the left leg below the knee had become painful and swollen, though not inflamed; and I suggested that probably some coagula from within the pulmonary capillaries, having entered the circulation, had led to the formation of coagula within the capillaries and veins of the leg. On the 4th of October, I again saw the patient. I found that, during the last few days, the painful swelling of the leg had entirely subsided, but this had been succeeded by urgent dyspnoea. The breathing was still upwards of 40 in a minute; the pulse rapid and feeble; the countenance anxious; the face pale; the lips livid. There was now normal resonance over the lower lobe of the right lung, the air entered freely, and there was scarcely a trace of crepitation left. Over every other part of the chest, percussion and respiration were quite normal. The blood which, for a time, had blocked a portion of the lung-tissue, had been expectorated, and evidently there was nothing in the state of the lung to explain the alarming dyspnoea. What, then, was its cause? Probably, a fibrinous clot, from the temporarily obstructed vein in the leg, had made its way to the right side of the heart, and there was embolic obstruction of the pulmonary artery. The dyspnoea was due, not to want of air in the lungs, but to want of moving blood. After my visit, the dyspnoea continued and increased, and she died early on the morning of the 6th October.

There was no inspection of the body, but there can be scarcely a doubt as to the sequence of events. Pulmonary hæmorrhage led to impaction of blood in the lower lobe of the right lung; this was soon expelled; but, meanwhile, either blood materials absorbed from the air cells by the pulmonary capillaries, or, probably, coagula formed within the pulmonary capillaries while the blood was rendered stagnant by pressure from without, passed into the circulation, and led to the formation of coagula within the capillaries and veins of the leg; thence coagula found their way back to the right side of the heart, where they increased and caused a fatal obstruction to the circulation.

You see that pulmonary hæmorrhage is a subject which demands a careful and diligent study. Each case requires a minute and accurate investigation before we can venture to give an opinion as to the cause or the probable consequences, or the appropriate treatment. There are few cases in which an off-hand inconsiderate opinion is more likely to be erroneous, and, therefore, unjustifiable, than in these alarming cases of blood-spitting. The fact that hæmoptysis is often associated with serious organic disease, either as a cause or a consequence, renders it the more important that the antecedents and the circumstances of its occurrence should, in every case, be thoroughly investigated.

The treatment of pulmonary hæmorrhage will vary somewhat according to the nature of the exciting cause. There are, however, certain general rules which are applicable to all cases. The patient must remain as absolutely at rest as possible. Bodily exertion or emotional excitement, by increasing the force and frequency of the heart's contractions, is apt to increase the bleeding, or to provoke a return. The patient should lie still, and neither move, speak, nor cough, more than is absolutely necessary. A cough is a powerful provocative of pulmonary hæmorrhage, and it is sometimes desirable to allay irritation

and cough by a dose of morphia, or by a cautious inhalation of chloroform from time to time. Do your utmost to prevent the patient from being alarmed and excited by the sight of the blood. Let him have a plentiful supply of cool fresh air. Let him keep lumps of ice in the mouth and swallow the cold water. Sometimes the application of ice to the chest has a powerful effect in arresting the bleeding. It should not be continued long enough to chill the patient; and it is more likely to be effectual in exciting sympathetic contraction of the deeper arteries when applied for a short time over different parts of the chest, both front and back, than when applied continuously over one part. Gallic acid I believe to be one of the best styptics in cases of pulmonary hæmorrhage. It may be given in doses of ten grains, in the compound infusion of roses, every three hours, if the bleeding be copious, and less frequently as the bleeding subsides. Another useful styptic is the liquid extract of ergot. This, too, may be given every three hours, in doses of half a drachm. It may be given alone or combined with the gallic acid mixture. The oil of turpentine in 20-minim doses sometimes succeeds when other remedies fail; and I have seen the bleeding quickly arrested by the inhalation of turpentine vapour. We directed our patient, Herring, to inhale, first turpentine, and subsequently creasote, for two reasons: first, to correct the factor of the expectoration and the breath; and, secondly, to promote the expulsion of the decomposing blood from the lung, and thus prevent the risk of mischief from its retention. You will find the inhalation of turpentine vapour an excellent expectorant in this and other cases where the object is to expel accumulated material from the air passages. An emetic of sulphate of zinc may sometimes be given for the same purpose. Dr. Weber, in the paper before referred to, recommends a combination of antimony and ipecacuanha as an emetic, partly to expel blood from the lung, and partly to arrest bleeding.

ON A CASE OF PARASITIC DISEASE PRODUCED BY THE LARVA OF THE CESTRUS BOVIS.*

By ROBERT WALKER, M.D., Carlisle.

IN September 1859, when I was in Shetland, a young woman, 22 years of age, consulted me, complaining of an acute pain behind the right shoulder. She had felt it first about a fortnight before in the lower part of the back, and it had gradually ascended and had lately become more intense. On examining her back, I found a red line somewhat resembling an inflamed absorbent, but having a rather tortuous course, and extending from the left loin to about the middle of the right scapula. Towards its lower extremity it was indistinct and had a purple tinge, but at its upper extremity was very red and acutely painful. A little in advance of this extremity was a slight puffiness of the skin, beneath which a hard substance could be indistinctly felt. Having read a paper on this disease written by the late Dr. Spence of Lerwick, I had no difficulty in diagnosing the presence of the larva of the *Cestrus Bovis*, and, cutting down upon the above-mentioned hard substance, I ejected the intruder.

The case, so far as it goes, is a good example of the disease as it is usually seen in Shetland; but when the larva—to whose travels beneath the skin the symptoms I have described owe their production—is undisturbed, it eventually becomes stationary; a furunculus forms, and at its apex a small orifice appears, through which a little bloody serum, mixed with pus, for a while exudes, and the grub at last escapes and falls to the ground.

The *Cestrus Bovis*, or gadfly, infests cattle towards the end of summer, and the sight of it drives them almost mad with terror. Although it is contended by some that the female deposits her eggs on the hairs of cattle, it is generally believed that she inserts them into the cellular tissue by means of an ovipositor. After a short time, small tumours appear on the back of the infested animal, each having an orifice at its apex, from which exudes a sanguineous discharge, mixed with pus, and from which at length a grub, commonly called a warble, emerges. Another species of *Cestrus* infests the reindeer, causing a very similar affection; but other species which infest the horse and sheep produce disorders of a very different nature.

In South America, this parasite seems to have been very frequently observed on man. It has been frequently seen in the human subject in Shetland; but invariably, I believe, in women. In stating it to be the larva of the *Cestrus Bovis*, I follow Dr. Spence. Mr. Bracy Clark also considers the human parasite to be the *Cestrus Bovis*; and Küchenmeister says we are not justified in supposing that there is an

Cestrus Humanus. The rapid motion of the larva under the skin of the human subject, whilst it is stationary in the ox, may probably be explained by the different characters of the areolar tissue in the human and bovine animals.

NOTES ON SOME FEIGNED CUTANEOUS AFFECTIONS.

By C. HILTON FAGGE, M.D.,
Assistant-Physician to Guy's Hospital.

IN a paper on "Feigned or Hysterical Diseases of the Skin", published in this JOURNAL for January 8th, 1870, Mr. Startin refers to the Museum of Guy's Hospital as containing a model of an affection produced by nitric acid. The model in question is, I believe, that numbered 298 in the Catalogue, and described as representing "the arm of a young woman, showing gangrene of the skin, believed by Dr. Addison to be the result of the application of some mineral acid." It was made many years ago, and has been altered by time, so that its original character is perhaps not clearly to be recognised; but it presents numerous irregular patches of very considerable size, which have sinuous outlines, and, as it were, fit into one another; being, however, separated by intervals of healthy skin. From some of the patches the sloughs are being detached; and in one instance this process has been completed, a granulating surface being left. No history of the case has been preserved.

Within the last few weeks, a patient has been under observation in Guy's Hospital who almost beyond doubt had produced an artificial gangrene of the skin of her face and chest by the application of nitric acid. She was originally a patient of Mr. Arthur Roper of Blackheath, who sent her up to me as an out-patient. I advised her to come into the hospital; and she was accordingly admitted into the Clinical Ward, under Dr. Habershon's care. With Dr. Habershon's permission, I communicate the following notes of her case, which were taken by Mr. Wilson Eager, the clinical clerk.

Caroline S., aged 18, was admitted December 14th, 1869. On admission, she had on her face and chest several dry, apparently gangrenous patches. These appeared to present different stages. The more recent ones were quite smooth, and of a yellowish-white colour; some with a more or less greenish tinge. Those of older date were uneven, and of a dirty brown hue. Over some of these patches there was absolute loss of sensation; others were exceedingly tender. Round the margins of two patches beneath the clavicles, the skin was distinctly raised into irregular blebs with a wrinkled flaccid membrane, very like blisters produced by some local irritant. Round the circumference of the patches there was a narrow well-defined zone of bright redness. In addition to the eschars or gangrenous patches, there were beneath the clavicles irregular erythematous spots, of about the size of five-shilling-pieces. On the neck, the cuticle was in several places peeling off in irregular brownish flakes.

The history of the affection given by the patient was as follows. In the afternoon of December 8th (eight days before her admission), she felt a burning sensation on the left side of her neck; she looked in the glass, and saw that there was a blister. Soon afterwards, other blisters appeared on different parts of the neck, preceded by tingling. On the following day, several patches were visible on her cheeks and forehead; and, on the 12th, one appeared on the chin. No fresh ones subsequently formed until the morning of her admission, when two or three were beginning to appear on the upper part of the chest.

In their examination of the case, the clinical clerks at once noticed that the tips of the middle, index, and ring fingers of the right hand, the palmar surface of the middle finger of the left hand, and the palm of the left hand itself, all presented stains of a bright yellow colour, exactly like those to which nitric acid gives rise. On the ring finger, the yellow colour ran along the sulcus between the side of the nail and the skin; the adjacent parts of both being stained in the same manner.

There could hardly have been stronger circumstantial evidence that nitric acid had been used than was afforded by this staining of the nail; but the girl maintained that she had used no acid or irritant substance of any kind, either in her work or otherwise. The marks on her fingers, she said, came of their own accord, with pain, like the patches on the chest and neck.

She was a well-nourished, healthy girl, but very impressionable. This was particularly noticed by Mr. Towne, who says that she started at the least touch.

On the 17th, while she was in a warm bath, her bed and locker were searched; but no evidence of trickery was discovered. No stains of

* Read before the Cumberland and Westmorland Branch.