The case of C. H. confirms the opinion of Laennec, that pulmonary gangrene does not follow ordinary, but a septic, type of pneumonia, as it also verifies his observation that it is often preceded by anthrax. The comparative lowness of the temperature throughout the disease, and the pathological appearances of the lung, sufficiently indicate the character of the inflammatory process which had preceded the gangrene.

C. H. undoubtedly experienced relief from the inhalation of carbolic acid vapour, which also afforded much comfort to those who had to wait upon him, and who were at first terribly inconvenienced by the odour of his breath. It is probably to pulmonary admittaria that we must look for alleviation in this intractable disease.

MUCOUS DISEASE.*

By WALTER WHITEHEAD, F.R.C.S.Ed.,

Surgeon to St. Mary's Hospital, Manchester.

THIS disease has been noticed by some of the ancient, and described by a few of the more modern, physicians, but owing to those two prominent, yet opposite errors, for ever being repeated throughout medical literature—the classifying a number of diseases under one genus, and the giving to one disease a multitude of names, this disease has been more frequently concealed under the name and description of other disorders; and, consequently, it has been imperfectly handed down, and received but occasional recognition even in the best of our standard works on medicine. In proof of this, I may state that out of the one hundred and twenty writers on this disease I have been able to collect, I cannot refer to any half dozen who have described it by the same name. The disease has been principally observed in connection with the intestines, the colon being by far the most frequent portion affected; and the remarks I have to make will refer principally to this form of the disease.

Before I enter into a general consideration of the disease, I will endeavour very rapidly to give you, in somewhat chronological order, a brief review of the labours of our predecessors in this field of inquiry. Amongst the earlier writers on this subject, I may mention Celsus, Fermelius, Van Swieten, Morgagni, Berengerius, Gabucinus, Platerus, Sennertus, Tollincoferus, Bonnet, Vesalius, Hoffmann, Trælingius, Brunner, Marcard, Stoll, Theden, and others, who evidently recognised, though with an imperfect light, the nature of this disease, as well as its grave importance. Celsus has described the pathological products of this disease, under the name of pituitous and viscous matters; and we find Morgagni, in his thirty-first letter, which treats of fluxes of the belly, saying:—"For, as the glands of the bladder when irritated secrete a greater quantity of humour, and not of the same nature with that which they secrete in perfect health, so the glands of the intestines likewise secrete a greater quantity of humour, and of a different nature; for which reason, in both cases, a white and mucous matter appears." He states that in this manner may be understood "the shreds of membranes, and even of large membranes that are said to have been discharged" by the bowels. He goes on to say—"That which we now call a mucous, or polypous matter, the ancients were in general accustomed to call pituitous, and viscid." He observes that Jacobus Berengerius has written the following words:—"and I myself have seen that concretions, like pieces of thick leather, have been generated in my intestines from pituita." Fermelius supposed the matter of a firm body which was a foot in length, and pierced through the middle with a duct, to be of the same kind; which body was discharged by the Ambassador of the Emperor Charles the Fifth, who was by that means restored to his former health. Lancisi confirmed, in many different ways, that the pituitous concretion which Justus Lipsius discharged by stool, in the shape of the intestines, and believed to be the very intestines themselves, belonged to this class of disorder. Tollincoferus was also of this opinion. In further discussing these pituitous or mucous humours, he (Morgagni) informs us that Lentilius has exceedingly well determined what intestine it was whence the membranes proceeded, which was excreted in the stools at intervals with or without blood, and in a greater or in a lesser number, "and when dried, they resembled the pellicles of a hog's bladder when shrivelled up with heat."

But Johann Maurice Hoffmann, when he examined very attentively membranes that were excreted by a woman, which every other person had taken for the internal coat of the intestines, found them to be a viscid mucus "coagulated upon the valvulæ conniventes, and condensed into the form of a membrane." Nor does the opinion of Trælingius, in the fifth volume of the *Acta Naturæ Curiosorum*, differ from this; for he, on examining a kind of tube or sheath, discharged by the

wife of a peasant, found it to be not really membranous, "although it resembled the figure of the colon, with its valvulæ conniventes." Again, the celebrated Verdriesius, having examined very closely one discharged by another patient, found it evidently to be nothing else but "a fistulous pituita which had concreted together, and accommodated itself to the figure of the intestine wherein it was contained." Apinus also mentions a patient who discharged membranes "of a form exactly tubular." Gabucinus, Platerus, and Sennertus taught that certain worms belonging to the bowels were not in reality worms at all. Kämpf, a Dutch physician, was the first to establish this morbid condition, under the title of Infarctus, as a particular disease; and he appears to have written a book with the express object of proving that there were few diseases which were not originally referable to infarctus. His exaggerated conclusions were, however, by no means warranted by his previous propositions.

Referring to more recent authors, Dr. George Man Burrows gives an account of a lady of phlegmatic temperament who voided by stool, some hours after the cessation of an extraordinary attack of pain, a white opaque substance resembling a piece of tape folded like a skein of thread, and weighing nearly an ounce; but it was believed twice that quantity, which had been thrown away, had passed with the stool. This lady had suffered from frequent pregnancies and repeated abortions. This substance was analysed for Dr. Burrows by Mr. A. T. Thomson, who concluded that it consisted of animal mucus combined with a small portion of albumen. The pathological opinions expressed in reviewing the complaint of this lady are well worthy of record, especially when we take into consideration the period of the writing. The writer says—"perhaps a conjecture might be advanced that it originated in a retrograde action of the lymphatics."

Next in order of time, we find a paper read by Dr. Powell at the College of Physicians, on the 17th of May, 1818. From this paper we gather that in all the cases to which he refers the leading symptoms were similar, and led him to suspect the passage of biliary concretions; but instead of finding, as he anticipated, gall-stones passed with the motions, he discovered a large quantity of flakes, mostly torn into irregular shapes, and appearing to have formed parts of an extensive adventitious membrane of no great tenacity or firmness. Dr. Powell considered it perfectly fair to suggest that they might be poured out from the exhalants. He further observed that the most remarkable circumstances in the history of his cases, was the production of an effect usually ascribed to inflammatory action without its previous existence.

We next refer to Dr. Good, who states, in his *Study of Medicine*, under the title of Diarrhoea Tubularis, and also Tubular Looseness:—"I have never hitherto seen this species arranged, and not often described, but it occurs frequently in practice." He describes the dejections as consisting more or less of membranous-like tubes, whitish, viscous, and inodorous. Dr. Good mentions the examples of the disease given by M. Bauer in his letter to M. de Hahn, and also that Spindler in his 45th Observation describes a like case, in which the secretion was worked up into a "materia alba longa compacta." Dr. Good considers that the disease appears to depend upon a peculiar irritability of the villous membrane of the large intestines, which, in consequence, secrete an effusion of coagulating fibrin, fibrin mixed with albumen instead of secreting mucus, occasionally accompanied with some degree of chronic inflammation.

About this time, Gendrin, in his *Philosophical Treatise of Practical Medicine*, tom. iii, p. 23, observes that the mucus which is deposited and agglutinated on the gastro-intestinal mucous membrane sometimes forms concretions of a pseudo-membranous appearance, which are excreted with the stools, and the same that one meets with in the intestines after death. These concretions, he says, never present the least trace of organisation, which authorises us to distinguish them from organised pseudo-membranous concretions which are formed on inflamed surfaces.

In order of time, we now come to, perhaps, the most perfect record of this disease in any language, with one exception, shortly to be mentioned. I allude to the paper contributed by Dr. Todd in the *Cyclopædia of Practical Medicine*, on Follicular Duodenal and Follicular Colonic Dyspepsia. Dr. Todd, as it appears from his titles, looked upon the disease as a form of dyspepsia, and he attributed it to a morbid condition of the mucous follicles.

Dr. James Simpson, in 1846, stated, that from observations he had made, he was convinced that this disease, which he termed chronic pellicular intestinal inflammation, and also eruptive inflammation of the intestinal mucous membrane, was often met with in practice. He considered the passing of shreds or membranes as a characteristic symptom, and that these substances as shown by the microscope consisted of plastic or coagulated lymph.

* Read before the Medical Section at the Annual Meeting of the British Medical Association, in Newcastle-upon-Tyne, August 1870.

Andral also makes mention of these membrani-form productions.

Graves considered the cause of these cases to be irritation of the mucous membrane of the large intestine, and of the rectum in particular.

Mr. Iliff of Newington read a paper in 1844, before the Physical Society of Guy's Hospital, on the membranous-like formations which are not unfrequently passed by patients, mingled with alvine evacuations. He read the particulars of eight or nine cases, one of which was the subject of malignant disease of the os uteri. His paper was illustrated by drawings of the membranes; and I am under a debt of gratitude to Mr. Iliff for the loan of them.

Cruveilhier acknowledges how frequently he had opportunities of observing gelatiniform pseudo-membranes voided by stool after the ingestion of energetic drastic purgatives, and the bark of pomegranate administered by physicians, who took for the remains of worms the mucous products expelled by their patients. He also refers to Villermé, who had witnessed many times the expulsion of these false membranes in individuals affected by the colic of Madrid.

Grantham, in his *Facts in Medicine and Surgery*, describes this disease under the title of Fibrinous Diarrhoea, and states that in every case which he witnessed, he invariably found the disease produced by the administration of mercury, conjoined with the too frequent use of aperient medicine. He also found that such patients experienced an exacerbation of symptoms by the use of mercury, even in small doses.

In the year 1854, M. Broca showed to the Anatomical Society of Paris, membranes which were frequently passed by a woman, and, curiously enough, by her brother-in-law; by the latter for ten, and by the former for fifteen years.

Dr. Potain also showed to the same Society, in 1857, shreds of gelatinous mucous matter expelled during defæcation by a young epileptic girl.

Dr. Lionel Beale, in his *Microscope in Medicine*, states that "mucous casts are sometimes expelled from the large intestine, and occasionally as complete tubes. Flakes, some of which are very firm, are common enough, especially after prolonged constipation. They consist of a firm mucus, in which the epithelial cells from the large bowel, and many corpuscles are embedded." He refers to an interesting example sent to him by Dr. Borrett. The patient was a little girl aged four years.

Trousseau, in the second volume of his *Clinical Medicine*, says that in many cases we confound the pains which have their seat in the transverse colon with stomach pains, and that it is not exaggerating the fact to say that, in a moiety of cases, what we are in the habit of calling gastralgia is nothing but colalgia. The pains experienced by the patient coincide with an obstinate constipation, sometimes followed by diarrhoea, which is accompanied by an excretion of more or less abundant thick mucus. The mucus presents itself under the form of bands of white ribbons, that are compared to macaroni.

In the year 1857, Mr. Jonathan Hutchinson brought before the Pathological Society of London, the case of a lady who voided what he termed tubular exudation casts of the intestine; and also the parallel case of a pale cachectic man, aged 23, who for at least three years, in spite of all treatment, continued to void these casts. Specimens of the matters passed by the lady just referred to, were submitted to Dr. Andrew Clark for examination. As the conclusions arrived at by Dr. A. Clark, after a chemical and microscopical examination of a most exhaustive nature, will frequently be referred to hereafter, I will not anticipate myself by giving you them at present. I shall also have frequent occasion to draw from his equally valuable views on the same subject.

Dr. Guyot brought a case of mucous disease before the Medical Society of Hospitals, at their sitting on the 28th February, 1848. Dr. Siredey, at a sitting on the 11th December of the same year, brought forward a case of a similar character. The paper read by Siredey is full of information and interest.

M. Grisolle, in his *Traité de Pathologie Interne*, in a chapter headed Chronic Enteralgia, states that the disease may continue whole years with remissions more or less long and complete. The patients of whom he speaks experience a kind of habitual uneasiness in the belly. After each crisis, a very abundant excretion of a white or yellow albuminous fibrous matter, resembling in form and colour large vermicelli. During the paroxysms of pain, the pulse remains natural. It is also, he states, more frequent with women than men.

M. Worms drew the attention of the Anatomical Society of Paris to some membrani-form flakes voided by a person of advanced age.

Dr. Merland, of Chaillé, in the *Union Médicale*, refers to about thirty cases of this disease which came under his notice.

Axenfeld has noticed and reported cases of expulsion from the anus of gelatiniform white ribbons, several inches long, which he considered to be formed by some concrete mucus.

It will now be my object to elucidate from these several quotations and allusions the borne-out facts recorded, and also to give a general description of the disease.

[To be continued.]

ALCOHOLIC POISONING.*

By G. S. ELLISTON, Esq.,
House-Surgeon to the East Suffolk Hospital.

I PROPOSE briefly to lay before you a few cases of alcoholic poisoning which I have met with in hospital practice during the last four years, the treatment which I have adopted, and the care that should be taken to correctly diagnose these cases from apoplexy and injuries of the brain, which they much resemble. Unfortunately, the only history you generally get is from the policeman who brings in the case; and invariably all that he can tell you is, that he found the patient insensible in the street—he supposes, from drink.

The first case is one which I had when one of the resident surgeons at the Royal Free Hospital. A law-writer, about thirty-five years of age, was picked up by the police in Gray's Inn at midnight, apparently dead, and conveyed to the hospital on a stretcher. I placed him on a couch, and found the body cold and pulseless, the pupils contracted, the conjunctivæ quite insensible to touch, and respiration scarcely perceptible. I at once applied a very powerful galvanic battery always kept in readiness, placing one pole on the nape of the neck, and the other over the course of the phrenic nerve. For two or three minutes he did not appear to feel it; but the respiration slightly improved. I varied the galvanism by placing the pole over the pit of the stomach, and sometimes on the nose; here it seemed to have the greatest effect, causing it to twitch violently; and shortly afterwards the head began feebly to rise from the pillow. This was followed by a gasp; and, by still applying the pole to the nose, I was able to bring the body up into the sitting posture; but, directly it was removed, he fell back a dead weight. I continued the galvanism; and in a few minutes he was so far roused as to say, rather thickly, "Take it away; don't burn my nose." At first, he was inclined to relapse; but a few touches of the battery soon brought him round sufficiently to take an emetic. But very little came from the stomach, and the odour of spirits was slight. He was placed in bed, and three or four days afterwards left the hospital quite well.

The next case shows that the serious symptoms of alcoholic poisoning may be caused by injuries received whilst partially under the influence of drink; and the surgeon may easily be misled by the statements of those accompanying the patient. Two nights after the preceding case, a gentleman was brought in by the police, picked up in the street in an insensible state. He appeared to have been drinking, for his clothes were in a disordered state; and he had evidently been robbed of his watch and other valuables. He was galvanised, and was soon sufficiently sensible to tell his name and address. As the hospital was quite full, it was suggested that the police should take him to his home. On his way there, he became much worse, and was taken to the nearest infirmary, where he died during the night. The *post mortem* examination disclosed a ruptured liver, which the subsequent inquiry proved to have been caused by a cab knocking him down. In this case, there was no clue to the injury, as not one of those who brought him in witnessed the accident.

Since being at the Ipswich Hospital, I have met with two or three well marked cases of alcoholic poisoning. A sailor was brought in one night by the police, who found him lying insensible in the street about a hundred yards from the public-house at which he had been drinking. A medical man had been called to him; but, as he failed to rouse him, he ordered him into the hospital. He was then in a profound state of coma, with all the symptoms which I have before described. I at once applied galvanism; and in about five minutes he began to show signs of animation, quickly followed by pugnacious attempts to knock everybody down near him, more especially the operator. A few more touches made him perfectly sensible; and he took an emetic as quietly as possible, but very little came from the stomach. He shortly afterwards fell into a quiet sleep, and got up next morning feeling as well as ever. Nothing could persuade him that he had been in any danger; and he left the hospital early, treating the whole affair as a joke.

A very similar case happened a few days ago, at the Whit-Monday fête. A young man was found in the afternoon on the ground in an insensible state. He was supposed to be suffering from sun-stroke; and, while a medical man who was present was trying to restore him, a bystander said he believed it was the result of some gin he had seen

* Read before the East Anglian Branch.