been well destroyed. There was no bleeding. On the 19th, the sloughs, which were very deep, had nearly come away. During the process of separation, the odour had been very foetid. The edges of the wound were beginning to be covered with mucous membrane. On the 23rd, she had improved rapidly; the wound was greatly cicatrised. A few dubious points remained here and there. On December 2nd, catamenia had appeared; they were much less copious than usual, and more like oozing: no clots. On examining her to-day, it was found that there was a large hole superiorly in the os and to the left side, which bled on touch. The fibroid enlargement was in no way diminished. A pledget of cotton, dipped in liquor ferri persulphatis, was placed within the os. On the 7th, there had been no more bleeding.

This patient made very little progress afterwards. Her mind was greatly distressed by the unkindness and desertion of her husband and daughter. She was removed into Middlesex Hospital, where, I learnt, she died some months afterwards.

[To be continued.]

NOTES ON APOMORPHIA.

By F. M. PIERCE, M.D., L.R.C.P.L.

IN the present communication, my object is to draw attention to the valuable properties possessed by apomorphia. By the kindness of my friend Mr. C. R. Wright, B.Sc., I was enabled, soon after the discovery of apomorphia by Messrs. Matthiessen and Wright, to make some investigations into its value in practical medicine. A detailed account of apomorphia will be found in the *Proceedings of the Royal Society for 1869*. It may suffice here to state that apomorphia is obtained by subjecting morphia to the continued action of pure hydrochloric acid at a high temperature for several hours. The base can be obtained from the resulting hydrochlorate of apomorphia, by dissolving in water, adding excess of bicarbonate of sodium, and extracting with ether or chloroform. In composition, apomorphia is morphia, minus an atom of water. In its chemical reaction, it is in nearly all respects different from morphia. It is soluble in cold, and to a greater extent in warm, water. In powder, it is of a snow-white colour. The watery solution is colourless at first, rapidly changing to a dark olive-green, and, at the end of a few weeks, almost black. The cause of this evident decomposition is at present *sub judice*. Morphia is not reproduced. The physiological effects of the coloured solution are the same, to a slightly diminished extent, as of the fresh preparation, but rather more irritating to the skin. An increase of the dose is, therefore, required on repetition.

Briefly to indicate its therapeutic value, the chief character of the drug—its power of producing vomiting—was noticed to a very disagreeable extent during its preparation. Its absorption through the cutaneous surface produced in the manipulators lassitude, weakness, frequent headache, constant nausea, and occasional sudden attacks of vomiting. From some experiments made by Dr. Gee, one-tenth of a grain was considered the average dose for an adult. I have found this too large, whether of a fresh or an old solution. The desired effect is readily obtained with one-fifteenth of a grain, or less; vomiting occurring in from five to twelve minutes, and the subsequent depression being slight and of short duration.

The solution which I have generally used consisted of a grain of hydrochlorate of apomorphia (the base being very unstable), dissolved in two hundred minims of water. Five minims of this preparation, or one-fortieth of a grain, was the dose generally administered to children, to whom I have frequently given it as an emetic. In nearly all the cases, it was administered subcutaneously. In private practice, the hypodermic use of drugs may appear a little formidable; but the more complete control of a drug thus acquired will probably in time outbalance the objections to this method.

In illustration of the action of apomorphia, I append one case.

W. B., aged 8, residing at Kingston-on-Thames, came under my notice suffering from well marked chorea. He was an intelligent, well-made boy, of sanguine temperament, the youngest of three. His father died of some strumous affection. The mother works much at her needle; she is nervous, but healthy. The patient's symptoms followed a severe scald of the chest when five years of age. At first, he was simply considered fidgety. Four months before I saw him, he had rheumatic fever. The twitches had increased very much since. The parts most affected were the muscles of the face, especially of the mouth (he could not protrude the tongue readily, nor keep it still when protruded), the muscles of the neck, arms, and trunk. The legs were least affected. A slight systolic *bruit* was audible at the apex of the heart. He had no worms. Three days after I first saw him, I injected

one-fortieth of a grain of hydrochlorate of apomorphia into the left arm. The pulse before the injection was 70, regular, weak; temperature and pupils normal. He did not complain of approaching nausea, but suddenly vomited at the expiration of six minutes and a half; and, fifteen minutes later, he again vomited. The pulse, nine minutes after the first vomit, was 100, weak; the temperature was not affected to any noteworthy extent. He felt a very slight drowsiness, lasting only a few minutes. He did not sleep, but returned to play. The involuntary movements were markedly better next day; they were less sudden, frequent, and complete. He could stand for two or three minutes almost quiet. At the end of three days, during which period his improvement was maintained, the fortieth of a grain of apomorphia was again injected. Previously to injection, the pulse was noted as being 76, irregular, weak; the temperature 98.9; the pupils normal. The muscles of the neck and left arm were still very much affected. Vomiting ensued, without premonitory symptoms, in fifteen minutes. The pulse was 88; the temperature-variation unimportant. The pupils were slightly dilated. He did not feel any sleepiness, and, after a few minutes' rest, felt quite well. The chorea was much diminished after this; and, ten days later, I injected one-thirty-third of a grain in the arm. The pulse was previously noted 80, regular; pupils normal; etc. Vomiting occurred suddenly at the end of ten minutes; the pulse counting 90; the pupils dilated a little. He was discharged quite recovered soon afterwards, not having had any choreic movements for the last eight or nine days. No other treatment was adopted.

In two other cases of chorea, the action of apomorphia was beneficial, though not as distinctly as in the above case, and, indeed, not more so than many of the vaunted remedies for chorea. It does not seem to have any specific action. It is only as an emetic that I would draw attention to it. With this object, I have used it at the commencement of pneumonia; to relieve the throat of exudation in diphtheria; in scarlatina, etc.; in cases of poisoning; and in drunkenness. As an emetic, it is depressant, but not to a dangerous extent. Though, in the example given, slight drowsiness was observed on one occasion after its use, it has not been a constant accompaniment, but the exception. The feebler the preparation, the less was this noticed. In both adults and children, with a dose not exceeding one-fifteenth of a grain, very slight inconvenience followed. The alterations of the temperature and respiration were so slight that they may be disregarded. The pulse was not peculiarly affected. The pupils were generally a little dilated. No action on the bowels or kidneys was observed.

Apomorphia is peculiar in the sudden accession of the vomit. The usual premonitory signs and symptoms are wanting, or all but coincident with the expulsive effort. The first attack of vomiting is sometimes succeeded by another, after a few minutes' interval, the patient quickly recovering from the temporary disturbance. Most frequently, the single vomit is short and final.

The occasional occurrence of sudden vomiting after the administration of morphia, subcutaneously or by mouth, may perhaps be due to the presence of this body—apomorphia—as a product of partial decomposition. Should further investigation complete the chain of chemical analysis and synthesis of the morphia salts, some very instructive chemical and physiological results may be obtained.

In conclusion: as an emetic, apomorphia is pre-eminent in the smallness of the dose required; the certainty, rapidity, and completeness of its action; the unimportance of its baneful effects; and its non-irritating character. It contrasts most favourably with the old-fashioned emetics. The only drawback at present is its costliness; but, no doubt, in a little time it will, as well as chloral, come within the reach of the most economical practitioner.

CASE OF A NAIL PASSING SAFELY THROUGH THE ALIMENTARY CANAL OF A BABY.

By B. BLOWER, Esq., Liverpool.

SOME years ago, a woman brought her baby in a state of suffocation. Passing my fingers into the fauces, I felt a piece of iron right across. A slight touch dislodged it; and, to my great dismay, it went down. The child immediately recovered its breath, and was soon at the breast. I desired the mother to give nothing but the breast-milk, and to watch carefully the motions. The next day, the poor woman came with the nail which I have now before me. It is two inches long, and a quarter of an inch broad at the head; the point is rather blunt. It is interesting to follow this intruder through its windings along the convolutions, as it wended its quiet way undisturbed by emetic or purgatives. I send this case in corroboration of the remarks of Mr. Newstead in the *JOURNAL* of February 5th, and to show what Nature can and will do if not disturbed in her operations.