at the same time as the chloral, the effect of the latter was less marked. On several occasions when I tried to give her a dose in milk, it seemed to disagree, and on one occasion she vomited it up again; the only time she was sick. The subcutaneous injection did not answer, the quantity to be injected being too large.

On another occasion I should begin with a drachm dose in sugar and water every half-hour till sleep ensued, and then continue drachm doses by the mouth every twelve hours (or half a drachm every six hours) regularly. If it be administered by the rectum double the above dose, at least, must be given. A note of ten other patients on whom the influence of chloral was tried will be found at p. 437 of this JOURNAL, Oct. 23, 1869.

THE THERAPEUTIC USES OF CHLORAL.

By STEPHEN MONCKTON, M.D.,
Physician to the West Kent Hospital, Maidstone.

SEVERAL short papers on the subject of chloral have appeared in the different journals. I would here give an account of my little experience.

I have given perhaps fifty doses, among about ten different patients, in the last three months—often with remarkable success; sometimes with partial effect; never with mischievous results, beyond a little delirium lasting an hour or two, resembling that of the second stage of chloroform, and proving (though not pleasant for the time to those around the patient) very transient and innocuous.

The drug, being pungent and unpleasant to take, should be freely diluted. Not less than an ounce and a half of water should be the vehicle for every dose of twenty grains or upwards. A little sugar or syrup is also desirable. Though disagreeable, the draught has never caused vomiting, over which, if already existing, it seems to exercise a remarkable control. When it is swallowed, the effect is generally speedy. It is well, therefore, that the patient should not take it till actually in bed. Less than ten minutes will sometimes suffice to send him fast asleep. One man (in the hospital with subacute rheumatism) became delirious—*i. e.*, noisy, irrational, and rather cheerfully excited—after a twenty-five-grain dose; and twice in the female wards the good effect has been very partial, perhaps from the light and movements surrounding the patient through the night. This suggested to me the practice of bespeaking darkness and quiet for every patient for whom a night-dose is ordered.

The soporific effect is sufficiently prolonged; five, eight, and ten hours' sleep, with generally a short break and semi-consciousness in the middle, being common enough; while a decided but not unpleasant sleepiness sometimes pervades the following day.

In the short series of cases on which alone I profess to base this paper, it has not once happened that headache, vomiting, or fever, harassed the patient on waking up. On the contrary, it has seemed to me clear that chloral, in doses of from twenty-five to fifty grains, is a very effective anodyne, antispasmodic, and hypnotic; that it possesses many recommendatory qualities and few drawbacks; while some of its partial failures are doubtless due to surrounding interferences and insufficiency of dose. It is very certain that twenty or twenty-five grains will sometimes give a strong man a wakeful and excited night, when forty or fifty grains would have secured sound sleep.

The following three cases, of different types, seem worth giving very shortly in detail.

CASE I.—Mr. C., a grocer, aged 76, came in October last with symptoms which led me to seek and find a recent left inguinal rupture, of only a few days' standing. He was recommended to procure a truss, and returned no more. Three months afterwards, his daughter sent an urgent summons at 7 A.M.; and I found him in the agonies of intestinal obstruction, symptoms having existed about twelve hours. No external hernia could be discovered; in fact, it had never come down since the first application of the truss. But there was a hard, globular, painful tumour, between the internal abdominal ring and the umbilicus on the left side. His pain was great, vomiting incessant, and belly rather distended. I ordered half a drachm of chlorodyne immediately, half as much to be repeated every two hours; and fomentations without measure. Twelve hours afterwards (7 P.M.), he was worse; the pain, sickness, and restlessness were most distressing. By care, he sipped down and retained forty-five grains of hydrate of chloral in an ounce and a half of water, with a few grains of sugar. At 10 the next morning, he had had a quiet night, with several hours' sleep; and had only retched once. The abdominal condition was unchanged. His countenance was good; his tongue moist; his head comfortable. Hesitating to credit the chloral with such power for good, I assumed that a quiet resolution of the intestinal difficulty had

occurred. But, in a few hours, pain and vomiting returned; the abdomen was more swollen; and the ejecta became distinctly stercoraceous. So it was proved that, in spite of unrelieved obstruction, the chloral had commanded quietude when strong opiates failed, without inducing that parched, stupefied, and miserable state which opium brings about, and never more surely than in the case of old people unaccustomed to its influence. On each of the next three nights, a fifty-grain dose of chloral was given, with satisfactory effect. Soon a liquid trickling *per anum* set in, passing gradually into natural defecation. Within a fortnight, the old man had fairly recovered.

CASE II.—Mr. D., an acute and sprightly old gentleman of 72, had been going wrong for some weeks with heart-symptoms, probably dilatation of the right cavities. There were dyspnoea, tumultuous action, no *bruit*, feeble and very irregular pulse, with some cough and expectoration. By February 1 his countenance had become rather livid, his dyspnoea urgent, and his legs swollen to the knees. After several nights of short and broken sleeps, from which he awoke in distress, his daughter begged permission to give him some opiate pills then in the house. Instead of them, he took thirty grains of chloral at 10 P.M. He was asleep in five minutes, roused and spoke to his daughter at 3 A.M., then slept again till 10 A.M. without disturbance, stupor, or dyspnoea. I found him comfortable, but sleepy, at 10.30; unwilling to remain in bed because of a mutton-chop that was waiting on the breakfast-table. He has taken a few more doses, and is in greatly better condition. Although the slumber was long and sound, he quite recollected waking and talking at 3 A.M. The sleep was therefore non-comatose in character.

CASE III.—Miss H., a tall slender girl, aged 20, with a large cavity at the right apex, in fact dying consumptive, was much worn with night-cough and nervous insomnia, expectoration not very profuse. She took thirty grains of chloral a month ago at bedtime, and slept more than eight hours, without being at all overwhelmed. She then rose to breakfast, and slept afterwards for the greater part of the day on a sofa. Since then she has taken eagerly the same dose four or five times a week with the happiest effect; and no constipation, headache, or impeded expectoration has ever appeared to follow.

Possibly, this homely record of my little experience may induce others in hard practice to lay aside natural and honourable misgivings and try this medicine for themselves. I shall watch for opportunities of giving it in acute rheumatism, chorea, neuralgia, spasmodic asthma, anæmic insomnia, and even inflammation.

CASE OF TRAUMATIC TETANUS SUCCESSFULLY TREATED BY BROMIDE OF POTASSIUM AND HYDRATE OF CHLORAL, ETC.

By EDWARD R. DENTON, M.R.C.S.Eng., L.S.A., Leicester.

GEORGE B., aged 18, fell from a cart January 31st, 1870. I was called in on the 16th February, six days after the first appearance of the tetanic symptoms. The patient was of the sanguine temperament, rather spare, short in stature, and of temperate habit. On examination, I found that he was unable to open his mouth or protrude his tongue; the lips were elevated from the teeth and gums, causing the characteristic "tetanic grin." The head was thrown back, and opisthotonos was well marked; the abdominal muscles being excessively tense, and the back arched and raised from the level of the bed. He was able to swallow only in very small quantity, even then appearing to suffer much from the spasm of the muscles of deglutition. The face was flushed and the skin hot; pupils dilated. The pulse was 120, small, hard, and wiry. Respiration 35. The bowels had been constipated for two or three days, and he had not slept for five or six nights.

The patient seemed to be aware of his precarious condition, but, as well as he could, expressed his readiness to die if nothing could be done for his relief. The wound presented a dry and unhealthy appearance, but was limited to the dorsal surface of the first phalanx of the second finger of the right hand. The nail had been removed three days previously, the back of the hand was swollen and inflamed, and an occasional shooting pain was felt extending along the posterior surface of the fore-arm and the anterior surface of the arm into the axilla.

The bowels were first relieved by a calomel and jalapine purge followed by a stimulant enema. Morphia, nepenthe, and chlorodyne, were administered every night from February 16th to 21st, with little or no effect in procuring sleep or relief from pain. I then procured some of the syrup of hydrate of chloral, of which small (three drachms) but repeated doses were given till sleep was procured, which at first lasted only a few hours. For the first three days ice-bags were placed in con-