LECTURE

ON

THE ACTION OF THE CHOLERA-POISON ON THE BODY; AND ITS NATURE AND HISTORY OUTSIDE THE BODY.*

By JOHN MURRAY, M.D.,

Inspector-General of Hospitals, Bengal Medical Department.

GENTLEMEN,—I have been requested by several of my professional brethren, whose opinion I highly value, to deliver a Lecture on the Nature of the Cholera-Poison, and its action on the human body, as a supplement or key to my practical Report on the treatment of the disease, already in your hands. It may possibly be asked why these views were not given as a preliminary to those on the treatment of the disease. My answer is, that the inquiry then conducted was limited by the orders of Government to practical points, and as the views I now offer must, to a certain extent, be theoretical, they may be wrong; and, whilst the facts I advanced are correct, the weakness or fallacy of my reasoning on theoretical points might have been reflected on remedies that are valuable. It is possible that similar objections may be made to the limited nature of the present lecture, in reference to the restricted views which I now propose to submit to your consideration; but the subject is too comprehensive to be treated satisfactorily on one occasion, and other branches of it may be left for future discussion. One great object I had in view in consenting to deliver this lecture was to derive benefit from the criticism and opinions of so experienced and talented an assembly, who have resided long in what is called in Europe the cradle of the disease; more especially as on some points our views do not coincide.

In no disease are there more marked and characteristic symptoms, during one of its stages, than in cholera; but these specific symptoms, which are only found in one stage, are so dissimilar to those induced by the poison in an earlier stage, or milder form, that the latter have by many been overlooked. The value and importance of the early detection of the presence of the poison cannot be over-estimated; as we all know from experience that, after a certain stage, the vital organs are for a time little sensible to the action of remedies. They are paralysed by the presence of the poison, in the same manner as the voluntary organs are paralysed by the action of chloroform. It is in preventing the super-vention of this stage, that most life has been saved; and the physician who can soonest detect the presence of the poison, will be the most successful in his profession.

The best method of ascertaining the symptoms induced by the presence of the poison, is to mark its action in its most intense form, before the other symptoms appear which follow the retention of those effete portions of the blood which, for the preservation of health, require constantly to be eliminated by the secreting organs. This intense form of the disease is mentioned by most writers, but is rarely seen in hospitals; in fact, it can only be observed by those who are searching for it among the people, as it proves fatal in from two to four hours. The patient falls down as if struck by a sudden shock, livid, cold, covered with clammy perspiration, with a broken voice and sinking pulse. The respiration becomes slow and intermitting; the only complaint is of uneasiness, burning and sinking in the epigastrium; the intellect is clear; there is generally an evacuation of the ordinary contents of the bowels, and sometimes one or two of the peculiar congee-like discharges; but this ceases, the pulse disappears, intelligence cannot be roused, and life is at an end.

In the more ordinary form of the disease the patient is purged, generally in the early morning; the evacuations are at first natural, copious, and liquid, with a little tormina, and uneasy feeling in the epigastrium, approaching to heart-burn; the body is cool, the pulse quiet, and the intellect clear; but the countenance is dark, the eyes congested, the nails look blue, and there are occasionally flying cramps. This is soon followed by the vomiting of undigested food, eaten perhaps one or two days before—then by copious congee-like evacuations from the bowels and stomach; intense burning pain in the epigastrium, low, sinking feeling, cold clammy perspiration, sometimes very profuse, feeble

pulse, livid characteristic countenance, and cold breath. There is no secretion of bile or of urine; the only gland that sometimes continues to act is the mammary in nursing mothers. There are generally cramps ascending from the toes and fingers towards the body or diaphragm; the higher they ascend the greater the danger implied; but they are often absent, and generally disappear in the worst cases for some time before death. In these cases the vomiting and purging also cease; restless moving of the body comes on, whilst the patient lies with the eyes sunk, half-closed, congested, and glazed; semi-conscious and occasionally wandering; when roused, he still complains of the sinking burning pain in the epigastrium, and calls for water. The pulse is not perceptible at the wrist, and there are long intervals in the breathing. It is wonderful how long this moribund state will sometimes last, before death closes the scene; and still more wonderful that in a few cases successful reaction takes place. There is great tendency to syncope or fainting, in collapse, and even after reaction has made considerable progress. The simple act of assuming the erect posture, or even sitting up in bed, has in many instances been followed by the patient falling back in a faint, from which he never rallied, or merely struggled on a little longer before death; in these cases, coagula are found in the heart after death. In most cases there is an effort at reaction. When it takes place the burning pain in the epigastrium disappears, the restlessness subsides, the stools become coloured, warmth returns to the palms of the hands, the colour improves, the pulse becomes stronger, and sleep ensues. If, after this, the urine appears, a favourable prognosis may be given, provided no strong remedies have been administered during the previous stage, and that collapse has not lasted many hours. In such a case there are no marked sequelæ for the disease. I have in several instances seen the first evacuation after sleep, natural in colour and feculent; but in general, especially when the collapse has been prolonged, the evacuations are first dark, sometimes pitchy, then green and yellow, and there is looseness for some time, and occasionally irritability of stomach. These symptoms are not essential, and may frequently be traced to the remedies employed. The sequelæ which follow collapse depend on different causes, either natural or artificial. The natural causes are imperfect reaction, when the secretion of the kidneys is not restored, when the liver acts imperfectly, and when uneasiness continues in the stomach with want of appetite. The pulse rises and becomes quick, the skin dry, the tongue coated, with sordes on the teeth; there are head-ache and wandering, sometimes delirium. Looseness continues, sometimes extending to bloody stools with tormina and tenesmus. Long protracted collapse, even though the reaction be satisfactory, induces slow convalescence. The secretions from the liver and kidneys are morbid, and the digestion imperfect. The accumulation of effete particles in the blood, caused by the suspension of their excretion during the protracted collapse, may have injuriously affected the brain or other organs. This has to be worked off. Rest, gentle nursing, and time are required to allow the feeble organs to restore the blood to its healthy state. The artificial causes of these sequelæ are those remedies which would directly induce inflammation of the stomach, intestines, or brain, or indirectly suppress the secretion, or diminish the action of the kidneys, liver or intestines, by which the specific poison and the retained effete portions of the blood are eliminated from the system. The two remedies which, when given in excess, are likely to act most unfavourably in this way are brandy and opium. The opinion of the practical profession in India, on the exhibition of these remedies in collapse, in large quantities, is almost unanimous that they are not beneficial, and in general that they are decidedly injurious. Such, gentlemen, is my opinion, and I would it had influence to abolish a practice which has prevented many from surviving, who would otherwise have recovered from the attack of this terrible disease. But this is not the time to discuss the treatment of the disease.

We shall now consider one of the most frequent forms in which cholera appears; viz., premonitory diarrhœa, called by many "choleraic diarrhœa," and by some considered a distinct disease. We cannot over-estimate the value of correct information on this point. In the first place, it is a stage in which the system is still amenable to remedies, and the progress of the disease can generally be arrested. In the second place, the poison may be in process of elimination through these discharges, and the disease unwittingly disseminated over the country—an evil which might be prevented by a correct diagnosis. Again, if the physician tell the patient that his disease is not cholera, should it advance to the next stage, the patient believes he has no chance of recovery; and at this critical period his heart sinks, and the despondency which ensues redoubles the power of the poison, and diminishes the hope of success.

Diarrhœa is always very prevalent during cholera epidemics. It is painless; the evacuations are light coloured, copious, creamy, gradually becoming more watery and devoid of feculent smell, with occasionally

* Delivered at a meeting of the Bengal Branch of the British Medical Association, on January 19th, 1870.

the peculiar smell of cholera perceptible. They may then become watery and colourless—the characteristic stool of cholera—with collapse and cramps. The period occupied by these changes varies from a few hours to several days. At first there is no appreciable difference between these evacuations and those passed in ordinary diarrhoea, which may accompany disease of the liver or of the kidney, or be induced by eating unripe fruit, or indigestible food, which forms one of the most frequent exciting causes in cholera.

There are other appreciable signs of the presence of the poison at this period. There are numbness, or want of power, in the rectum, a want of appetite, a desire for stimulants, and sometimes uneasiness in the stomach or heart-burn, and occasionally flying cramps. The secretion of urine is scanty, the countenance dark, and the eyes congested. The frequency of the stools may be diminished, and the consistence of them improved by opiates and astringents, without improving the colour, removing the tendency to recurrence, or curing the disease.

Diarrhoea is the first active symptom induced by the presence of the poison in the system; but, previously to this, an experienced physician will detect its presence in what I have called the stage of "*malaise*"—a feeling of depression of spirits, and want of appetite, with weight in the head, torpor of the bowels, scanty urine, and a want of brightness in the colour of the face. I have several times observed this state to disappear suddenly after an inordinate flow of urine.

These are the earliest symptoms of the presence of the poison which I have been able to trace; but in many instances it appears to be present without exciting observable symptoms, if we may judge from the fact of individuals leaving an infected locality, apparently in perfect health, being attacked on the following or subsequent days in a distant place, where the disease had not previously existed. That the *malaise* which I have described is caused by the presence of the poison of cholera is an opinion strengthened by the fact, that during an epidemic attack the action of a purgative, especially salts, will almost always be followed by collapse, cramps, and death.

We have now traced the action of the poison on the system from *malaise* to collapse, through diminished power of the sympathetic nerve on the functions of the organs of organic life, inducing deficient secretion, or total suspension of excretion through complete paralysis. This diminished action produces want of appetite, with defective supply of gastric juice, the absence of which allows the natural changes of fermentation and putrefaction to take place in the food, rendering it irritating, and inducing diarrhoea. In ordinary circumstances, this is Nature's cure for improper food; but at this time there is also present in the system a foreign body, which I shall now endeavour to describe. It may not be visible, but its nature can be traced by its effects. It is evidently vital, as it reproduces itself. It is infinitely small, like other lower vitalities, which produce fungi in fermentation, and vibriones, etc., in putrefaction. There are known chemical agents, called antiseptics, by which fermentation and putrefaction are suspended or restrained. It may, therefore, be inferred that there are agents by which this other vitality can also be restrained or destroyed. Nature provides a powerful agent of this character, viz., the *gastric juice*; and it may naturally be inferred that it has a similar power over this analogous low vitality. The whole history of the progress of the disease tends towards this conclusion, and affords an explanation of the cause that induces trifling depression, which may be developed in a few hours into a violent collapse. In *malaise*, the presence of the poison in the system is evident, but it frequently passes off without appreciable injury. The germ of the disease in this case is restrained in its reproduction by the presence of the gastric juice; but if, during the presence of the poison, the secretion of the gastric juice be diminished or suspended by exhaustion, mental anxiety, terror, or grief; or if, along with the bile, it be removed from the bowels by a purgative, the restraining power being absent, the germ is allowed to multiply with its ordinary rapidity, and from tens becomes hundreds of thousands. That the germ can multiply in this excessive number, is known from large bodies of water becoming contaminated in a very short period. This sudden development from the removal of the restraining power, affords an explanation of what long formed a difficulty in my mind—how so virulent a poison exerts so little influence as it does in one stage. That the gastric juice is absent in the most virulent form of the disease, is evident from the fact that food eaten one or two days before is frequently vomited, or found undigested in the stomach after death. There is another fact in the history of the disease, which would be elucidated by this explanation: the most frequent hour of attack is in the early morning, about 3 or 4 A.M., when, the process of digestion being long completed, the secretion of the gastric juice is for a time suspended. It is easy to imagine that, if a small number of germs could diminish the secretion, which in its turn restrained their growth, and if the natural multiplication of these germs were allowed for a short period, they would be

present in such numbers as completely to prevent the future secretion of the restraining power. The existence of such a specific germ is deduced from the production of all the effects which such a body, and only a living and reproducing body, could cause. That it is visible, or appreciable by the optic nerve, with the assistance of the powerful microscopes of the present day, is the conviction of many. Hallier thinks he has detected it in the shape of a fungus, and Parkes saw a multitude of animalcules in cholera-stools. The most searching investigation is now being made in Calcutta, by Drs. Cunningham and Lewis, on this point. The presence of lively animalcules in thousands, in cholera-evacuations, is constant, and might lead to the inference of their being the cholera-germ. The course of their existence passes through one stage, in which it has the form described by Hallier. It is intimately associated with the cholera-germ, if it be not the real one. "*Noscitur a sociis.*" My doubts of its being so rest on the fact, that a similar animalcule is found in every tank and drain in Calcutta, and in the evacuations in other diseases, though in comparatively trifling numbers. At the same time, I do not consider that the individuality of the germ depends on its being visible. There are many agents that cannot be seen that are appreciable by the olfactory nerve. Gases or vapours, that are invisible, yet may cause death, or lingering illness.

For practical purposes we have gained an important point. It may reasonably be inferred that this animalcule or infusorium thrives best where the cholera-germ is present, and that whatever acts on it will act in a similar manner on the cholera-germ. A series of most interesting experiments have been begun to investigate this point. All ordinary antiseptics have been used, and also all the remedies recommended in cholera. Their action was uniform, and the result curious, and likely to lead to valuable consequences. In the first instance, the remedies were applied to the animalcule in its lively state, and the effect was watched under the microscope. In some cases it was sudden death; in others, the motion slowly but certainly ceased; in others, they turned round about for some time and then ceased to move; whilst with other supposed valuable remedies they continued lively and unaffected. In estimating the value of these experiments, it must be borne in mind that they were conducted in glass tubes; whereas in the human body they have to be conducted in the stomach and intestines. The effect of these agents was very curious, and watched with great interest. The gastric juice was first used in the shape of *pepsine*, a neutral, or rather alkaline preparation. For some time it had apparently no effect, but gradually the motions diminished, though they had not entirely ceased in half an hour. The most interesting experiment was the addition of a very small quantity of acid to the pepsine, which renders it more like the natural gastric juice. The suspension of animation was instantaneous, more rapid than from any of the other agents that were employed; the outer coat of the corpuscle was then dissolved. When bile was diluted and applied, the motions gradually subsided; and when it was applied undiluted, the cessation was immediate; but there was no solution of the germ, as under the action of pepsine; in fact, after an interval of three days, the lively action returned. The effect of the cholera pill was curious; for some minutes there was little change, but gradually the liveliness subsided, till in from five to eight minutes only one or two continued in motion, and in ten minutes all were quiescent. It was here that I noticed the whirling round before death. The bodies remained in a quiescent state; but were not dissolved as under pepsine, or under acetic acid, which was the next remedy in rapidity and power. Carbolic acid, when strong enough, acted instantaneously in the glass tube; but it is too strong to put into the human stomach with impunity. Sulphuric acid had no apparent action unless combined with pepsine. In administering these antiseptic remedies to the human subject, their influence on the secretion of Nature's antiseptics—gastric juice and bile—must be considered. If they diminish these secretions, it is questionable if the substitute would prove more powerful in arresting the development of the germ. Even remedies which have no direct action on the germ, may prove powerful cures of the disease, by stimulating the secretion of the gastric juice and bile. This will account for the action of many remedies strongly recommended.

Another series of experiments was conducted by adding a little of the congee-stool to tank-water, and, to separate specimens of this in test-tubes, adding the various antiseptics and cholera medicines, and watching the effect on the reproduction of the animalcules, compared with water in a standard test-tube to which no addition had been made.*

The explanation of the action of the poison from excessive reproduction is very probable and satisfactory in cases which terminate fatally; but in many cases the violent prostration or collapse suddenly ceases, and

* These experiments were commenced in October last by Drs. Cunningham and Lewis, but have been suspended for some weeks, as the animalculæ have ceased to reproduce, probably in consequence of the cold season.

the ordinary healthy state of the body returns in a few hours. This may be explained by supposing that the symptoms are caused by the movements and living action of the germ; and that, on these movements ceasing, the symptoms would subside; but this cessation could not arise from the reappearance of the gastric juice, for, in collapse, none is secreted. In the innumerable cases which recover without the use of medicine, Nature alone must supply the remedy. Simultaneously with the cessation of the collapse and commencement of reaction, there is a change observed in the character of the evacuations, showing the presence of bile. Could this be Nature's repressing influence? *Post mortem* examinations show that there is bile accumulated in the gall-bladder, which, in the most intense cases, is often much distended. The bile evidently has not been allowed to enter the intestines, from some unknown cause. If this cause were spasm of the *ductus communis*, which would give way or cease on the total paralysis of the nervous system when death approaches, the bile might then escape, and, entering the intestines, come into contact with the active germ, and cause its death or suspend its action, so as to allow the restoration of the natural functions and of health.

The primary action of the poison is on the nervous system, and is limited to that portion supplying organic life. The secondary action is on the mucous surface of the intestines. I attribute the diminished secretion and excretion to diminished nervous energy, and the feeble action of the heart to the same cause; which, when intense, induces total suppression of the functions, or paralysis of the sympathetic and ganglionic nerves. The term paralysis is generally associated with loss of muscular power, and may be considered inappropriate where the muscular contraction of the intestines forms one of the most prominent symptoms in one stage of the disease, after the other functions of the sympathetic have been suspended. But they yield in progression—first, the secretion of gastric juice; then that of bile; then of urine; then the purification of the blood in the lungs; and, last of all, muscular action. Here, the name of paralysis is appropriate, and the meaning of partial paralysis is explained as indicating diminished functions. The secondary action of the poison I consider to be on the mucous surface of the intestines, causing irritation of the villous coat by the destruction of the epithelium. This arises from a change in the course of the vitality of the cholera-poison from a state of dormancy or inactivity to one of active life and reproduction, somewhat analogous to that of the *cysticerci*, which are quiescent and unproductive in the body until they reach the mucous coat, where they reproduce. This is an increased, not a paralysed action; paralysis here only comes on with the final expiring efforts of nature; when the spasm of the *ductus communis*, which retained the bile, gives way, and may permit the oozing out of the accumulated fluid, and allow reaction by discharging this vital active antidote to the poison.

Some who have thought much on this disease, consider that the primary action of the poison is to produce spasm of the small arteries; that the vomiting and purging indicate spasm of the muscular coat of the intestines; and that the spasm in the *ductus communis* and ureters causes the retention, not suppression, of the bile and urine. One of our most intelligent writers, Dr. G. Johnson, considers that the primary action is spasm of the pulmonary artery, impeding the circulation in the right side of the heart; that the lividity in collapse is caused by the impeded venous circulation, and "that, when this is not present, lividity is absent, and the lips of a patient in pulseless collapse are often as florid as in perfect health."

This must be a different disease from what we have in India, where I have never seen florid lips until reaction has set in. The *post mortem* appearances, described by Dr. Johnson, in the heart and lungs, are in accordance with my experience; but the gorged state of the right side of the heart may depend on diminished power of the heart to empty itself, as much as on increased power of resistance to the passage of the blood through the pulmonary arteries. A theory to be correct must explain all the symptoms, and not be incompatible with the course of the disease. This theory does not explain all the symptoms of the disease, either in its milder development of *malaise*, when there is no oppression of breathing, or in its most intense form, when there is neither vomiting nor purging. In the former there is torpor of the intestines, with a dull dead feeling in the abdomen, well known to all who have had much experience in cholera epidemics, which contraindicates active spasm. It is evident that the muscular contractility of the intestines and *ductus communis* survives the failure of the secretions, in the milder form and intermediate stage; but this also subsides as the disease approaches a fatal end. I have not observed any priority in the symptoms in the lungs to those in the liver and kidneys, or to the feeble action of the heart. The circulation can be restored, in the most advanced stage, by the transfusion of saline fluids; the pulse revives, colour and warmth return, the spasm of the vessels in the lungs is no longer present;

but health is not restored, the abdominal secretions—the bile and the urine—do not appear, and death ensues.

The *post mortem* examination of those who die from the most intense form of the disease, shews no signs in the intestines, different from what is found after sudden death from lightning, with the exception of the presence in the duodenum and jejunum of a whitish fluid, which, when diluted with water, resembles congee-evacuations. When death takes place after partial reaction, there is often a pinkish tinge over the intestines; they feel doughy and thickened, with the mucous coat abraded, and deficient in epithelium. When reaction has been completely established, before death, there are often signs of inflammation of the mucous coat, with ulceration. These secondary signs are not always present, yet they are considered by some writers as indicating that the disease is essentially an inflammation of the mucous coat of the intestines. As the ordinary symptoms of such inflammation are totally wanting in the earlier stages of the disease, and the morbid changes are not found in its most intense form, it may be inferred that this state supervenes as a secondary change, induced by the impurities retained in the blood, by irritation from the vitiated contents of the bowels, or from stimulant remedies exhibited, as well as from the action of the poison on the epithelium. These signs are most prominent after protracted sickness. The rapidity with which convalescence takes place in slight cases, or when collapse has not long endured, contraindicates any extensive inflammation of the bowels. In the stage of *malaise*, the employment of depletion, or of depressing remedies, is most dangerous, often leading to collapse and death.

[To be concluded.]

TINCTURE OF PERCHLORIDE OF IRON IN ACUTE RHEUMATISM.

By W. T. GREENE, M.B., L.R.C.S. Edin., etc.

FOR a considerable time I had been aware of the beneficial effects of the perchloride of iron, in small doses (ten to fifteen minims), during convalescence from acute rheumatism, but had never prescribed it during the acute stage until after the publication of Dr. Russell Reynolds' paper in No. 468 of the BRITISH MEDICAL JOURNAL, when the success which he met with induced me to give it in several cases.

W. T., aged 33, had suffered from acute rheumatism on two previous occasions. I commenced the treatment by ordering thirty-minim doses of the tincture every four hours. On the following day, the pain in his knees was considerably better; the shoulders and wrists, however, had become affected, and he complained of headache. The dose was reduced to twenty minims, and convalescence was complete on the tenth day.

In the next case I was not so successful. Mrs. Y. had an attack of scarlatina, which had been immediately followed by acute rheumatism, the lower extremities being principally affected. I ordered her twenty-minim doses of the tincture of perchloride of iron every four hours, and found her decidedly improved the next day, but complaining of great debility. I increased the dose by ten minims: next morning, the pains in her joints were better, but she had a slight pain in the region of the heart. I ordered a blister to be applied, which was done. In the evening, the husband sent for me, saying that his wife was dying. On my arrival, I found her quite unconscious, with great dyspnoea and tumultuous action of the heart. I sent for some tincture of digitalis, and applied such remedies as were possible under the circumstances; but the power of swallowing was gone, and very shortly afterwards she breathed her last. No *post mortem* examination was allowed.

Mr. C., aged 36, had suffered from acute rheumatism when a child. When I first saw him, the rheumatic affection was confined to one knee and ankle. I ordered him thirty-minim doses of the tincture of perchloride of iron, and next day found that both knees and one shoulder were affected. He could not stir, and was suffering most acute pain. In addition, every now and then all his limbs were convulsed by a tetanic spasm, especially violent as he was dropping off to sleep, when it almost amounted to opisthotonos. He said, "You have been giving me steel, I am sure." I replied that I had done so. "Yes," he said, "I was sure of it; I have taken it once or twice before, and it always has the same effect." Although he had only taken four doses, it was nearly a week before the spasmodic twitching of his limbs entirely subsided. He made a good and speedy recovery under alkaline treatment.

I may have been specially unfortunate in my cases, but I certainly shall hesitate about prescribing iron in acute rheumatism in future—at least until the more violent symptoms have been subdued.