end, a stove in the centre, and ten extempore wooden beds along each side. These *Baraken*, as they are termed, form an admirable *Feld-Lazareth*. They are only suitable, or intended, for temporary occupation; but for such a purpose are, I think, very suitable, and much better than tents, which are liable to numerous inconveniences. Stromeier had a large staff of surgeons and assistants under him; and his duties as *Generalstabsarzt* did not confine him to one ambulance. The great civil surgeons who hold the highest posts in the German armies visit the different field-ambulances at fixed times, when there are regular consultations with the surgeons attached to them on all cases of importance; and at these visits the operations are performed, which are decided upon after consultation as being needful. There can be little doubt that the advantages of such a system, both to the army medical officers themselves and to the patients under their charge, in having the counsel and assistance, at all times and in all cases of difficulty, of the most matured surgical advice in the country, are very great. There is no undue interference, however; and I have not heard that any jealousies are excited, or that the system does not work to the great benefit of all concerned. Certainly, in the case of Stromeier, who is beloved and respected by all who come into contact with him, nothing could be more harmonious than the relations which subsisted. On the occasion of my visit to him, he insisted on my performing an amputation of the thigh, which was required for an injury to the knee. We had afterwards a long conversation together, in the course of which he asked me if I had arrived at his opinion that the practice of military surgery was unsatisfactory in the extreme. He said he was now passing through his third campaign; and that the longer he lived, the more deeply was he alive to the seemingly unavoidable evils which attend the practice of surgery during war.

CASES OF FRACTURE OF THE BASE OF THE SKULL: WITH REMARKS.*

By C. E. PRIOR, M.D., Bedford.

IN these cases I have no novelty of treatment to advocate, or theory of interest to propound; I wish simply to set before you a plain unvarnished history of several cases of an injury not very infrequent. A history of this sort, collected from a rather extensive field, may serve to qualify some important points of diagnosis, and may lend accuracy to the prognosis of one of the most formidable accidents to which the human frame is liable. My cases are collected partly from my own experience as a coroner, some from my own practice, some communicated by my friends Mr. Thurnall and Mr. Johnson, both of Bedford.

CASE I.—Fracture of the Anterior Fossa from Blows with Fists: Death within Six Hours: Post Mortem Examination.—In May 1863, about 11 P.M., a gentleman and lady, who had been spending the evening with friends, were returning home through the back streets of Bedford, when the former was insulted by two ruffians who had been "on the spree." On his attempting to defend himself, he was struck violently on the face by both, and knocked down. He rose with difficulty, and was followed by them for some distance, when the assault was renewed with at least equal violence, and he was knocked down again. He lost much blood at both places. Being at length at liberty to proceed, he went to the residence of the chief constable of the county, and endeavoured to ring the bell; but his wife said she thought he would have died on the doorstep. Soon after this, he reached home; and a medical man was summoned, on whose arrival, however, he seemed better. Hæmorrhage had nearly ceased; and he went to bed. Internal hæmorrhage probably took place soon after this, and he died rather suddenly about 4 A.M.

The *post mortem* examination showed a slight wound below the left eye; contusions of the forehead and nose, and under the eyes; fracture of both nasal bones, and clotted blood in the nostrils, and bruises on various parts of the body. There was extravasation of blood between the pericranium and frontal bone above each orbit; a clot of about four ounces was effused over the occipital bone, and about two drachms between the arachnoid and pia mater. There was also effusion on raising the calvarium; and a zigzag fracture of about an inch and a half long extended across the left orbital plate of the frontal bone, with some depression.

CASE II.—Fracture of Middle Fossa (Pars Petrosa) from a Fall on the Vertex: Death in Three Days: Post Mortem Examination.—A boy, aged 9, fell from a tree in a village four miles from Bedford.

When taken up, he was bleeding in a stream from the left ear, and there was a compound fracture of the left thigh. When found, he requested that he might be laid down. He was taken to the Bedford Infirmary. He was sensible on his arrival there, and continued so for some hours; then the pupils dilated, and a copious colourless fluid escaped from the ear. He remained insensible until his death, three days after the injury. On *post mortem* examination, a fracture was found extending through the petrous part of the temporal bone, and upwards through the squamous part; the brain was much congested, and suppuration had commenced immediately over the fracture.

CASE III.—Fracture of Middle Fossa (Pars Petrosa) from a violent Blow on the Head: Recovery.—J. S., an engine-driver on the London and North Western Railway, left his post at the engine at some part of the line between Bletchley and Bedford, and, crossing over the top of the carriages, arrived at one in which there were some persons that curiosity induced him to watch. It is supposed that, while he was leaning his head over the side of the carriage to look into the window, his head came into contact with the brick-work in passing under one of the railway bridges. On the arrival of the train at Bedford, the man was found on the top of the carriage quite insensible; he was at once taken to the Infirmary in an unconscious state; his head was then shaved, and he was placed in bed. No fracture of the upper part of the skull could be detected. Blood escaped copiously from one ear, and this was followed by an escape of a considerable quantity of cerebro-spinal fluid. He remained insensible for a fortnight; breathing was stertorous for the first few days. About the fifth day after the accident he had three or four epileptic fits. About the fourteenth day, he became slightly conscious, and from that day gradually mended; and a deafness from which he had at first suffered was ultimately lost. He left perfectly well at the end of three months, and from that time to the present has suffered from no head-symptoms of any kind. He follows his old occupation as an engine-driver. The treatment was of the ordinary kind; cold to the head, counterirritants and warmth to the feet, and mercury to slightly affect the gums.

CASE IV.—Fracture of Middle Fossa (Pars Petrosa) from a Fall on the Vertex: Recovery.—A boy, aged 12, subject to epilepsy, was in a tree getting birds' nests, and is supposed to have fallen from the tree with his head downwards, in one of his fits. He was brought to the Bedford Infirmary unconscious, and suffering loss of blood from the ear; the blood was followed by clear fluid, which continued to escape for some days in largish quantity. After remaining unconscious for several days, he began slowly to regain his senses, and continued to improve until he left the hospital. I believe his recovery to have been perfect and permanent.

CASE V.—Fracture of Middle Fossa (Pars Petrosa) from Fall on Vertex: Death after Three Days.—T. W., aged 29, was engaged in painting a portion of the Bedford Union Workhouse. He had occasion to ascend a ladder to a height of forty-five feet; the ladder broke at about twenty-five feet from the ground, and he fell, striking a high wall as he came down, which turned him so that he fell with the upper part of the head on a brick pavement. I found him, on my arrival, bleeding from a wound above the right eye, and from the ears. He was in a state of collapse, and quite insensible. On the next day, a discharge of clear fluid came on from the left ear, which was very profuse, as much as an ounce to an ounce and a half passing in an hour; nevertheless, he appeared a little better, and was able slightly to recognise persons; but he afterwards gradually sank, and died three days and a half after the accident. No fracture of the upper part of the skull could be detected.

CASE VI.—Fracture of Middle Fossa (Pars Petrosa) from a Fall on the Head: Recovery.—W. P., aged 55, labourer, was knocked down in a fight at Sharnbrook on June 4th. He said that his head went on to the pavement with a bump, and rebounded. He was admitted into the Bedford Union Workhouse on June 7th, bleeding from the left ear, confused, and suffering great pain in the head; pulse 84. There was great deafness; but this, he said, had already been the case. The treatment consisted of cold to the head and calomel with aperients. A discharge of clear fluid from the ear came on June 8th; a dessert-spoonful passed in twenty minutes, probably, therefore, several ounces in twenty-four hours. He was removed to the Infirmary on June 9th: after his admission there the discharge of fluid gradually abated, and he slowly and partially recovered. He was readmitted into the Workhouse on July 2nd, still very deaf, and complaining of giddiness and pain in the head. He left the house on July 19th in that condition; and after some days he returned, not much better.

CASE VIII.—Fracture of Middle Fossa and Pars Petrosa by a Fall on the Vertex: Death in Four Days.—J. G., aged 33, was thrown out of a runaway cart, and fell on his head. While he was in this position, an

* Read before the South Midland Branch.

empty cart appears to have gone over him, at Kingston, three miles from Bedford. He was immediately taken to the Bedford Infirmary. On his admission, he was slightly sensible. There was a superficial cut on the back of the head, and a copious flow of blood from the right ear. On the following day, clear fluid began to escape from the ear, and continued to do so. The same day he was better, and still better on the third. On the fourth day he became sleepy; comatose symptoms set in, and he died early on the fifth day. On *post mortem* examination, the posterior surface of the petrous bone was found fractured across; there was also a considerable clot of blood (an ounce and a half to two ounces) between the dura mater and the skull on the left side; also considerable recent inflammation on the under surface of the cerebellum.

CASE IX.—Extensive Fracture of Base of Skull from Violent Pressure on Both Sides: Death after Two Hours.—W. F., an agricultural labourer, aged 34, fell from the shaft of a waggon on which he was sitting driving at Bromham, Bedfordshire, when the waggon-wheel ran over him near the head. When taken up, he was insensible, bleeding from both nostrils, and from the right ear. He died immediately after admission to the Bedford Infirmary, about two hours subsequently to the accident. On *post mortem* examination, there was found a large quantity of extravasated blood on the surface of the brain, the side of the skull fractured through the squamous portion of the right temporal bone, and the base extensively fractured through the petrous portion.

CASE X.—Extensive Fracture of Base of Skull from Violent Pressure on both Sides: Recovery.—About twenty-five years since, a bricklayer had his head jammed between two buffers of a train (cracked like a cocoa-nut, in fact). Upon his arrival at the Bedford Infirmary, he was quite insensible, with blood flowing from both ears, the nose, and mouth. For several days after the accident, there was a watery discharge from both ears. Consciousness returned in two or three weeks, and he progressed favourably but slowly, and left the Hospital in about five months. During the time he was in the Infirmary, he suffered from severe pains in the head, which continued for a long time after he was discharged, and the muscles on the right side of the face became paralysed. He is at the present time following his usual employment as a bricklayer, and is in good health; but there is still want of power in the muscles of the right side of the face.

CASE XI.—Extensive Fracture of Base of Skull from a Violent Blow, and Lateral Pressure: Death.—T. S., a child aged 5, in the summer of 1867 was watching the men at work at the Midland Ouse Bridge at Bedford. One of the men engaged was carving a balk of timber, when the iron rod broke, and the end of the balk fell on the child's head, crushing it into the ground. Death was almost instantaneous. Blood flowed abundantly from the mouth and ears. On examination, no fracture could be found at the upper part of the skull; but it was detected at the junction of the occipital with the parietal bone, and probably extended across the base of the skull.

The last three cases are examples of the greatest degree of violence which this part can sustain with the faintest possibility of recovery. In each the violence was excessive; in each pressure operated from both sides; in one, death was almost or quite instantaneous; in another, the patient survived for two hours; in the third astonishing, and, I should think, almost unique case, the subject is living to this day.

CASE OF CARBOLIC ACID POISONING.

By ALEXANDER OGSTON, M.D.,

Assistant-Professor of Medical Jurisprudence in the University of Aberdeen.

ON October 26th, 1870, Patrick McGrath, aged 47, sergeant-instructor of volunteers, Tarves, found in a rifle-shed a bottle of crude carbolic acid, part of which he swallowed, in mistake for bitters or other fluid. It is calculated that he swallowed, at most, an ounce or two of the fluid. This, when examined, seemed to be of considerable strength, crude, black, and oily. He then proceeded to a druggist's shop, but left this at once, without taking any drug. Shortly afterwards, he was found lying insensible, and was taken to Dr. Irvine's house. Dr. Irvine kindly handed me his notes of the case, which are as follows.

"Last night, October 26th, P. McG. was brought to my house at 5.30 P.M. He was reported to have been accidentally poisoned. He had a strong smell of carbolic acid, was unable to speak or walk, and was quite unconscious and in a state of apparent apoplexy, with stertorous breathing and puffing of the lips at each expiration. The pulse was not much disturbed. The heat of surface was natural. I laid him on a couch, covered him warmly, and administered castor-oil to dilute

and carry off the poison." The bottle containing the poison was distinctly labelled "Carbolic Acid, Poison." The notes continue: "On coming back, I tried with my patient olive-oil and a powerful emetic, which had no effect. I then sent for my son, who brought a stomach-pump; it would not work, and was consequently not applied. Subsequently, carbonate of ammonia and brandy were tried to no purpose. The patient's lips, gums, and tongue were white, as if sodden. After he had been laid on the couch, there was for three or four hours no muscular movement at all. The eyelids were closed, and the pupils contracted to the size of a pin-point. The first indications of returning sensibility were in the eyelids. About 12 o'clock, he began to open his eyes to some extent. Subsequently, he moved his arms, then his feet and legs; and afterwards made an attempt to raise his shoulders. After this, the pupils gradually expanded, but never became large. He began to speak, and asked for cold water to drink; but he never acquired complete consciousness. The next phase was affection of the lungs. There was very hard and laborious breathing, and difficult muco-purulent expectoration. After some time, the breathing became easier, just as suddenly as the coma had improved. He then complained of severe pain in the back. The pulse, which at first was not much disturbed, ran up to 84, 90, 96, and 106—not always alike, but fluctuating. Respiration from 46 to 48. At 2 o'clock in the morning, he passed naturally a quantity of very dark coloured urine, having the smell of carbolic acid. He afterwards attempted again to pass urine, but without success. No real reaction ever took place, notwithstanding the frequent use of warm applications. The brow, face, and hands were covered with cold clammy perspiration. Collapse came on, and at 7 A.M. death closed the scene, after thirteen hours and thirty-five minutes of suffering."

On proceeding to examine the body thirty-two hours after death, it was found in the following state. Rigor mortis was present. There was lividity of the back parts, and of the front of the neck and chest. The face was sallow. The pupils were natural. Watery froth and yellowish fluid, smelling of carbolic acid, were present at the mouth and nose. There was a clot in the superior longitudinal sinus. The dura mater and arachnoid were congested, the blood smelling of carbolic acid. Serum, with a distinct odour of carbolic acid, was found in the subarachnoid tissue and in the ventricles. The cineritious brain-matter was pinkish in hue. There was intense congestion of the vessels on the under surface of the pons Varolii and medulla oblongata. The mucous membrane of the mouth, throat, and gullet was unnaturally white and sodden in appearance, and at parts was soft and easily detached. There were two fluid-ounces of thick brownish pulp, smelling of carbolic acid, in the stomach. At its greater curvature, the mucous membrane of the stomach was studded with hard, reddish, and elevated points, of the size of small shot, without any marked congestion around them. Similar but smaller points existed on other parts of the stomach—showing, however, a more linear form, and tending to follow the course of the vessels. The stomach was otherwise natural, or perhaps a little more vascular than usual. Portions of the small intestine were slate-coloured, especially in the duodenum and upper part of the jejunum, but otherwise natural in every way, and containing a brownish pulpy matter. The salivary glands, pancreas, and mesenteric glands, were all pale. The heart was healthy, and full of dark firmly clotted blood on both sides; the right side containing four, and the left side two ounces. A minute fibrinous clot was found in the pulmonary artery, imbedded in a large black clot. The trachea was congested, and contained fluid similar to that found at the mouth and nostrils. The lungs were bulky and very cedematous. Fatty patches were present on the upper surface of the liver. The gall-bladder was enormously distended with bile. The spleen was soft; the kidneys fatty. Four ounces of turbid urine were found in the bladder. The blood, urine, and solid textures of the body, all smelt of carbolic acid. The blood was every where very firmly clotted.

ON SYPHILITIC RENAL DROPSY.

By S. MESSENGER BRADLEY, F.R.C.S.,

Lecturer on Human and Comparative Anatomy, Royal School of Medicine and Surgery, Manchester.

MEDICAL reasoning is too often hastily deductive in character; in other words, we too often reason from the particular to the universal, and rush from an unproven hypothesis to an erroneous and dogmatic conclusion. This constitutes the very sin of the Aristotelian method, and, if followed constantly, would bring medical science into equally well-deserved disrepute, as it did the old Grecian philosophy.

Still, reasoning thinking men will draw deductions; nay, more, we