times as high as 114, 118, nay, it is said, 125 deg., and even by night the temperature may not fall below 105 deg. And it is only what we should expect, when we are told that moisture of the atmosphere and muscular exertion, the one diminishing the loss of heat by evaporation, and the other increasing the heat set free by the body, are powerful excitants of sunstroke, in addition to the mere heat of the air. The deleterious effect of the heat seems to be largely explicable even in the present state of animal chemistry. Weikart declares that the blood begins to clot in the vessels at 109 deg.; the muscles of warm-blooded animals set instantly at 118 deg., and set more gradually at lower temperatures; 113 degs. Fahr. of heat abolish the excitability of nerve in frogs. The rational treatment of heat-stroke which suggests itself is strongly supported by experience; we must aim at reducing as quickly as possible the heat which is killing the patient.

[To be continued.]

LECTURES ON DERMATOLOGY.

DELIVERED AT

The Royal College of Surgeons of England.

By ERASMUS WILSON, F.R.S.,

Professor of Dermatology in the College.

LECTURE VI.

IN reference to treatment, eczema must be regarded as a "solution of continuity", and as such requires a dressing to defend the lesion, for the purpose of keeping its broken edges in contact, and of excluding the operation of external irritants of every kind. A dressing with the benzoated ointment of oxide of zinc fulfils this indication completely. The ointment should, in the first instance, be smeared upon the eczema; the smearing process should be repeated twice or more frequently in the day, so as to secure a permanent covering; and, wherever practicable, strips of linen rag spread with the same ointment may be placed over the eruption and maintained in position by a roller, so as to keep the ointment and the dressing in permanent contact with the diseased skin. In this relation the ointment fulfils the purpose of a light unirritating plaster; it preserves the natural moisture of the skin, and excludes the stimulant operation of the atmospheric air. It is easily replaced when accidentally removed, and in many cases will accomplish every object of treatment from the beginning to the cure of the case. Let me instance an eczema infantile, in which the body is covered with eruption from head to foot; it is hardly possible to conceive a remedy more admirably suited in such a case to meet the exigencies of treatment than the benzoated ointment of oxide of zinc.

It has long been my custom to combine with the zinc-ointment a small quantity of spirits of wine, in the proportion of one drachm to the ounce: the spirit softens the ointment and facilitates its application; it produces a sensation of coolness which is agreeable to the heated surface, and it has besides a gently stimulant effect on the nerves of the skin. The application, as I have already observed, relieves the heat, the stiffness, and the itching, and the relief continues until the ointment dries up or is accidentally removed. When either of these events takes place, the ointment must be repeated in the same manner as before. Our aim should be to favour a thin deposit or concretion on the surface, which should occupy the place of the broken or excoriated epidermis; such a deposit or concretion is, in fact, an artificial cuticle, a kind of natural plaster, under which the skin is preserved in a state of repose, and the inflamed and irritable tissues have time given them to recover their normal status. You will perceive that eczema brings before us a surgical lesion in which the application of a plaster or dressing of any kind is often perfectly impracticable, and therefore we are obliged to have recourse to a remedy which shall possess the intrinsic property of adhesion to the skin without the aid of other coverings. This will explain our anxiety to have the inflamed surface thoroughly coated with the ointment, and to have the ointment repeated as often as, from any accident, it may by chance be displaced; and, of course, wherever practicable, means may be contrived, such as bandages and compresses, to render the dressing permanent. Above all, it is important to avoid washing the skin at this stage of the disease: washing would remove the ointment and open up the excoriation afresh; it would be undoing that which it had taken much

labour to do; it would be unsurgical, by creating disturbance where rest and position were of the utmost importance. There can be no want of cleanliness where a pure oxide of zinc ointment is properly used; and if secretions be poured out from the skin, they may always be absorbed by means of a soft napkin, and, together with crusts, may be gently wiped away. Mothers will sometimes hear with wonder the command to avoid washing their children in a case of eczema infantile; but they are generally ready to admit that the eruption is always more uneasy after the washing than it was before, and their instinct quickly assures them that their instructions to avoid washing are correct.

Eczema enjoys the peculiarity of presenting, in a general attack of the eruption, a variety of degrees or forms of manifestation on different parts of the body; thus it may be erythematous in one place, ichorous in a second, and squamous in a third. The zinc-ointment is equal to the relief of every one of the forms of the eruption; but it not unfrequently happens that we may find an advantage in using a powder in one situation, the lime-water and zinc-lotion in a second, and the ointment in a third. And in the adoption of these means we are to be influenced rather by the sensations of the patient than by any preconceived theory of the disease. In general terms, it may be said that whatever gives rise to pain or uneasiness is bad, and will require to be changed.

Eczema is essentially chronic as to its course; but it manifests stages in its progress which are as decidedly acute. The treatment which we have just been considering is strictly palliative, and is therefore especially adapted for the early and acute stages and acute manifestations of the disease, but is equally applicable for the relief of heat, itching, and dryness, in every stage of the affection. If I were asked how I should distinguish between an acute and a chronic eczema, I should say that the acute stage was denoted by erythematous congestion and moist excoriation, and the chronic stage by infiltration and desquamation, the infiltration giving rise to thickening and induration. I shall now suppose that the eczema has reached its dry, squamous, thickened, and indurated stage. It has assumed that condition which, in ulcers, for example, is termed indolent, inveterate, and so forth. The zinc-ointment removes the scales, heals chaps and fissures, and relieves in some degree the itching, but does nothing towards the dispersion of the thickening and induration of the tissues—that is, towards the cure of the disease. The time, in fact, has come when we must contrive to awaken the tissues from their indolence and lethargy, and direct, as far as we are able, the newly awakened life towards a normal state of action—that is, towards a restoration of healthy function and tone.

Our purpose, therefore, is no longer palliative; it is stimulant or tonic; the palliative treatment has accomplished all that it can; now we have to consider the nature and mode of application of the stimulant treatment. Our stimulants possess a great range of power, and, if we knew exactly the amount of resistance which we should meet, we might apportion our power with exact accuracy. But as this can never be the case in dealing with so delicate a machine as living organisation, the laws of surgery would prescribe to us that we should begin with the mildest of our stimulants and advance progressively and to the best of our judgment to the higher degrees. The philosophic surgeon will not hesitate to adopt this course as one in every way consistent with his knowledge of vital phenomena; and, therefore, we may proceed to inquire, In what do our stimulants consist?

The first and best of our tissue-tonics or local stimulant remedies is undoubtedly soap. It will be remembered that I have prohibited soap in the earlier and acute stages of the eruption; therefore soap is new to the morbid skin, and capable, consequently, of producing a more powerful effect than would otherwise be the case. The chronic eczema should be thoroughly washed with soap, combining with the solvent action of the soap upon the epithelium such an amount of moderate friction and compression as may reach the vascular and the infiltrated tissues. After the washing, the surface is to be dried with a soft napkin, and, as soon as dried, dressed with the zinc-ointment like an eczema of an earlier period. The next day we examine very carefully our eruption, and, if we find no signs of excessive irritation present, we renew the saponaceous washing and repeat it again and again until the eczema is healed. It will soon be apparent how frequently the process may be repeated, whether once a day only or twice, or whether the reapplication must be deferred for several days. The immediate consequence of the stimulant treatment will probably be the development of an exudation on the surface, which will relieve the infiltration and turgescence within; and a few repetitions of this exudation will so far tend to empty the tissues of their excess of fluids that they will be enabled to return by degrees to their normal state.

This, in fact, is the explanation of the *modus operandi* of the powerful stimulants which we hear of as being from time to time useful in the cure of chronic eczema; such as strong alkaline soaps, strong solutions of potash, strong mercurial ointments, strong solutions of nitrate

of silver and tar. And, in illustration of the same principle, I may call your attention to a remark which I made in reference to the treatment of No. 27; namely, that, after an obstinate resistance of a variety of remedies, the disease was suddenly cured by one application of the tincture of croton, pencilled over the inflamed surface.

If I wished to reduce the principle of local treatment to an aphorism, I might do so by the use of the two words *palliative* and *stimulant*. Among the palliatives, there is one which is occasionally of much service for the relief of heat, tension, and pruritus; namely, water-dressing, whether the moist application be made in the ordinary way with an impermeable covering, or employed in the shape of a cold starch-poultice. The water-dressing may be used constantly for a period, or during the night only; and its use may be combined with that of the lime-water lotion, zinc ointment, or saponaceous frictions and ablutions. The water-dressing is very useful in accelerating the separation and removal of crusts, in promoting exudation from infiltrated and cedematous tissues, and in preparing the tender skin for the treatment by ointment or the treatment by desiccating lotions. If a water-dressing had been kept applied to the leg in Case XLVII for twenty-four hours, the whole of the dark crust visible in the model might have been washed away. The morbid skin would then have been prepared to receive the dressing of zinc-ointment; and this, with the support and moderate pressure of a properly adjusted roller, would have effected a vast improvement in the disease in a very short space of time.

One symptom which more than any other torments both the surgeon and the patient in eczema is *pruritus*; and very frequently we are called upon to change our plan of treatment in order to combat this annoyance. When the pruritus proceeds from ordinary irritation, it may be subdued by the oxide of zinc ointment; and a moderate friction with this ointment may be had recourse to as often as the itching returns. If allowed to continue, the pruritus will sometimes assume a neurotic and neuralgic character. It will come on after any slight exertion, after the taking of food, upon changes of temperature, and especially at night, sometimes on first assuming the reclining posture, and at other times in the early morning. It is always intermittent, and presents such a variety of manifestation as to make evident that sometimes the peripheral nervous plexuses are the seat of the painful sensation, sometimes the perforating twigs of the cutaneous nerves, and sometimes the larger branches and even the trunks of the nerves. There may be a gradual transition from a feeble itching to a pruritus which vibrates through the whole nervous system, or to a state of neuralgic pain such as has suggested the term *eczema neurosum*.

When the zinc ointment, with the addition of spirits of wine, camphor, tannic acid or carbolic acid, fails to relieve the pruritus, we may sometimes meet with help from hot water, from a cold starch-poultice, from water-dressing; from lotions of soda, hydrocyanic acid, juniper tar, or a solution of nitrate of silver. But all these remedies must be regarded simply in the light of adjuvants to the principle of treatment already discussed; and, when the neuralgia is decided, we must seek for relief in constitutional means.

The stavesacre ointment is often very successful in the relief of the pruritus which sometimes follows scabies; and as this remedy, next to sulphur ointment, is one of our best means of destruction of the acarus, it may possibly happen that the pruritus is protracted by some such overlooked and unlooked-for complication. The pruritus of lichen urticatus is best relieved by Hebra's lotion—a lotion consisting of an ounce each of soft soap, juniper tar, and alcohol, diluted with five ounces of water. And the sometimes ungovernable itching of lichen planus is with the most certainty mitigated with hydrocyanic acid suspended in an emulsion of bitter almonds.

Another modification of the local treatment of eczema is the natural consequence of its seat. Neither powder, nor lime-water lotion, nor zinc ointment, would be suitable for application to the eyelids, or within the meatus aurium; but for both these regions an excellent substitute will be found in an extremely diluted nitrate of mercury ointment, or in the glycyrrine of tannic acid of the *British Pharmacopæia*. Again, the oxide of zinc ointment is unsuitable on the scalp, in consequence of clogging the hair, and thereby giving rise to much inconvenience; but, in the latter situation, a far better remedy presents itself—namely, the nitric oxide of mercury ointment, diluted with benzoated lard in the proportion of one part to three of the diluent. The case of dermatitis exfoliativa (No. 6), suggestive of an extensive burn, was most efficiently relieved by the linimentum calcis and the carron-oil.

In bringing to a conclusion my remarks on the local treatment of the eczematous affections, I should fail in my intention unless I succeeded in leaving on the minds of my audience a forcible impression that the treatment of diseases of the skin is in nowise different from that of surgical disease in general; that the same enemy is to be combated in all; and that the principle of treatment is in the first place *alleviative*, and

in the second *stimulant* or *tonic*. But, just as the Abernethian doctrine of the "constitutional origin of local diseases" is universally recognised by the physician of surgery at the present day, so, in the instance of the diseases before us, the constitution of the patient undoubtedly enacts an important part, and constitutional treatment possesses a corresponding value.

[To be concluded.]

ABSTRACTS OF CROONIAN LECTURES

SOME POINTS CONNECTED WITH THE ELIMINATION OF NITROGEN FROM THE HUMAN BODY.

Delivered at the Royal College of Physicians, London.

By E. A. PARKES, M.D., F.R.S.,

Professor of Hygiene in the Army Medical School at Netley; Emeritus Professor of Clinical Medicine in University College.

LECTURE II.—*Friday, March 17th.*

IN the consideration of the way in which nitrogen is eliminated from the body, it is necessary to be acquainted with some simple facts. One of these, of great importance, has only lately been made out; viz., that the amount of nitrogen removed from the body may be very nearly accounted for by the quantity ingested—that is to say, the exit of nitrogen is regulated by its entrance, and to a very slight extent only by the muscular action. This fact was not recognised until it was pointed out by Pettenkofer and Voit. Dr. Parkes, from some experiments on healthy men, has found that, by simply varying the amount of nitrogenous food, he could increase or diminish the amount of nitrogen eliminated. Thus, to one man he gives for ten days food containing 195 grains of nitrogen *per diem*; the amount excreted in the urine was 183 grains. He then increased the amount of nitrogen to 215 grains; the quantity of excreted nitrogen rose to 204 grains; and, on giving 274 grains of nitrogen for another period of ten days, the amount eliminated rose to 242 grains. Other similar experiments were referred to; all of which showed that the amount of nitrogen excreted can be varied at will by varying the quantity ingested.

In some circumstances, there is not this correspondence between the exit and entrance of nitrogen. This occurs, for instance, where much nitrogenous food is given at once. If the excess consist of the albumen of fowls' eggs, it is not assimilated, but passes off in the urine. In dogs, however, and occasionally in man, it may be assimilated.

Again, even where the amount of nitrogen in the food is greatly reduced, there may be a considerable elimination by the urine; even in animals starved, nitrogen has been found in their urine for three weeks. Dr. Parkes, having fed three healthy men on a diet of starch and fatty food, without nitrogen, found that still a large amount of nitrogen was eliminated.

Why should the egress of nitrogen from the body be modified by its ingress? An attempt has been made to explain this by supposing that the nitrogen taken in with the food was in excess of the wants of the body—a mere surplussage, to be got rid of. This doctrine, however, of "luxus-consumption" (as the Germans call it) has now but few adherents. If the amount of nitrogen in food be cut down, its exit will be lessened; and when the quantity of nitrogenous food is again increased, the amount excreted will not at once correspond to the quantity taken, a part being evidently retained in the body to compensate for the previous deficiency. Thus a man, on nitrogenous diet, excreted 110 grains of nitrogen daily for five days. Nitrogenised food was then withheld for two days; and was then restored for four days in the same quantity as before; he now excreted only 85 grains instead of 110, showing that a portion had been retained to make up for the loss. In another experiment performed by Dr. Parkes, the results were the following. For twenty days, a man had food containing 300 grains of nitrogen, and excreted 305 grains. For five days, he had only half the quantity of nitrogen; and for another five days his food contained none. After this, the amount of nitrogenised food was restored to its quantity at the commencement of the experiment; and, instead of the nitrogen excreted corresponding at once with that taken in, it rose gradually, the quantities being, during four days, 189, 242, 253, and 261 grains; thus showing a gradual return to the normal amount of excretion. These