GULSTONIAN LECTURES

ON

THE HEAT OF THE BODY.

DELIVERED AT

The Royal College of Physicians, London,

MARCH 1871.

By SAMUEL J. GEE, M.D., F.R.C.P.,

Assistant-Physician to St. Bartholomew's Hospital and to the Hospital for Sick Children.

LECTURE III.

WE have proved that in pyrexia there is augmented generation of heat; also that the products of combustion are increased in quantity; a third point remains to be discussed. Is there a corresponding consumption of combustible? In other words: Do the tissues of the body waste more rapidly in the pyretic state than in the state of health? In health, the ingesta and egesta must exactly counterpoise each other, so that the weight of the body may remain unaltered. If the consumption of the tissues be increased (which we presuppose to be the case in pyrexia), whilst the quantity of the ingesta remains as before, the body obviously must lose weight. But, in the pyretic state, the problem is not so simple. The ingesta are usually diminished; moreover, the digestive powers are impaired; so that, for these reasons, a fever-patient would lose weight, independently of the pyrexia. And the difference in these respects is very great in different persons: a fact which explains the different conclusions to which different observers have come; some declaring that the loss of body-substance is greater than in a state of hunger, others that the loss is inconsiderable.

It is more easy to ascertain the loss of weight which corresponds with the different stages of a febrile disease in the same patient. Upon this point I will refer once more to Leyden. A number of careful weighings, in different forms of fever, led to the conclusion that the greatest loss of weight occurs during the crisis—a loss equal to about one per cent. of the whole body-weight in twenty-four hours. The loss of weight shortly after the crisis, and during the height of the fever, is about one-half per cent. Still less is the loss during the period of remission; and the loss at the beginning of convalescence is the least of the whole illness; there is a loss, however, even then—that is to say, the loss of body-weight does not cease with the pyrexia.

Loss per cent. of body-weight in twenty-four hours.

Whole illness	.66	High fever	.57
Crisis	1.06	Remission	.45
After crisis	.59	Early convalescence	.24

Taking the fever as a whole, without regard to its stages, Leyden estimates the average loss of weight in the febrile state to be about half that of absolute hunger. Hence he infers that, in ordinary cases, the danger of death from inanition does not begin until the fever has lasted eight weeks. But, obviously, all these data must differ greatly in different cases; death from inanition is certainly at hand so soon as the body has lost forty per cent. of its natural weight.

It would be a mistake to suppose that this loss of weight always indicates a corresponding loss of body-tissue. On the contrary, the great loss during the crisis is undoubtedly due to increase in the insensible excreta, chiefly water. By comparing the loss of the body-weight with the weight of the ingesta and of the urine and fæces, Leyden arrived at the following average results:—

Excreta by skin and lungs per mille of body-weight in twenty-four hours.

Crisis	1.55	After Crisis	.85
Remission	1.20	Early convalescence	.64
High fever	1.00	Apyretic person	.73

You will observe that Leyden has no facts to bring forward concerning the cold stage of fever. However, notwithstanding this defect, his figures enable us to draw the following conclusion; that, comparing the hot and critical stages of fever with the non-febrile state, the insensible excretions are as ten in pyrexia to seven in the apyretic condition. The increase is greatest during the critical period of the fever, at which time the æriform excreta are double the standard of health; now we have already seen that, in the sweating stage, the carbonic acid expired is not increased at all, so that the whole doubled loss during that period must be due to evaporation of water.

The increased temperature of pyrexia does, then, signify increased chemical change. You will remember that, when we discussed the physiology of the natural heat of the body, we dwelt upon sundry chemical processes which go on outside the blood-vessels and within them. And we found that there was much to be said in favour of Mayer's assertion—that not one-hundredth part of the oxidation-processes of the body go on outside the blood-vessels. Are there any grounds for belief that the oxidation of the blood is augmented in pyrexia?

The artificial production of pyæmia in the lower animals, by the injection of putrid matters into the blood, is an experiment of old date, which has been oftentimes repeated. But, more recently, Billroth and Hufschmidt found that even perfectly laudable pus, injected either into the blood-vessels or into the subcutaneous tissue, had power to produce both local inflammation at the spot where the injection was made, and also a general pyretic state. C. O. Weber went a step further, and showed that the same results could be brought about by the fluid of inflamed tissues, and by the blood of pyæmic or even of simply febrile animals. Three or four years ago, many dissertations were published by students of the Dorpat school upon this subject; the chief facts which they established being as follows: that to remove the fibrin from blood, or the corpuscles from pus, by whipping the blood or filtering the pus, does not remove the pyrogenic and phlogogenic power of the blood or pus; and that the height and duration of the pyrexia set up depend upon the amount of pyrogenic liquid injected; when it is large in quantity, death ensues. When a local inflammation is set up, this modifies the pyrexia and prolongs it. For, when the injection is made into the veins without the production of any local phlogosis, the acme of the pyrexia is reached in two or three hours, and in a few hours longer the temperature regains its normal.

I have spoken of the injection of febrile or of inflammatory blood. But healthy blood, injected in large quantity, also raises the temperature. Not so, however, if blood be removed from one vein at the same time that blood is being injected into another. And, with regard to venesection alone, although for a short time (say half an hour) it may depress the temperature, yet afterwards the body-heat not only rises to its original point, but even goes beyond it. That is to say: like unto injection of pus, of pyæmic or febrile blood, or of healthy blood, so venesection also is pyrogenic. The rise of temperature consequent upon blood-letting occurs both in febrile and in non-febrile animals; but in non-febrile animals, the venesection must be very copious to affect the temperature in this way; the rise of temperature begins immediately, reaches its maximum in an hour or two, and then a fall ensues. Even when the body-heat is falling rapidly, renewed blood-letting delays the fall, or even causes a renewed elevation. Under venesection such as this, healing ulcers often begin to open again. In febrile animals, injection of healthy blood sends down the temperature.

I will not follow these experiments further. Underlying them all is the inference that the products of the disintegration of the tissues are productive of local inflammation when injected into a texture, and productive of pyrexia when injected into the blood. Pus is notably a result of increased metamorphosis of tissue. But even healthy blood must contain more or less of the products of regressive tissue-change. In the first lecture we noticed some experiments of Alex. Schmidt which seemed to leave no doubt possible that the blood does, in health, convey materials which are prone to the speediest oxidation; so that an excess of natural blood raises the temperature, because involving an excess of highly oxidisable matters. Again, venesection is pyrogenic, and possibly because it favours the absorption of the intermediate results of the consumption of tissue. And inflammation implies a disturbed metamorphosis, with an unnaturally rapid decay, whereby the tissues locally, and the blood generally, become overloaded with those half-degraded products of tissue-change, which need oxidation so as to arrive at that state in which they can be separated from the body. Hereafter we shall find that the reverse sequence of events, also, is probably true—that pyrexia increases the destruction of the tissues.

Before quitting this part of our subject, let us investigate a little further the source of the pyrexia which is secondary to local inflammation. It is a very old notion that the local disease is a kind of hearth, where the increased heat is generated, and that the universal pyrexia is due simply to the heat carried off by the circulating blood from the local phlogosis. This doctrine has been held by living pathologists, but there can be, nevertheless, small doubt that it is not consonant with fact. The most recent experiments, made by thermo-electric needles, confirm the dogma of John Hunter that the temperature of a local inflammation is never above the temperature of the blood. More than this: the temperature of highly inflamed skin, or of muscle even in the deepest layers next the bone, never reaches the internal temperature of the body as measured in the vagina, the rectum, or the peri-

toneal cavity. In the severest inflammation, the difference almost always amounts to two degrees or more. Next, the inflamed parts in the acute stage of the inflammation, although cooler than the internal viscera, are yet generally hotter than the corresponding non-inflamed parts on the other side. But the difference between the two sides is less, the farther the parts concerned are from the surface. Thus, in the case of a rabbit's ears, an inflamed ear may be eight or nine degrees hotter than the other; whilst in deep-seated muscles, the difference rarely exceeds a degree, and may be none at all. When the inflammation becomes less acute—for example, suppurating wounds—the inflamed parts are often actually cooler than the corresponding parts in the sound limb. Even with regard to the internal viscera, the peritoneum or the pleura may, when inflamed, be nearly a degree less warm than the left ventricle; usually, however, the temperatures are equal. Now, if the pyrexia can but partly depend upon heat abstracted from the local inflammation, there seem to be two alternatives left to account for the influence of a phlogosis upon the general body-heat. Either the pyrogenic influence is conveyed by the nerves to a supposed heat-regulating centre (a subject which I have postponed hitherto), or the vessels absorb from the wound some matters which set up the fever. The experiments of Brener and Chrobak, although not free from objection, seem to militate against the nervous hypothesis. They removed portions from all the nerves supplying a dog's leg. They then suffered the wounds to heal; and this required, in general, one or two months. They then caused severe inflammation of the paralysed limb, and noted that the body-heat was raised as much as if the nerves had not been resected. Hence by way of exclusion we are almost forced to accept the doctrine, which for other reasons we have seen to be highly probable, that traumatic pyrexia depends upon pyrogenic compounds absorbed by the vessels from the local inflammation.

So much for the chemical theory of pyrexia; that, just as the heat of the body in health is due to chemical combinations, so the increased heat of pyrexia is due to increased chemical combination. I will now briefly allude to the old mechanical theory. It being believed that the heat of the body in health was due to the friction of the blood, it was naturally supposed that increased heat was due to increased friction, that is to say, to increased velocity of the blood, and increased resistance in the capillaries. The one was corollary to the other. And, no doubt, the immediate cause of some of the febrile heat is in most cases really mechanical; but this is no gain of heat to the body. I will not occupy your time by going over this old ground again.

Let us pass to the other possible alternative cause of pyrexia; namely, diminished loss of heat. No more than the ordinary amount of heat is generated, but less than ordinary is lost: therefore, the temperature of the body rises. The proofs which we have given of the opposite hypothesis, that there is increased generation of heat in all stages of fever, have already refuted by anticipation the doctrine we are now considering. Yet it may be profitable to go shortly over the arguments in favour of the hypothesis of diminished loss of heat.

As you are no doubt well aware, this doctrine has been chiefly upheld by Traube. The lectures unfolding his views were published eight years ago. He begins with a consideration of the cold stage of fevers, in which, he says, the amount of blood in the skin is diminished. This anæmia must be due either to want of power in the heart, or to contraction of the muscular coat of the smaller arteries. But there are no reasons for supposing that the heart is weakened. Therefore, there is a contracted state, a tetanus, of the small arteries of the skin. And, supposing that the arteries of the whole body are in the same condition, Traube believes that he can account in this way for the dry tongue, anorexia, dyspepsia, constipation, diminished secretion of urine, cerebral and other symptoms, generally attendant upon the febrile state. He points out also how, when the vessels relax, and a sweat breaks out, the fever ceases.

Surely it is very curious to note in all this a revival of Boerhaave's doctrine concerning the increased arterial resistance in pyrexia; nay, even of Cullen's reference of that resistance to spasm of the vessels. Of course the supposed vascular spasm is presumed to act differently in the two hypotheses; Cullen supposed it to increase friction, Traube to diminish loss of heat. It is curious also to recal Marey's doctrine of pyrexia, which we formerly discussed, and which, you remember, was based upon the supposition of a paralysis, not a spasm, and of a dilatation, not a contraction, of the arteries in a febrile state. But like Cullen's and Marey's, so also Traube's hypothesis must be classed amongst ingenious attempts to make a process which may be potent to a certain extent, exclusively potent. Leyden's calorimetric experiments have shown us that, even in the cold stage, there is no diminished loss of heat. Doubtless the loss is relatively diminished; relatively, that is, to the increased production: but absolutely the loss is increased in all but the cold stage. I am bound to add that, in the most recent devel-

opments of Traube's hypothesis, the increased production of heat during pyrexia is not denied.

Now that we have reviewed the most important facts which illustrate the nature of pyrexia, let us come to consider it as a cause, and ask if it produce any consequences of its own. We have already found that the combination of tissues which we call the body cannot continue to live under an internal heat of 113 deg. or more. And, in the case of individual cells, Max Schultze and others have shown that the law is the same: death upon a small elevation of temperature. Moreover, anatomical changes are found in the bodies of animals which have died under the condition of mortal elevation of their temperature. But nature does nothing by leaps: we should expect before hand, that temperatures which are unnaturally high, although not straightway destructive of life, would yet exercise a deleterious effect upon the tissues. This is no new notion. Boerhaave's 689th aphorism runs thus: "The augmented heat drives off the most liquid parts from our blood, that is to say, the water, the spirits, the salts, and the most subtle oils; it dries up and condenses the remaining mass, makes it concrete into an impassable irresoluble matter," and so on. Boerhaave's particulars are clearly created at will out of his gross chemical and mechanical pathology; but his general idea has been revived of late years, and supported by careful observation of fact. For example, Liebermeister, who has especially worked at this subject, thus writes: "The longer I study fever, the more I am strengthened in my conviction, that the manifold symptoms which make up the idea of fever are, in great part, mere consequences of one of these symptoms, namely, the elevated temperature. The elevated temperature is not only the pathognomonic symptom of fever, but also the immediate and sufficient cause of most of the other symptoms peculiar to fever, or characteristic of it." We will now consider the facts which are deemed to support this doctrine.

You will remember that, when we were discussing the combustible parts of the body, we found that the heat was furnished in chief part by the glands, the muscles, and the blood; what heat was derived from the nervous system remained uncertain. It is upon these very same organs that pyrexia reacts.

First, with regard to the glands. No doubt it forms part of the practical experience of all who are here present, to know how frequently the cortex of the kidneys is found *post mortem* in a pathological state. In fact, except in the case of perfectly healthy persons who have met with a sudden death, it is very uncommon to see the kidneys free from that change which has been called cloudy swelling, parenchymatous swelling, or albuminous infiltration. Both the glandular epithelium and the intercellular material become charged with minute opaque molecules, which behave like albuminous matters to chemical reagents. In all probability the immediate antecedent of this state consists in an afflux of nutritive material beyond what the organ can use up; either the afflux of blood is increased, or the consuming activity of the organ is diminished, or both of these processes are at work at the same time. Be this as it may, there is little or no doubt that a pyretic state is attended by clouded swelling of the cortex of the kidney, more or less. Not that it is by any means peculiar to pyrexia: on the contrary, local irritation, serious chronic diseases, are accompanied by the same change. But a febrile state is able to produce a high degree of parenchymatous swelling in a few hours; and the degree of the change is mostly proportional to the height of the fever, or to its duration. Function is probably disturbed at the same time; at any rate, the secretion, the urine, sometimes becomes albuminous. A complete return to health may follow the lesser degrees; but when the change has been great, fatty degeneration and disintegration of the tissues ensue. Changes of precisely the same kind take place in the liver. Out of twenty-four fatal cases of typhoid (enteric) fever, Liebermeister found that in eighteen the degeneration of the liver had gone so far as to lead to partial destruction of tissue, and that in the remaining six the degeneration was well marked, although not so far advanced. And, as to the kidneys, in sixteen cases the epithelium was fatty, and in the remaining eight affected with simple albuminous infiltration.

Next, the muscles undergo similar changes; that is to say, they become infiltrated with albuminous and fatty molecules. But, besides these morbid states, you will remember that apparently peculiar condition which Zenker has more particularly studied, and which has been called the waxy or colloid degeneration of voluntary muscle. In half his fatal cases of typhoid fever, Liebermeister found this waxy degeneration more or less advanced. The symptoms which depend, during life, upon these muscular changes, are those so admirably described by bedside physicians under the name of febrile debility. "If that part of the animal actions," says Stoll, "which comprehends voluntary movements, through fever languish so that they are exerted with unwonted difficulty and with a sense of weakness, debility of the animal function is said to be present." But what of the most noble muscle? what

changes does the heart undergo? Softening of the heart in typhoid and typhus fevers has long been familiar to morbid anatomists. In Liebermeister's cases, the heart was fatty in twenty-one, and in the remaining three was affected with simple albuminous infiltration. And the symptoms which attend these degenerations of the heart are those which are classed under the head of vital debility. "If the muscular action of the heart," says Stoll, "the arteries, the lungs, and thereby the circulation of the fluids, be diminished beyond that point which is wont to constitute health, there will be a vital debility, differing in degree, duration, and kind." This is what some persons have called *asthenia*, or *adynamia*. Every fever in which the temperature remains continuously high tends to take on the characters of *asthenia* or vital debility. I will only very briefly allude to these symptoms. First in importance comes the increased frequency of the pulse, which, as a general rule, is proportional to the debility. Moreover, frequency and feebleness of the heart's contractions go together. It has been noted by Cyon and Panum that the contractions of the heart cut out of a frog are greatly influenced by the surrounding temperature: generally speaking, the number of the contractions increases proportionally to the temperature. The irritability of the heart also is more quickly exhausted by a high temperature. The weakening of the first sound of the heart is another sign of the deterioration of the muscular tissue. And the passive congestions and pulmonary oedema, which are so often the immediate causes of death, are further consequences of the vital debility. But perhaps the most striking form of *asthenia* is that which Galen named *lipuria*. "If the external parts be cold, but the internal be burnt up, it is a mortal symptom," says Hippocrates—and with reason; for the debility of the circulation must be great indeed when the blood of the rectum is 104 or 105 deg., and that of the extremities 50 or 60 deg.

Thirdly, concerning the blood itself: upon this topic, chemistry and microscopical anatomy do not give us the definite information which we might wish. The fibrinogenic part of the blood seems to be diminished as a consequence of prolonged pyrexia. And in drops of blood drawn from the finger of febrile patients I think I have found changes in the lymph-corpuscles similar to the changes which we have seen to be present in the case of the glandular epithelium; viz., a granular infiltration of the cells. These granular lymph-corpuscles sometimes seem to be swelled also—a trifle larger than the unaffected corpuscles. At the same time, the blood contains very many free granules, apparently of the same kind as those present in the corpuscles. I confess, for my own part, that it does not appear improbable that these changes are really indicative of a degeneration of the blood, augmented by pyrexia. Any how, the increased metamorphosis which forms part of fever must necessitate the blood being overloaded with the products of those changes. The clinical aspect of the matter is better understood. The symptoms which the ancients considered to be characteristic of what they named putridity are partly due to the change which pyrexia induces in the blood. "The great sign," says Galen, "of putrid fevers is the quality of the heat; for it has nothing mild, nor moderate, nor ephemeral, but, as the best physicians before us have said, it is biting." Translated into modern phrase, putridity is due to a severe fever or a long fever. So Boerhaave, in his 698th aphorism: "The hottest fever is acute, swift, putrid: heat putrefies." And we have not yet forgotten De Haen's vigorous polemics against the heating treatment of fever, which was orthodox in his day, and which he declared, truly enough, to be the cause of all the putrid symptoms, petechiæ and others. Van Swieten endeavours to explain some of the symptoms by the supposition that coagulation of the blood is a possible consequence of the elevated temperature; and this hypothesis receives some support from the declaration of Weikart that the blood begins to clot in the vessels at 109 deg. But the expression "dissolution of the blood" probably comes much nearer to the truth. Liebermeister has shown that parenchymatous degeneration of the liver, however induced, has a strong tendency to be associated with hæmorrhages; and we cannot overlook the great likelihood that degeneration of the coats of the small vessels may be a consequence of pyrexia. But, although we are not yet able to explain the exact nature of putridity, yet the state itself is very real, and the old name, I think, very good. I have often been surprised that the medicine of the present day seems to have forgotten not only the name, but almost the thing signified. I cannot do better than refer once more to Stoll, and on this occasion to his chapter upon Putrid Fever, the first aphorism of which, by your leave, I will quote: "That synochus has been called putrid, which is due to the causes of fever, but greater than usual, and applied longer; with a greater degeneration of the solids and liquids; a more universal degeneration also, swifter, inducing putridity." For the name putridity is not used in a sense metaphorical; but, as matter of fact, the patients sometimes become actually putrid during life, whilst after death decomposition

advances with astonishing rapidity. No doubt some fevers become putrid much more rapidly and with less elevation of temperature than others; so that, although the pyrexia by itself has much to do with the generation of the group of symptoms which we are now considering, yet much must be conceded to the specific nature of the fever—much also to the constitution of the patient. Rheumatic fever is a remarkable example of a severe fever, and a long fever which is, as a rule, singularly free from any disposition to putridity; yet the ardent form of rheumatic fever always passes into putridity. On the other hand, in epidemic diphtheria, two or three days of moderate pyrexia sometimes suffice to plunge the patient into the severest degrees of putridity; the urine becomes loaded with albumen; the blood during life is of a dirty brown colour; uncontrollable hæmorrhage occurs from the mouth and gums; and, *post mortem*, we find parenchymatous degeneration of the glands and heart in the most decided form.

Lastly, what direct influence does pyrexia exert upon the nervous tissues? Now here we are compelled to confess still more our ignorance. The morbid anatomy of the nervous tissue is in a most backward state: so that it is a pure hypothesis (more or less plausible, we may think) to deem that the nervous symptoms which accompany the febrile state are due to changes induced in the nervous tissues by the elevated temperature. Nevertheless, although we do not know in the least of any such anatomical changes referable to pyrexia; yet that it is the cause of much of the disturbed function, would seem to be indicated by the good influence of abstraction of heat over these nervous symptoms. I allude more particularly to the treatment of the so-called ataxic form of scarlet-fever by cold water applied externally: the method introduced by Currie. But the more rapidly fatal cases of scarlet-fever, those which exemplify the true or primary malignity, would seem to show that the elevated temperature alone cannot account for the sudden and complete collapse of the animal functions. When a child in perfect health is suddenly seized with incessant vomiting and purging, delirium and tremors, rapidly falls into a state of complete prostration, and dies in twenty hours, the temperature never rising, even under the influence of convulsions, above 105 deg., it is difficult, I say, to attribute the perfect paralysis of the nervous system to the elevated temperature alone. Yet certainly there is, in many respects, a close resemblance between a case of malignant fever such as I have described, and a case of sun-stroke. Perhaps it is more plausible to see in the paralysis and intellectual defects which follow long fevers, evidences of deleterious change of tissue.

If it be, then, highly probable that pyrexia induces change of structure identical with those consequent upon inflammation, it is hardly less probable that these local changes react upon the body at large in the same way as inflammations act (the pyrogenic power of which we have already discussed)—that is to say, fever produces fever. And the same is probably true of putridity in fevers; it tends to perpetuate and to multiply itself. This opinion is rendered still more plausible when we call to mind the baneful effect which fevers in general, and putrid fevers in particular, exercise over local inflammations. Moreover, the opinion of Liebermeister is certainly very probable, that the unfavourable course which fevers run in the aged, the intemperate, and the fat, is partly due to the degenerations which have already taken place in their tissues—degenerations of much the same kind as those produced by pyrexia.

[To be continued.]

TOXIC ACTION OF QUININE.

A LITTLE time since, I ordered half a grain of quinine as one of the components of an aperient pill for a lady who consulted me. Half an hour after the dose was taken, she was conscious of some irritation over the skin of the head and face, ending in erysipelatous redness and oedema of the right side of the face, eyelid, etc. This lasted more than twelve hours. Since that time, she has kindly sent me a very vivid description of the effects produced upon her by quinine; and I have condensed the following notes therefrom.

"My first recollection of taking quinine is that, when about sixteen, I was weak, and had medicine ordered. The first dose I took at bedtime. I passed a very restless night; and in the morning my whole head was much swollen, and the face so altered that I could with great difficulty open my mouth to take food. The cause was unsuspected. Some time afterwards, I took a glass of port wine from a wineglass in which medicine had been previously taken. But a very small quantity remained in the glass of the quinine solution; yet the same condition of swelling of face, etc., came on, but not so severely. Once again, when recovering from small-pox, I took some quinine, and had large wheals with local redness over neck, chest, and arms. All preparations of bark (*cinchona*) affect me more or less in this way."

W. NEWMAN, M.D. Lond., Stamford.