ated students from other German Universities. The ordinary undergraduate student has, as with us, but little time for extra courses, so that the supply of more advanced special instruction is closely connected with the attractiveness of the school as a whole, which causes it to be visited by learners from a distance, who have already acquired their necessary qualifications. It would be impossible here to discuss the question how such an attractiveness is to be given to the schools of London.

LARGE FLUCTUATING ABDOMINAL SWELLING: RAPID DISAPPEARANCE, WITH EXCESSIVE URINARY DISCHARGE.

By WILLIAM R. LANE, Esq., Surgeon Second Battalion Grenadier Guards.

CORPORAL —, of the Second Battalion Grenadier Guards, was brought to hospital in a cab on December 28th, 1869, at 11.30 P.M. He was on leave, and had fallen down a flight of stairs. On admission, he complained of pain over the crest of the ilium of the right side, the skin over which was slightly discoloured from bruising. There were also pain and tenderness over the region of the bladder. The tongue was white and loaded; the pulse full and rapid. The bowels were constipated. He was ordered to have three grains of calomel immediately, and a dose of castor-oil in four hours.

On December 31st, three days after admission, he was attacked with severe pain and tenderness in the region of the right kidney. He passed a large quantity of blood with the urine. The pulse was very small and rapid. He was ordered to have a grain of powdered opium three times a day. From this date to January 8th, eleven days after admission, the blood in the urine gradually diminished; but on the 10th he complained of a return of pain in the back; and, on examination, there was found enlargement with extensive dulness over the hepatic region. The enlargement gradually increased, distending the right hypochondriac, epigastric, right lumbar, and iliac regions; and, from the swelling pressing upwards on the lung, breathing was much impeded. Fluctuation was distinctly discovered at the back; there was every appearance of a large abscess forming and pointing posteriorly just below the floating rib, and the fluid felt very superficial. Matters remained very much in the same state up to March 1st, and it was then decided that on the following day an exploring needle should be used. On paying my visit to the hospital on the 2nd, the patient said that he felt very much better; and I was surprised to find that all the swelling had entirely disappeared, and that nothing but redness of the skin remained. He stated that the swelling gradually began to subside after I had seen him the day before, and that he passed during the day and night upwards of six quarts of urine. Convalescence was very rapid; and on March 18th he was discharged from hospital perfectly recovered, and has ever since done the whole of his duty.

What was the swelling? and how was it produced? At first it was taken for a large abscess of the liver; but, considering the rapidity with which it disappeared, and the absence of pus in the enormous quantity of urine which passed concurrently with its subsidence, this supposition must have been incorrect. It could not have been urine from rupture of the kidney, or there would have been the symptoms of extravasation of urine—viz., inflammation, suppuration, and sloughing. If it were serum, how did it find its way so rapidly into the bladder? The rupture of large cysts into the peritoneal cavity, and the passing off of their contents by the kidneys after absorption, are not very uncommon, especially in females from rupture of an ovarian cyst. Could this have been an urinary cyst, the urine being deprived of its irritating qualities by the cyst-wall while resident in it? If so, the rupture of the cyst into the peritoneal cavity will account for its disappearance and the enormous quantity of urine passed. The case is so obscure, that I scarcely like to venture a theory to account for it.

PROFESSOR FLOWER'S HUNTERIAN LECTURES ON THE TEETH AND ALLIED ORGANS IN THE MAMMALIA.

Delivered at the Royal College of Surgeons of England.

LECTURE IX.—*Wednesday, March 8th.*

THE Pinnipedia or Aquatic Carnivora are mainly distinguished by the characters of their limbs, which are organised for swimming. The incisors have not the ordinary formula $\frac{3}{3}$, but vary much. The canines are large and well developed. The molars differ from those of other

Carnivora, being very uniform in character, and never having more than two roots. Their crowns are always compressed and more or less pointed, and may have some little accessory tubercles. None of the teeth presents sectorial characters. There are usually five molar teeth above, and the same number below, on each side. The Pinnipedia are divided into several families; viz., Otariidæ, Phocidæ, and Trichechinæ.

The Otariidæ comprehend about twelve species, of which the Sea Lion or Sea Bear (Otaria) is an example. In this animal, the hind limbs are more adapted for walking than those of the Seal; and small external ears are present. There are three incisors in the upper jaw, and two in the lower, on each side. The middle incisors are the smallest; the upper outer ones are very large. Each upper central incisor is divided by a deep groove into an anterior and a posterior cusp. The two lower incisors do not differ much in size; in the young animal, they appear to be lobed in a different direction from the upper teeth. The canines are large and recurved. There are generally five teeth of the molar series, of which the second, third, and fourth are preceded by milk-teeth; the first has no predecessor, nor has the fifth (the true molar). The roots are simple, often connate. The milk-teeth appear at an early period, and are mostly shed before birth. The permanent teeth, in the Otaria and the true Seals, appear to come up at about the same time. In an Otaria about a fortnight old, the milk-canines only were present; a few months later, all traces of the milk-dentition were gone, and several crowns of the permanent teeth had appeared above the surface. In the old animal, the canines become greatly developed through the growth of the roots, and wear away. It is the Otaria that furnishes the article used for clothing known as "sealskin". In another form of Otaria from Behring's Straits, the last molar is separated from the others by an interval. Some Otariidæ have six molars on each side in the upper jaw.

The Phocidæ or true Seals comprehend three groups—Phocinæ, Stenorhynchinæ, and Cystophoridæ. In the Phocinæ, the dental formula is $i \frac{3}{3}, c \frac{1}{1}, p \frac{2}{2}, m \frac{1}{1}$. The upper incisors are sharp and curved backwards, the outer ones being the largest. The canine is large and curved backwards. The molars are all alike; each has a small cingulum, and, except the first, two roots. The crown has a main cusp, with small ones before and behind. The milk-dentition is perhaps even more rudimentary than in the Otaria. In a new-born Seal (Phoca) at the Zoological Gardens, the tops of the permanent teeth were all on a level with the alveolar ridge, and the milk-teeth were nearly absorbed. At an earlier stage, the milk-molars have two roots. In a Seal a week old, scarcely a trace of the milk-teeth was left. In the Phoca barbata, all the teeth become worn down to stumps, until not a fragment of tooth is left. In the Halicæurus or Grey Seal, the teeth are very simple, and are single-rooted—except, perhaps, the posterior molars.

The Stenorhynchinæ are all—with the exception of the Mediterranean Seal—inhabitants of the Southern seas. There are two incisors above on each side, and two below. The molars are simple, and have two roots. In the large Seal of the Southern seas, the molars have a strong lobe, with accessories behind and before. In the Lobodon or saw-toothed Seal, the molar teeth have each a compressed cone with a number of accessory cusps.

The Cystophorinæ or hooded Seals have two incisors on each side in the upper jaw, and one in the lower. The molar teeth are very simple, and generally have each only one root; they have a slight tendency to the formation of a cone with accessory lobes. In the great Elephant Seal of the Southern seas (Macrorhinus), the canines are large and much furrowed on the side. The molars are single-rooted, and have simple slightly compressed crowns, almost like the homodont teeth of the Cetacea.

The Trichechus or Walrus, which is found in the Arctic seas only, has an enormous tooth projecting at each side from the upper jaw. These tusks are massive, and somewhat resemble those of the Machairodus, but have not sharp edges. They are the upper canines. In the lower jaw, the canines do not project; they are not in a line with the other teeth. The skull is specially modified for the attachment of the large tusks. The tusk is composed of true dentine about half an inch thick, covered with about one-tenth of an inch of cement; there is no enamel on the surface. The growth of the tooth is continuous; and, as it wears, the pulp-cavity becomes filled up with osteodentine. There are generally also four round massive teeth, with flat surfaces on each side above and below. They have single roots; the enamel covering the crowns soon wears off. In the upper jaw, the first of these molar-like teeth lies between the maxilla and premaxilla, and is generally regarded as an incisor. In the lower jaw, the first of these teeth may be called a canine, the next three præmolars or molars. In the young Walrus there seem to be more teeth than in the adult. In a young animal, which