You see that in both instances we had to combat disorder of long standing under many unfavourable circumstances, and that success was obtained in each, not by excogitating some new and peculiar remedy, some antidoloric oil or drops, but by the steady pursuance of a rational method, involving simply the use of old well approved remedies. In both cases, improvement came very slowly; some weeks elapsed before any was apparent; but the ultimate results were fairly satisfactory. Now I want you to lay this well to heart in your future course. You will surely meet with many similar cases; and you will be often puzzled and wearied by them, and tempted to give them up in despair; and, if you allow this to appear, the patient's confidence in you will necessarily be shaken. But be prepared for all this discouragement *ab initio*. Let your patients or their friends know that progress, especially at first, will be very slow, and probably interrupted by numerous relapses; but that, nevertheless, the wished-for result is attainable, if they second your perseverance with patience. Remember that diseased action is much like a railway-train in motion; it cannot be brought to a stand-still all at once, though you apply the breaks and shut off the steam. The momentum acquired cannot be safely neutralised *instantly*, and time is essential to set up counter-actions of a restorative character. To those who with fashionable scepticism may be inclined to suggest that Time is the healer in such cases, and not Art, I reply that all my experience leads me to think that the longer space elapses without the intervention of Art, the longer also must be the period during which remedies must be employed. A sciatica of fourteen days' and one of fourteen weeks' standing are vastly different things. Trousseau embodied the same truth in his aphorism—*longue maladie, longue traitement*. In urging upon you perseverance, I may appeal to the estimates formed by foreigners of our national character, from which I trust you do not intend to degenerate. Napoleon sent this message from the field of Waterloo: "I have beaten the English, but they won't go." A Swiss canon, recounting the attempts made at scaling the Matterhorn, writes: "Les Anglais ne reculent jamais, c'est leur caractère." In both these instances, pluck and tenacity of purpose carried the day; and so they often will, under God's blessing, in the strifes which you have to wage against disease and death.

HYDATIDS OF THE LIVER.*

By EDWARD LONG FOX, M.D., F.R.C.P.,

Physician to the Bristol Royal Infirmary.

At the end of May 1869, a girl, aged 14, was brought as an out-patient to the Bristol Infirmary, and was admitted for a short time as an in-patient, and treated for some months afterwards at home.

The mother gave the following account of her. In July 1866, she had an attack of shingles; and whilst she was under treatment for this, an enlargement of the right side was observed by her medical attendant. No further notice, however, was taken of the swelling until Christmas 1868, when it increased very much, interfered with respiration, and was sometimes the seat of acute pain. In March 1869, she was admitted into the Westminster Hospital, under Dr. Fincham, and hydatids of the liver were diagnosed. She remained there for about six weeks, being measured several times a week, but taking no medicine. It is an interesting point of the case that, during her sojourn in the Westminster Hospital, she craved very much for pork, and was clandestinely supplied by her mother with pork and ham.

In May 1869, her parents removed to Clifton, and thus she fell under my notice.

On admission, she presented the following appearance. She was a fairly nourished girl, short, but rather stout. The upper parts of the abdomen and lower parts of the thorax were occupied by a tumour, quite dull on percussion, smooth, and slightly resilient, extending from the fourth rib on each side downwards to the extent of seven inches, causing a considerable bulging of the right side, especially posteriorly. The tumour in front extended seventeen inches horizontally across the body. The right side of the thorax was about two inches and a half larger than the left. The heart was pushed somewhat upwards, the apex beating about the level of the fourth rib. Respiration was rather rapid, especially if she hurried, or walked upstairs, or lifted anything. Digestion was fair. The urine was healthy. She was now menstruating for the first time. The temperature was slightly raised, probably in consequence of slight peritonitis over the tumour. She liked to eat her meat almost raw. Her mother suffered from tænia solium, and did so for the first time when pregnant with this child.

She was under treatment about six months, and month by month gradually decreased in size, until at last no trace of the tumour could

be detected, the diaphragm returning to its proper position. Her medicine all through was iodide of potassium, either alone or in combination with mild tonics. She came to see me a year afterwards, in perfect health.

Besides her recovery, the interesting points in the case are: first, the position of the growth of the hydatid cyst, pushing up against the diaphragm and the ribs, instead of the more yielding walls of the abdomen; and secondly, that recovery took place apparently by the death of the echinococci, from the absorption of iodide of potassium, as it is certain that they were not passed by the stomach, bowels, or bronchial tubes.

Had the patient been a male, the sex in which the diaphragm plays an important part in respiration, the difficulty of breathing would probably have been much more marked.

Frerichs, in advocating surgical measures in cases of this kind, remarks that certain medicines have been employed, which are thought to pass from the blood into the cysts, and to kill their inhabitants. Baumes believed that calomel was endowed with this property; Laennec common salt; and Hawkins iodide of potassium, but as yet (he says) no case is known, in which any such plan of treatment has succeeded.

No doubt, iodide of potassium is often given without success; but I claim this case as an example of absorption of this drug into the cyst. Of course, it may be said that this was an instance of spontaneous death of the parasites, wholly unconnected with the medicines employed. We have in the Infirmary Museum a beautiful specimen of an hydatid cyst of the liver into which a bile-duct discharged itself, and so caused the death of the parasite, and spontaneous cure. But when the disease has advanced as far as in the case before us, spontaneous shrivelling up of the sac is, I believe, unknown. Nor is the other most usual form of cure more common in large hydatid tumours; viz., that in which the acephalocysts are destroyed by the secretion of a thick putty-like material within the sac, a species of caseation.

The iodide of potassium has been used by physicians in this country for many years in these cases. Dr. George Budd mentions it in one of the early editions of his book on the liver, but not with any approval.

CASES OF ALLEGED DISPLACEMENT OF THE UTERUS: WITH REMARKS.*

By C. J. GIBB, M.D.,

Consulting Surgeon to the Newcastle Infirmary.

MR. PRESIDENT,—I would not have promised to lay before you the present cases, had I not felt ashamed to see in the circular calling the January meeting, that, with the exception of Dr. Philipson's valuable Report, no member had a pathological specimen, or even a paper, to communicate to the meeting. It happened that two of the following cases had consulted me that morning; and, the subject being one of great interest, although somewhat out of my line, I thought you would, under the circumstances, excuse my attempting to draw the attention of the Society to it. Three of the five cases have come under my notice during the last month.

CASE I.—A married woman, aged 25, who had had her third child two years previously, consulted me as to the propriety of submitting to the performance of two operations—to set straight, as she said, her twisted womb. She complained of pain under the left ribs, of flatulence, nervous palpitation, and general depression of spirits. She had had all the symptoms at intervals, even before her marriage, and had enjoyed very fair health since her last confinement, until her old symptoms relapsed a few weeks ago. She then consulted a surgeon in her town, whom I know to be emulous of uterine renown. Her complaint proved obstinate in his hands; and a few days before she visited me, he startled her by proposing to perform two operations for the cure of her twisted womb. The word "operation" being rather formidable in her eyes, she came to Newcastle for further advice. I examined her very carefully, and, from her foul tongue, flatulent and distended stomach, pain under the left ribs, and other symptoms, found her suffering from aggravated dyspepsia, dependent very much upon chronic gastritis. Of womb-disorder or displacement there was not a trace. The uterine sound passed up a healthy uterus, in its proper axis, and there was no uterine or ovarian tenderness, nor a trace of any leucorrhœa or of any sexual disorder.

CASE II.—Two years ago, a married gentleman consulted me for chancre. A few weeks afterwards he brought his wife, suffering from a large sloughing chancre on the swollen mouth of the uterus. She

* Read before the Bath and Bristol Branch.

* Read before the Newcastle Medical Society.