peritonitis, creamy lymph is poured out, do bad consequences follow operations, which, though bold, are yet well considered. I saw Mrs. S. in November last. She was then stout and well; and, on examination, the abdomen and uterus were found to be quite free from anything indicative of disease. I have again seen Mrs. S. during the present month, and, except that she is troubled with dyspepsia, she is very well.

NOTES ON DISPLACEMENTS OF THE HEART.

By R. DOUGLAS POWELL, M.D., M.R.C.P.,

Assistant-Physician to the Hospital for Consumption and Diseases of the Chest, Brompton; Physician to the Evelina Hospital for Sick Children.

THE position of the heart* affords us very important information in studying diseases of the chest; for it is particularly constant in health; hence the heart may be said in a manner to serve as an index, warning us of disturbances within the thoracic cavity, and this index should be early consulted in the rapid general survey of the chest which precedes more careful examination.

The great readiness with which the mediastinum becomes displaced has been much insisted upon by Dr. Stokes as of great diagnostic value in pleural effusions; and we see at once the cause of this susceptibility when we remember that the mediastinum is, as it were, *poised* by the contending elasticities of the two lungs; and it is very necessary to bear in mind, that the lungs exercise the same traction power upon the mediastinum that they do upon the thoracic wall, in order rightly to understand the mechanism of cardiac displacements; the physiological bearings, too, of this fact in its relation to the circulation of the blood in the great vessels of the heart are of extreme importance.† This view, however, of the relation between the mediastinum and lungs is entirely opposed to that stated by Dr. Townsend in the *Cyclopaedia of Practical Medicine* (art. "Displacements of Heart"; and which has passed unquestioned up to the present time, though it is demonstrably incorrect.‡ Dr. Townsend, after enumerating the attachments by which the heart is kept *in situ*, states that it is also maintained in position "by the walls of the mediastinum, and by the equal pressure of the lungs which oppose its displacement to either side."

Malpositions of the heart may be conveniently divided into those which are congenital and those which are the result of disease, to the latter alone can the term displacement be properly applied. It is only intended in the present communication to dwell upon a few special points of interest connected with the more uniform and common causes of cardiac malposition.

Displacement of the heart is always a mechanical result either of traction or pressure; displacements by mechanical violence, an interesting case of which is related by Dr. Stokes, are extremely rare, and rather curious than clinically important.

Traction.—This is a more common means by which cardiac displacement is produced than is generally admitted: it is the first displacing force that comes into play both in pneumothorax and in pleuritic effusions: in the former case, when air enters one pleura, causing the lung to collapse, the opposite lung, hitherto also in a state of forced partial expansion, instantly collapses to a certain extent, and carries with it the mediastinum (*Med. Times, loc. cit.*). In early pleuritic effusions, too, the lung collapses exactly in proportion to the amount of fluid effused, and its active elasticity is thus gradually neutralised; in the same gradual manner the elastic power of the opposite lung becomes predominant, and draws the mediastinum to its own side. From the pressure point of view alone, it is not easy to understand how a small quantity of fluid should so readily displace the heart, acting as it must do through the medium of so spongy and compressible a substance as the lung. In those cases where fluid effused into the pleura becomes reabsorbed, but the lung does not return to its former degree of expansion, also in cases of contracted lung from old disease (cirrhosis, the fibroid phthisis of some authors, etc.), the pleural space is filled up by the flattening of the chest wall and the encroachment of neighbouring organs, including the heart, which, with the mediastinum, is displaced towards the diseased side. The mediastinum is regarded by Dr. Stokes and Dr. Walshe as being *drawn over* in these cases; but several other observers, Dr. C. B. Williams, Sir W. Jenner, and others, regard the mediastinum, in common with the chest wall, diaphragm, etc., as being *pushed* towards the con-

tracting cavity by atmospheric pressure. There seem to be two classes of cases to which these two modes of displacement severally apply. 1. In cases of contracting lung without any effusion (cases of cirrhosis, etc.), it is clear that the atmospheric pressure is equal in the interior of each lung; it is also equal within and without each thoracic cavity; it might, therefore, be imagined as removed altogether, and the contracting force of the diminishing lung would alone remain as a displacing cause. The atmospheric pressure in these cases *permits* the healthy lung to enlarge, and to follow the mediastinum. 2. In cases where the pleurae (visceral and parietal) are separated by fluid which is gradually being removed, the conditions are much changed; atmospheric pressure acts equally on every side of this fluid, and tends equally to expand the lung and to depress the chest-wall, to raise the diaphragm and to curve the mediastinum; and the variable success it meets with in these several directions is determined by the varied resistance to its action; viz., by bony resilience, muscular action, membranous resistance, and lung elasticity. Each of these gives way, to a certain extent; in adults, the mediastinum becomes more displaced; in children (especially rickety children), the chest-wall becomes more distorted. Here atmospheric pressure being the displacing force, these cases really belong to the next category, but are conveniently referred to in this place.

In emphysema of the lungs, the heart is displaced downwards by the descent of the diaphragm, which remains, in common with the thoracic parietes, more or less in the position of inspiration, and does not return to the normal arched condition of health owing to the impaired elasticity of the lungs, which prevents their resuming, in expiration, their natural size. The heart is often so completely covered over by the large lungs that no dulness is to be obtained, nor impulse to be felt, except that of the dilated right ventricle at the ensiform cartilage.

Pressure, as a displacing cause, is induced (1) by accumulation of fluid in one pleura; (2) by accumulation of air (or of air and fluid) in one pleura; (3) by accumulation of air or fluid in the pericardium; (4) by the growth of a tumour, thoracic or abdominal.

It has already been observed that the first stage, so to speak, in the displacement of the heart, both by air and fluid, is one of traction, the effusion by annulling the elastic tension of one lung rendering that of the opposite lung free to act. This amount of displacement (an amount by no means inconsiderable) takes place before there is any pressure within the chest. It is instantaneous in cases of pneumothorax, because the elastic air permits the affected lung to collapse at once, and the opposite lung is at once free to act to its fullest extent. In fluid effusion, however, this process takes place gradually step by step, the inelastic liquid only permitting the lung to collapse according to the amount effused. In many cases of pneumothorax (those in which the opening is valvular), and in probably all of pleuritic effusion, further accumulation rapidly takes place, and further displacement of the mediastinum is produced by direct pressure. In cases of reabsorption of fluid effused into the pleura, when the lung is bound down by adhesions, the displacement of the heart is, in common with other organs as above shown, by atmospheric pressure. Before referring to other displacements of the heart, it is necessary briefly to consider to what extent the axis of the heart is altered in these lateral displacements. Thus far the heart and mediastinum, including the pericardium, have been mentioned indifferently, as if the terms were synonymous, because the heart is only affected with and through the medium of the mediastinum and pericardium.

In all the cases of lateral displacement above mentioned, whether to the right or left side, the direction of the heart's axis is slightly altered; in cases of displacement by accumulation of air or fluid, it becomes more vertical, but the axis never diverges beyond the vertical line; the apex never points to the right. In cases of contraction, the axis of the heart is either but little altered or rendered more transverse according to the degree of contraction, since the diaphragm is raised at the same time that the heart is drawn over.

It is difficult to understand how, by any general displacing cause as either fluid effusion into one pleura or contraction of one lung, the apex of the heart could be displaced so out of proportion to the base as to greatly alter the direction of the axis, when the fixed attachment to the diaphragm, the attachment of the right auricle below to the liver, above to the vena cava superior, the other attachments of the heart and its position with reference to the diaphragm and liver, are considered; moreover, on reflecting that the displacing forces under consideration act, as above said, upon the mediastinum and *pericardium*, and only through the latter upon the heart, it appears, I think, impossible, on anatomical grounds, that any pendulum-like movement of the apex of the heart upon its base can ever take place, save within very narrow limits; can at all events ever proceed beyond the vertical line. This is a point of great practical importance; and since it has been stated by observers of the highest authority (Dr. Williams, *Diseases of the Chest*,

* For precise details concerning the normal position of the heart, see Dr. Sibson's *Medical Anatomy*.

† The return of the blood to the heart is greatly aided by the constant action of this force; and in the same cause the tendency to the entrance of air into the veins of the neck during the performance of operations in that region, no doubt, finds its true explanation.

‡ See *Medical Times and Gazette*, January 30th, 1869, art. "Pneumothorax"; also Lecture at College of Physicians, Dr. Salter. *Lancet*, 1865.

4th edit., p. 119; Dr. Sibson, *Medical Anatomy*), that in cases of effusion into the left pleura, or of contraction of the right lung, the apex of the heart passes over until the axis points to the right, the following simple experiment was undertaken with the object of testing the accuracy of the above view.

Six pints of water were injected into the pleura of a female subject, causing displacement of the cardiac dulness to the right of the sternum; a stilette was then introduced at the third right interspace, close to the sternum, and passed horizontally through the chest; a second stilette was then passed in the same manner through the fifth space in a line with the first; the sternum and cartilages were then removed without disturbing the instruments, and the exact position of the heart noted. The upper stilette had passed close to the *left* side of the pulmonary artery at its insertion into the ventricle; the lower stilette had passed through the right ventricle nearer its left than its right border; a line drawn vertically from the middle of the right clavicle touched the border of the right auricle (the extreme right of the heart); a similar line, drawn from the left sterno-clavicular articulation, touched the apex of the heart (the extreme left); the axis of the heart was slightly more vertical than natural.

Mr. Henry Arnott was present at this and at a previous experiment, in which the result was exactly similar, and was kind enough to take diagrams of the exact position of the heart in each case.

In June 1868, a patient died at the Brompton Hospital while under the care of Dr. Pollock. The case was one of extensive excavation from apex to base of the right lung, with thickened pleura and contracted side; the left lung was greatly enlarged, and encroached on the middle line; the cardiac impulse was distinct, both visibly and to the touch, at the fourth right interspace, and no impulse could be felt anywhere else. *Post mortem*, the heart was found to be displaced generally somewhat to the right, but the relative position of the apex and base unchanged, the apex being opposite the fifth cartilage, an inch and a half within the left nipple; the pericardium was completely uncovered on the right side by the retraction of the lung, and the portion of the heart opposite the fourth space was the upper part of the right ventricle. The character of the impulse in this case was indistinguishable from that of the apex beat.

A similar case, one of cirrhosis of the right lung, has been this session brought before the Pathological Society by Dr. Fagge. I have notes of three such cases which have come recently under my observation, and are still living. Dr. Stokes relates an analogous case of displacement of heart, secondary to reabsorption of pleuritic effusion, in which it clearly appears that the impulse on the right side of the sternum was produced by the base of the heart. It thus appears that in cases of displacement of heart, following upon contraction of one pleural cavity, the axis of the heart is but slightly changed (usually, as above explained, slightly more transverse); and that, particularly when the right lung is affected, the displacement often appears to be much greater than it really is, owing to the covering up of the apex by the large left lung, also to its getting more under shelter of the sternum, and the uncovering of the base by the retraction of the right lung.

When the left lung is contracted, the heart is also uncovered as well as displaced; and the pulsation which is felt at the second interspace is produced by the right ventricle at the base of the pulmonary artery; the apex can be felt at the same time below the nipple, or, if the contraction is great, the apex is raised by the ascent of the diaphragm; the axis of the heart is thus rendered more transverse, and the impulse is felt in the axillary region.

Air or fluid within the pericardium causes displacement of the heart backwards, since the attachments of the heart are at its posterior part; in the case of fluid in the pericardium, the apex is slightly raised.

But little can be said, in a general way, concerning the displacements of the heart caused by the growth of tumours, since every case has some peculiarity of its own. Tumour growths, when they approach the mediastinum, usually bear the heart before them in the direction in which they are growing; and this obvious fact is often of great importance in deciding between a lung consolidation and a tumour. There are, however, exceptions to this rule; a noteworthy example of which occurred in a patient of Dr. Cotton's at the Brompton Hospital, in whom there was a cancerous tumour between the right lung and the pericardium, involving the lung and leading to secondary pleurisy, with moderate effusion; the heart in this case was scarcely at all displaced, and this was explained by the fact, ascertained *post mortem*, that the morbid growth which had originated in the bronchial glands on the right side had also affected those on the left, and had invaded the pericardium and also the left auricle, by which means the heart was fixed in position.

Abdominal tumours affecting the position of the heart being mostly of liver origin, give rise to displacement upwards, and to the left. As

before remarked, however, no general rule can be laid down. The aim in practice should be to ascertain, in all cases, the exact position of the heart; and by knowing the results of the more regular displacing causes, which are also by very far the more common, any special peculiarity of position will suggest further inquiry as to some special cause producing it. A continuance of this subject into further minutiae of detail might be considered as an endeavour to elevate what is, after all, but a very important clinical feature of many diseases, into a disease by itself—a mistake most carefully to be guarded against.

CASE OF TRAUMATIC STRICTURE.*

By CHRISTOPHER HEATH, F.R.C.S.,

Assistant-Surgeon to University College Hospital.

THE method of treating stricture of the urethra by rapid dilatation or splitting, introduced by Mr. Holt, has been received with considerable favour in professional circles as regards the ordinary forms of stricture, but I have been surprised to find that many surgeons entertain a doubt whether strictures of a traumatic origin are equally amenable to this form of treatment. Having treated several cases of bad traumatic stricture, complicated with fistula, by this method with the greatest success, I venture to bring before the meeting the last of the series as a fair specimen.

All surgeons are aware of the serious nature of strictures resulting from rupture of the urethra, from falls on the perineum, even if the primary dangers connected with retention and extravasation of urine be safely surmounted. These strictures are extremely dense; in cases of long standing, even cartilaginous in structure, and they, therefore, owing to the non-vascular condition of the parts, are the most satisfactory cases to submit to urethrotomy, whether external or internal. Both these operations, however, possess dangers of their own into which I need not enter now; but I must for a moment institute a comparison between internal division and Mr. Holt's method, in order to mention a fact which has been but recently ascertained. In a patient who was the subject of stricture, and who died of fever before being submitted to operation, I assisted Mr. Holt to split the stricture in the *post mortem* room, and to examine the preparation, which showed a longitudinal rent in the stricture, as may be seen in the illustration to Mr. Holt's work. It was naturally concluded that a similar split occurred in the living subject, though neither Dr. Cruise of Dublin nor myself had been able to discover anything of the kind with the endoscope in patients examined a day or two after the operation. Recently, however, two different surgeons have had the opportunity of examining the urethras of patients upon whom they had operated shortly before, and they found that the mucous membrane was entire, and that merely the indurated submucous tissue or stricture proper was torn. This may not, of course, be the case in every instance; but it is certain that in many Holt's dilator does not tear the mucous membrane, and hence the very slight symptoms produced. In internal division, on the contrary, the mucous membrane must necessarily be incised; and the depth to which the submucous tissue is divided will depend upon the extent of knife protruded, and not upon the full size of the urethra being attained.

In cases of traumatic stricture, complicated with fistula, my experience is that, if the urethra is restored to its natural calibre, the fistulae close spontaneously; but it is possible that severe cases might require a touch with a hot wire to make them heal up.

CASE.—Mr. W. H., aged 18, a midshipman, in June 1867, fell off the wharf at Trincomalee, astride the gunwale of his boat, and ruptured his urethra. He was landed and put under the care of a naval surgeon, who unfortunately died of an apoplectic fit the same day. Being consequently neglected, he got extravasation of urine, and his life was only saved by the arrival, on the following day, of a military surgeon, who made free incisions into the distended scrotum. He eventually recovered, with a fistula of considerable size behind the scrotum. He was taught to pass a No. 8 silver catheter for himself, and continued to do so for a couple of months; but then, being on active duty, unfortunately discontinued the practice.

The patient came to me on May 14th, 1868, in good general health. There was a minute fistulous opening behind the scrotum; from which pus exuded, and considerable thickening about the urethra in the neighbourhood. He passed water in a very small and twisted stream, which was occasionally forked. The urethra was so sensitive that he could not bear me to pass an instrument, and could not pass a No. 3 silver catheter for himself.

On the following day, under the influence of chloroform, I suc-