to ask a physician's opinion as to the possibility of removing her. His answer was, that it was too late; and too late it proved, for she died in a few days."

A system which renders possible such results as these is a cruel irony on charity, absurd in principle, and most disastrous in practice.

ON THE OCCURRENCE OF AMAUROTIC AMBLYOPIA, LONG AFTER THE INJURY, IN CASES OF CONCUSSION OF THE SPINAL MARROW.*

By THOMAS WHARTON JONES, F.R.S.,

Professor of Ophthalmic Medicine and Surgery in University College, London; Ophthalmic Surgeon to the Hospital; etc.

CASE I.—In the early part of last year (1868), my attention was directed to the following case in University College Hospital, under the care of Mr. Erichsen. R. S., a man aged about 35, two years before fell from the top of a house on his back and injured his spine. Since then he suffered from muscular weakness, especially of the right extremities, and was now altogether in a broken down state of body. His mind and spirits were at the same time much depressed. Three weeks before I saw the man, the sight of his right eye had become so dim that he had a difficulty in making out even the largest letters with it. This failure of the sight was accompanied by photops, or an appearance which he compared to "fireworks", before his eye; by haloes all round the gas-lights; and by pains in the region of the right eye and side of the face, extending to the back of the neck. The left eye presented indications of having been, at some former period, affected with iritis. On inquiry, the patient stated that the attack occurred many years before the accident. The sight of the eye in question had not been impaired by the inflammation; but subsequently to the occurrence of the dimness of the right eye, the sight of the left began to fail also, and that in a similar manner. Two or three months before his attention was drawn to the failure of the sight of his right eye, the man found that after reading for five minutes or so the eyes began to water, and that after reading for half an hour he had to give up altogether, in consequence of the occurrence of pain over the eyebrow. The pupil of the right eye contracted freely on exposure to light. Under the ophthalmoscope, I observed in the right eye a bluish whiteness of the optic disc on the side next the temple (apparently on the nasal side), with congestion and blackish discoloration of the retina all round, from granular pigmentous deposit—appearances indicative of degeneration of structure.

CASE II.—In the beginning of June 1868, I was consulted respecting the sight of Mr. O., a gentleman aged 48, who, in November 1866, suffered a shock of the brain and spinal marrow from a railway collision, while in a second-class carriage. Since the accident, the patient's energies, both bodily and mental, had become much impaired. He was unable to read longer than five or ten minutes at a time; and it was only with the right eye that he was able to do this. In November 1867—that is, one year after the injury—dimness of the sight of the left eye was first discovered. On examination in June 1868—that is, eighteen months after the injury—I found that the patient could see with the left eye no better than to make out the large letters called Two-line Great Primer. The pupil was still quite natural in its movements. On making an ophthalmoscopic exploration, I found the optic disc the seat of vascular injection and granular pigmentous deposit, so that it was of a blackish red hue. The retina, all round the disc, presented a similar discoloration. The centre of the disc, where the retinal trunks emerge, was free from vascular injection and pigmentous deposit, but was unnaturally white. Though Mr. O. could still see distinctly enough with the right eye to make out letters of every size, he could not, as before mentioned, continue to read any length of time with it. Under the ophthalmoscope, I found in the right eye congestion and discoloration of the disc and retina similar to what I found in the left, though in a less advanced stage.

CASE III.—Mr. L., aged 31, suffered a shock of the brain and spinal marrow in a railway collision, while in a second-class carriage, on the 4th of September 1866, and was laid up for three months in consequence. Since then he had found himself unhinged, both bodily and mentally. On the 3d of July 1867, ten months after the injury, he first consulted me respecting his eyes, which had been failing him. He complained of a haze which now and then came over the sight, especially that of the right eye; of inability to exert the eyes as formerly; of pain in his head when he made the attempt; of motes which he saw

floating before him; of the appearance, sometimes, as if of flashes of light in the dark; of haloes around the lights; and of undue retention of impressions. Since the patient first consulted me, I have seen him every now and then. Of the several examinations of the eyes which I made, the following is the result of one instituted on the 3rd of July of the present year—just one year and ten months from the date of the accident. Both pupils were active, but when the left eye was kept closed, the pupil of the right eye became somewhat more dilated than when both eyes were exposed to the light. There was some external congestion. The right eye being examined under the ophthalmoscope, the optic disc was seen to present on the temporal (apparently nasal) side a bluish white aspect, and to be opaque and irregular-looking at its circumference all round. Both it and the adjacent retina were in a state of anæmic congestion. Similar appearances, though in a less degree, were observed in the left eye.

CASE IV.—The following is an extract from a newspaper report of a trial at Westminster, on the 25th of June, 1868. The railway accident occurred on the 20th of September 1866—"The plaintiff, about 44 years of age, appeared to be in a very nervous and feeble state. Being asleep at the moment of the collision, he had only an indistinct knowledge of a blow on the head. When he recovered consciousness, he found himself on his knees, leaning over on one side. Since the accident, has suffered a constant pain in the head, with numbness and weakness of the left extremities. *Later, there was the symptom of a shadow before the left eye.*"

In the cases which I have now related, the failure of sight, it is to be remarked, was not experienced until some considerable time after the injury. This would seem to shew that the affection of the eyes arose from a disturbance of the circulation, and consequent impairment of the nutrition, leading slowly to degeneration of structure of the optic nerve and retina.

Inquiry into the nature of the connection between the injury of the spinal marrow, on the one hand; and the affection of the eyes and sight, on the other.—The part of the sympathetic nervous system, on which the healthy circulation in the eye and the due nutrition of the organ depend, has its roots in the spinal marrow in the region of the lower part of the neck and upper part of the back. Thence the nerve fibres pass to the sympathetic in the neck through the hypoglossal nerve and the anterior roots of the two last cervical and two or three uppermost dorsal spinal nerves. From the sympathetic in the neck, the internal carotid plexus arises, and from this are detached fibrils, which, having passed from the cranium into the orbit, enter the eyeball, and are distributed to the muscular walls of the arteries of its internal tunics. Through these fibrils the sympathetic governs the contractions of the walls* of the arteries, and so regulates the variations in the width of their calibre. Variations in the width of the arteries of an organ imply, it is to be remembered, modifications in the flow of blood in the part, independently of the general effect of the heart's action. Thus it is that the healthy circulation in the eyes and certain other parts of the head, and their due nutrition, depend on the integrity of the sympathetic nerves in the neck.

Lesion of the sympathetic nerves in the neck is followed by such a disturbance of the circulation in the eye, and consequently such changes in the nutritive process, as to lead to degeneration of structure and impairment of function.

In the cases which have been related, the roots of the sympathetic in the neck must necessarily have participated in the injury which the spinal marrow sustained from the concussion in the accidents, on which the failure of sight supervened. The effect has been the disturbance of the circulation in the optic nerves and the internal tunics of the eyes, from which the deteriorated nutrition, causing the impairment of sight, has directly resulted.

That the eyes thus suffer from injury of the sympathetic nerves in the neck, the old experiments of Petit, and the more recent ones of Reid, Claude Bernard, Brown-Séquard, and many others have demonstrated. Section of the sympathetic in the neck of a dog, cat, or rabbit, in the experiments referred to, was followed by vascular congestion and disturbed nutrition of the eye of the same side, leading, in some cases, to destruction of the organ by penetrating ulceration, or even sloughing, of the cornea, and evacuation of the humours.

The occurrence of inflammatory congestion of the eye after section of the sympathetic in the neck was, in the first edition of my work on *Ophthalmic Medicine and Surgery*, attributed to the consequent paralysis of the walls of the blood-vessels of the eye; and microscopical observations recorded in my essay on *the State of the Blood and the Blood-vessels in Inflammation*, published in 1850, shewed, in illustration and cor-

* The radiating muscular fibres of the iris are also under the government of the same part of the sympathetic nerve, being supplied with nerve-fibrils therefrom. But this is a point which does not immediately belong to our present subject.

* Read at the Annual Meeting of the Shropshire Scientific Branch, October 1868.

roboration of this view, that section of the ischiatic nerve in the frog, which contains, mixed up with the ordinary sensitive and motor fibrils, the sympathetic filaments which are distributed to the arteries of the limbs, was followed by dilatation of the arteries, with a fuller and more rapid circulation in the web. The blood in the capillaries and veins being, at the same time, unusually loaded with red corpuscles, the general effect to the naked eye was increased redness, not only of the web, but of the whole limb.* An eventual result was opacity of the web, indicating an altered state of nutrition.

In the cases related in the first part of this paper, the eyes, though seriously impaired in function, have not suffered disorganisation such as was observed in the animals subjected to the experiment of dividing the trunk of the sympathetic in the neck. In the latter cases, the influence of the sympathetic on the circulation in the eye was completely cut off. In the former, the injury to the sympathetic, from the concussion of the spinal marrow, has been such only as to impair and pervert its influence on the circulation in the eye. It is to be remarked that the subjective symptoms which first attract notice in such cases may exist for some time without any material alteration of structure being distinctly observable under the ophthalmoscope. It is also to be remarked, in conclusion, that the inability which the patients laboured under to exert their sight for any ordinary length of time, which accompanied the amaurotic failure of sight, was different from the common form of asthenopia. The inability to exert the sight was owing, partly, to the irritable congested state of the eyes, and was, partly, one manifestation, among others, of the impaired energy of body—sense and mind generally—which we saw to be a characteristic of all the cases which have been passed under review.

TWO FATAL CASES OF CONVULSIONS OF A PECULIAR CHARACTER.

By THOMAS DALTON, M.D., Llandudno.

I VENTURE to send for publication a short record of two cases of convulsions which have come under my observation during the last three years, as they seem to me to possess several features of interest: 1, in the suddenness of the seizure; 2, in the nature of the convulsions; and, 3, in the apparent uselessness of all remedial measures, at least of such as were tried. As the cases are very much alike in their leading characteristics, I propose to give the history of one somewhat in detail, and then to mention any important points of difference observed in the second case.

CASE I.—The patient was a boy aged nine. He was small for his age, but appeared to have enjoyed uniform good health. He had never been known to have convulsions of any kind. His parents were poor, but he had been adopted by the persons with whom he was living at the time of the attack. He had been with them for about three weeks, and during that time had eaten voraciously of food to which he had not been accustomed—chiefly meat and pastry. He had appeared to be in perfect health till the day I was sent for. He went to bed quite well the night before, but about four o'clock in the morning he awoke complaining of great pain in the stomach; after having rather a large dose of brandy and water, he went to sleep again; about nine he awoke, and almost immediately afterwards “had a fit.” When I arrived, the patient was in bed, and was just beginning with a convulsion, apparently of an epileptic character. He was immediately put into a warm bath with some mustard in it. The convulsion soon passed off; the patient was put to bed again; a bladder of ice applied to the head, and a powder of calomel and jalapine given. When I called again, in a little more than an hour, I was told that the convulsions had returned, and that they came on very frequently. I now saw that they had changed in their character, for during the attack the patient presented very much the appearance of one suffering from tetanus, save only that he was insensible both during the attack and the interval. After the paroxysm passed off, he remained quiet for about ten minutes, and during that time the pupils were natural and quite sensible to light; he then began to breathe quickly (the respirations often amounting to 50 per minute); the pupils became fully dilated and the eyes fixed; there was slight mus-

cular twitchings of the arms and legs; next, the sterno-mastoid muscles became rigid; the rigidity soon extended to the muscles of the limbs and back, so that there was complete opisthotonos; the arms and legs were semiflexed, and the thumbs bent in on the palm. There was no foaming at the mouth, but the tongue was protruded, and the jaws clenched so firmly that it was with the greatest difficulty any thing could be kept between the teeth. During the paroxysms, respiration was almost in abeyance, there being merely an occasional sobbing gasp. This state of spasm usually lasted from one to two minutes, then it went off quite suddenly, and all the muscles were almost instantly and perfectly relaxed. Though unconscious, the patient was very susceptible of any noise or motion in the room; in fact, a paroxysm might at any time be induced by merely shaking the bed, or still more certainly by making any attempt to get him to swallow. The symptoms continued much the same till death, which took place about ten o'clock next morning, or about twenty-four hours after the first attack. I should mention that the bowels were freely moved during the night, and that for the last two hours of his life there was a very loud churning sound in the abdomen during the paroxysms. A *post mortem* examination was made the same afternoon by Dr. Nicol and myself. We examined the head, thorax, and abdomen, but not the spinal cord, as the friends, unfortunately, objected. All the organs appeared perfectly healthy. The only approach to any morbid appearance was in the brain, the whole substance of which was somewhat soft and friable, but there was no true softening of any particular part—no congestion and no effusion. On examining the stomach and intestines, we found that they contained a very large quantity of bile; in largest quantity in the small intestine. There was not a trace of any undigested food or feculent matter; in fact, the bilious fluid formed the sole contents of the intestines.

CASE II.—In this case, the patient was a little girl aged eight; not naturally robust, but generally enjoying fair health. She had been languid and heavy the day before, but in the evening seemed better. When I first saw the child, she was cold, pale, and collapsed, with a weak laboured pulse of only forty. She had been put into a warm bath before my arrival, and I was told that the fit had lasted about five minutes. I had her put to bed immediately, and gave her a little brandy and water, which she swallowed; the pulse soon improved, and she presented the appearance of a child after an ordinary epileptic fit. I then left, promising to call again in three or four hours. On my return from visiting a patient in the country, I found that my partner, Dr. Nicol, had been called to the patient, as the convulsions had returned. The pulse was now 120, and the convulsions of a different character. It struck both of us that the case was likely to prove similar to that just narrated. Our fears turned out to be but too well founded, for the disease followed almost an identical course both in its progress and results, except that in this case the child lived longer (forty hours) after the first attack.

I must now say a few words regarding the treatment adopted. In both cases we tried warm baths in the first instance, then ice to the head and spine (by means of Chapman's bag); at first this seemed to be of use, especially in the first case, but it soon lost its effect, and indeed, after a time, it seemed to do positive harm. We next had recourse to chloroform, and certainly, as a *palliative*, with most satisfactory results. If administered in time (*i.e.*, as soon as the breathing began to be hurried), it entirely prevented the spasm, but if a little longer delayed, it seemed to hurry on the attack, and to render suffocation more imminent; in fact, the symptoms it produced were at first rather alarming, but we found that, when persevered with, the spasm was overcome in a few seconds, and breathing resumed quite naturally. It was a remarkable fact, that in the second case the pulse invariably became stronger, slower, and more regular, whenever the chloroform was given. The administration of the chloroform was kept up in this case, by Dr. Nicol and myself, without intermission, for sixteen hours, about five ounces of chloroform being used. During this time there were only half a dozen decided spasmodic attacks. However, greatly to our disappointment, as soon as it was given up, the attacks returned just as frequently and violently as before. We next determined to try the subcutaneous injection of atropine, still giving the chloroform when a paroxysm was threatened. The first time I injected about $\frac{1}{10}$ of a grain; but, as this had no apparent effect, in about two hours I injected again, increasing the dose to about $\frac{1}{5}$ of a grain. This seemed to do some good, for the patient soon became very quiet, the breathing regular, and there was no sign of spasm for upwards of three-quarters of an hour. At the end of that time the spasms returned; and though the injection was again tried, it entirely failed. I should mention that in this case (the second) the bowels were relieved by means of an enema, the long tube being used. The motion was copious, dark, and offensive; afterwards the strength was kept up by means of enemata of beef-tea and port wine.

* In 1852, Dr. Claude Bernard, in repeating the old experiment of cutting the sympathetic nerve in the neck of a cat or rabbit, discovered that, in addition to the effects on the pupil and on the circulation in the eye previously known, there supervened redness and heat of the ear and side of the head. Dr. Brown-Séquard, having cut the sympathetic in the neck, galvanised the end of the nerve above the section, and found that the increased redness and heat of the ear and side of the head were thereby diminished. This effect was owing to constriction of the arteries, in consequence of their circular muscular fibres having been excited to contraction by the stimulation to which the sympathetic was subjected by the galvanism.