various stand-points, and in the interpretation of her language by different minds, that our store of knowledge is extended; and he who works in an honest spirit, even though he fail to read her lessons aright, may still have done some service in helping forward the cause of Truth.

ADDENDUM.

THIS woodcut, from a drawing by Dr. Stewart, the Curator of the Museum at St. Thomas's Hospital, exemplifies the condition alluded to in a former Lecture, in which the primitive muscular fibres are found

ecchymosed, disintegrated, and ruptured, in tetanus. The specimen was taken from the patient referred to in the text, who died of acute traumatic tetanus; and is magnified 300 diameters.

ON THE USE OF PROTOXIDE OF NITROGEN GAS; AND ON A NEW MODE OF PRODUCING RAPID ANÆSTHESIA WITH BICHLORIDE OF METHYLENE.*

By RICHARD RENDLE, Esq.

It is my intention to place before you the results of my experience in the use of protoxide of nitrogen gas and the bichloride of methylene, which, I think, from the variety of cases in which these agents were used, and the varied modes in which they were administered, may prove interesting to you. Having had, since the last meeting of the Association, some opportunities of studying the use of anæsthetics in various departments of surgery, I have been often struck with many disadvantages in the use of chloroform; such as the length of time in the production of anæsthesia, the occurrence of vomiting and muscular excitement, and the tardy recovery which in some cases attended its use.

With the idea of lessening, if not of remedying these evils, I began to administer the bichloride of methylene with a modification of Dr. Junker's apparatus, which is here before you. I gave it in this way in forty cases taken indiscriminately—19 males and 21 females, from 14 months to 76 years of age. In 32 of the cases, the time required to produce anæsthesia was—over 20 minutes in 3; over 10 minutes in 4; over 5 minutes in 9; over 3 minutes in 6; over 2 minutes in 2; and over 1 minute in 8. In eight cases, all over 20 years of age, after continuing the administration for 3, 5, 10, 11, 14, 15, 16, and 17 minutes respectively, using from 30 minims to 4½ drachms, chloroform was substituted, and in 5 of the 8 produced anæsthesia in 1 minute; in 1 in 2 minutes where the methylene had been used 14 minutes; in 1 in 3 minutes where the methylene had been used 15 minutes; in 1 in 9 minutes where the methylene had been used 16 minutes; in this last case, complete muscular relaxation was required and produced. Of the 32 cases, in 1 anæsthesia was kept up for 45 minutes (ovariotomy); in 2, for 10 minutes; in 9, for 5 minutes; in 15, for 2 minutes; in 5, for 1 minute. On an average, three or four fluid-drachms were used. It was given for iridectomy in 11 cases; for strabismus in 11; for extraction of cataract, needle-operation for cataract, excision of eyeball, applying nitric acid, setting fractures, reducing dislocations, ovariotomy, and various other operations, in 18. Vomiting occurred in 8 cases—that is, 1 in 5. Thus, in the majority of the cases, it took longer to induce anæsthesia, required more of the anæsthetic, and was consequently much more expensive (bichloride of methylene being at present more than three times as costly as chloroform). Vomiting occurred as often as with chloroform; and the only advantage which I noticed the bichloride of methylene, administered in this manner, to possess over chloroform, though a very important one, was, that there was scarcely any muscular excitement.

I next gave the protoxide of nitrogen gas in 74 cases. The details of manufacture and of administration having been published in the *Odon-tological Society's Transactions*, and in papers read by Mr. Fox and Mr. Clover at Oxford last year, I shall proceed at once to mention the modes in which I have given it, and the observations I have made. Its mode of action, I believe, is similar to that of all other anæsthetics; viz., that the blood supplied to the nerve-organs is not duly oxygenated,

or is not able to interchange its oxygen for carbonic acid sufficiently for the due performance of their duties. Anæsthesia is a consequence of the air supplied not producing the necessary changes in the nerve-organs, and not of a deprivation of respirable air, as has been stated.

At first, I obtained the gas compressed in iron bottles from Mr. Barth, and gave it in the manner recommended by Mr. Coleman, which consists in causing the gas, when expired, to pass through a chamber of slaked lime (as shown in this diagram). This economises the gas very much, about three gallons being sufficient for each patient for almost any length of time, but the apparatus required is rather complicated; moreover, there must be some arrangement for getting rid of the air already in the lungs, instead of allowing it to return into the bag. This is effected by a valve near the mouth-piece, which opens on expiration, and after the first three expirations can be closed. Here are two made for this purpose. After this the gas, when expired, returns to the bag over the slaked lime, by which any carbonic acid is removed, and the gas is rebreathed. It was given in this way in 24 cases, 13 males and 11 females, from 3 to 73 years old: for strabismus, in 7 cases; for iridectomy, in 6; for operations on the eyelids, in 8; removal of opaque lens capsule, in 1; extraction of teeth, in 2. The shortest period in which anæsthesia was produced was 60 seconds; the longest, 150 seconds; average, 93 seconds. Anæsthesia was maintained 13 minutes in 1 case; 6 minutes in 5; 4 minutes in 5; 2 minutes in 12; and 1 minute in 1. Intervals of air were allowed in all but one. The gas was given until deep anæsthesia was produced, air being then admitted until the natural colour returned, when the gas was again turned on, and so on alternately as long as desired. The gas was required for a shorter period at each successive time. The period during which air was admitted varied very much, and it would require a much larger experience to state whether it had any relation to the time of each inhalation of gas or to the whole period of anæsthesia. In the earlier cases, before I succeeded in getting well-fitting mouth-pieces, some failures occurred from the admission of air with the gas; yet I noticed that, if the quantity of air admitted were small, anæsthesia was produced, though less rapidly, and the blueness was less marked. One of these patients laughed uproariously on recovery, and had lost all idea of the time. One, a woman, aged 48, on whom iridectomy was performed in both eyes, was under the influence in 60 seconds. The gas was continued five minutes longer without any admission of air; at the end of this time she was most unusually blue; the respiration and pulse very slow and failing. The gas was removed; the patient was turned slowly over on her left side (a proceeding, I believe, peculiar to the ophthalmic department at Guy's, and which has never failed to restore cases manifesting unpleasant symptoms from anæsthetics). She gradually recovered completely, and walked away in five minutes more. I next had a small gasometer made, a modification of one made by Mr. Barth (seen in section in this diagram); it has a core to diminish the quantity of water, and on the top of this is a tray containing slaked lime; a pipe passes in from beneath for the passage of the gas in and out; the upper part is exactly counterbalanced, and, as the gas is breathed out and in, this falls and rises. The expired gas is thus exposed to the slaked lime before being again inspired. Of course the same arrangement has to be made in the mouth-piece for getting rid of the air in the lungs, by the first three expirations being allowed to escape instead of being expired into the gasometer, but all other valves are done away with. This is a very suitable apparatus for an operating-room, but is not portable. This was used in 15 cases—7 males and 8 females, from 4 to 70 years of age; for strabismus, in 3; for needle operation for cataract, in 1; for operations on the nasal duct, extraction of teeth, removal of toe-nails, and various others, in 11. The shortest period in which anæsthesia was produced was 30 seconds; the longest, 85 seconds; average, 64 seconds. Anæsthesia was maintained half a minute in one case; 1 minute in 8; 2 minutes in 3; 4 minutes in 1; 6 minutes in 1; and 8 minutes in 1. One of these cases, a man, aged 30, short-necked, plethoric, who began to inhale it for the extraction of a tooth, pushed away the inhaler and refused to continue. He complained of very unpleasant sensations in his head for some hours after, but this eventually passed off. One woman complained of headache afterwards.

Having now a store of gas at my command, economy was not studied, and it was given in the ordinary way from a large bag, with a mouth-piece having an inspiratory and expiratory valve; in this way a much larger quantity of gas was used, about five gallons for each minute's inhalation. Thirty-four inhaled it in this way—22 males and 12 females, from 3 to 58 years of age; for strabismus, 3; enlarging lacrymal duct, 2; fistula in ano, 6; opening of abscess, 4; extracting teeth, 8; removal of nails, reducing paraphimosis, removal of small cysts, attempt to reduce a dislocation, 5. The shortest period in which anæsthesia came on was 40 seconds; the longest, 100 seconds; the average, 61 seconds. Anæsthesia was maintained 1 minute in 23; 2 minutes in 6; 3 minutes

* Read in the Surgical Section before the Annual Meeting of the British Medical Association in Leeds, July 1869.

in 4; and 5 minutes in 1. One had had chloroform and bichloride of methylene on previous occasions, but she preferred the gas. Three had a second inhalation. In the case of dislocated humerus where it was used, the patient, a strong muscular man, was brought fully under the influence on two occasions; but, though considerable force was used as the muscles remained rigid, we failed to reduce it. Chloroform was given about twenty minutes afterwards, and the dislocation easily reduced.

A medical student, aged 22, had it for the extraction of an impacted molar tooth four times in succession on one occasion, and twice on another. Each time, immediately on the removal of the inhaler, before any instrument was used, he gave a shrill scream; and, on recovery, had no knowledge of it, but described the sensation as pleasant. A child four years old had it on two occasions, and, on the second, vomited about two hours afterwards. This is the only case in which vomiting followed the use of the gas. In a child three years old, where anaesthesia was maintained 90 seconds, two intervals of air having been allowed, the respiration ceased, and the pulse sank so as to be scarcely perceptible at the radial artery for several seconds. The gas was removed, and the child turned slowly over on his left side. He gave a deep sigh; the pulse and respiration gradually returned; and he recovered completely within five minutes.

There was no appreciable difference in the effects under the three modes of administration, excepting a slight one in the time of induction of anaesthesia; being, on an average, 93 seconds with the small bag; 64 seconds with the gasometer, where the same gas was re-breathed after passing over slaked lime; 61 seconds with the large bag, where it was breathed only once. This difference, however, appears greater than it should, from the average being much raised in the first mode by badly fitting mouth-pieces. I generally prefer the large bag for use in private practice, being less complicated and more portable, the extra expense of gas being of minor importance.

It may be said, then, that the advantages of the gas are, the rapid production of, and recovery from, anaesthesia; the absence of sickness, etc.; and the agreeable taste. That it is safe in all short operations, there can be no doubt; and, I think, also in many longer, for even twenty minutes or more, provided there be a due admission of air at proper intervals. But that great care is required in its use, is shown by the alarming symptoms of one or two cases which I have related.

Its disadvantages are, the trouble and time required in its preparation, which I know to my cost, having occasionally had to remain watching it through the night, though this has been much lessened by the beautiful automatic apparatus invented by Mr. Porter; its bulk and non-portability, even when compressed into iron bottles; the complex apparatus necessary for its administration; its unsuitability in certain cases, especially where muscular relaxation is required. The rigidity and congestion preclude its use for operations within the eye where there is any increased tension. The muscular twitchings interfere with all delicate operations. Moreover, it has the objection common to all gaseous anaesthetics, as pointed out by Dr. Richardson; viz., that they remain gases at the temperature of the blood, and do not condense when introduced into it. Hence it is necessary to give it in very large quantities, to produce anaesthesia; and this renders its use expensive. But, notwithstanding all these disadvantages, some of which may yet be overcome, it is very useful in certain cases, especially dental.

I must now pass on and describe the new method of giving bichloride of methylene, by which effects are produced resembling, in rapidity and absence of unpleasant symptoms, those of the protoxide of nitrogen gas, and which are free from some of the objections to the gas. It was suggested by Mr. Bader, who had seen it so used by a dentist in Brighton, that, if I could succeed in making the patient inhale a large quantity of the bichloride in a short space of time, anaesthesia would be rapidly induced. For this object, I constructed a cylinder of cardboard, seven inches long, and about three in diameter, perforated at the sides, and covered inside and out, and at one end, with some absorbent yet porous material. The other end was left open, and shaped to fit closely over the nose and mouth.

I have had several modifications made, in size, shape, and material. The last are those exhibited in the Museum, and these on the table before you. They are made in various sizes, of leather sufficiently thick to retain the shape, yet thin enough to yield a little in fitting on the face. The top is dome-shaped, and perforated to admit just sufficient air to enable one to breathe without effort. The sides are not perforated, and the open end is shaped to fit nose and chin. In the interior is a flannel bag, the mouth of which is turned over the edge of the leather, and secured by an elastic band. Thus the edge is made soft to the face; and the flannel lining is kept in position, and, when soiled, can be readily changed. Into this inhaler, for an adult, one fluid-drachm of the bichloride of methylene is sprinkled, and the inhaler applied closely over the

nose and mouth: the patient is directed to breathe freely, anything tight around the chest or neck having previously been loosened, and the respiration and pulse being closely watched. In some cases, the respiration proceeds naturally; in others, chiefly from fear, there are convulsive efforts at inspiration for a few seconds. This generally ceases, and respiration becomes natural. If not, it is advisable to remove the inhaler, and allow one inspiration only; and then reapply it; and all goes well. But all unnecessary admission of air must be avoided, as the rapid effects are dependent on the rapid inhalation of the bichloride with a minimum of air.

The pulse is invariably strengthened and accelerated at first, but soon returns to its normal state. Insensibility to pain comes on, in a large number of cases, in twenty seconds; and Mr. Bader generally commences ophthalmic operations in about thirty seconds. I usually prefer, however, in general surgery, continuing the inhalation for about sixty seconds; but even then I know no one sign indicative of insensibility to pain, as distinct from complete insensibility; the former, as must be well known to many, being a prior stage to that for which a callous conjunctiva is the guide.

The duration of anaesthesia also varies much. In dental operations, where the inhaler has of necessity to be removed, and the pain is peculiarly acute, anaesthesia is short, though quite long enough for one, and often for three or four extractions, according to the skill of the operator. In other operations, where the inhalation can be continued until the drachm of bichloride is exhausted, the effects last about five minutes, and then the patient recovers, and is able to walk away with slight unsteadiness in gait, which passes off in a minute or two, and recovery is perfect. No sickness or headache follows, unless the inhalation have been continued many minutes, or a second dose is given to keep up the effect. Of course, if the inhalation be prolonged, the after-effects resemble, though in a less degree, those of chloroform, save the one before mentioned—absence of muscular excitement.

The following is an analysis of 123 cases in which I have given it in this way. The ages varied from 6 months to 70 years. It was given for iridectomy in 40; for operations on eyelids and ducts in 9; for strabismus in 8; for extraction of cataract in 6; for needle-operation for cataract in 1; for excision of eyeball in 2; for various other ophthalmic operations in 11; for extraction of teeth in 18; for examining and setting fractures in 9; for applying nitric acid in 7; for opening abscesses in 4; for reducing dislocations in 2; for circumcision, excision of testicle, removal of growth from hand, amputation of finger, dividing fistula, in 6.

Anaesthesia was produced in 30 seconds in 18 cases; in 60 seconds in 70; in 2 minutes in 25; in 3 minutes in 5; in 5 minutes in 3; in 9 minutes in 2.

Anaesthesia was maintained 1 minute longer in 20; 2 minutes in 31; 3 minutes in 19; 4 minutes in 14; 5 minutes in 11; 6 to 10 minutes in 10.

Less than 1 fluid-drachm was used in 20 cases; 1 fluid-drachm in 54; an additional half-drachm in 20; an additional drachm in 16; additional 2 to 4 drachms in 13.

Fifty recovered within 1 minute; 23 in two minutes; 9 in 3 minutes; 28 in 5 minutes; 11 in 10 minutes; 2 were continued with chloroform.

Vomiting occurred in 15—*i. e.*, 1 in 8; but in all of these it was continued beyond the second minute; and in 9, more than 1 drachm had been used. Some had just been eating.

Three female adults and one child became rather blue, and the pulse slower than normal; but, on removal of the inhaler, and turning them slowly on the left side, they recovered well and rapidly. One, a man aged 70, spat up a little bright frothy blood during the inhalation, and once or twice after, during the next twenty-four hours. He stated that he had never done so before. One woman became very hysterical. Two had had the gas, and liked methylene equally well. One preferred the gas. One screamed while under, but had no knowledge of any pain.

The rapidity of action and recovery appears to be due to the great volatility and solubility enabling a large quantity to reach, and escape from, the nerve-organs at once; its safety, to its rather stimulant action on the heart, and its rapid elimination. The bichloride, being very volatile, requires to be kept in a well stopped bottle; and it is an advantage to keep it in the dark, inverted under the water.

To sum up, then, I claim for the new mode the following advantages. The patients retain their normal colour and appearance. The anaesthesia is good, is rapidly induced, and can be maintained for any length of time. The recovery is rapid and complete; there are no unpleasant after-symptoms, nor muscular rigidity. The apparatus required is simple and portable. As regards safety, I may add that I have not had a fatal case.

I hope, gentlemen, the observations I have placed before you have been sufficiently novel and interesting to repay you for the attention you have given me.