decrease. The wasting of large overgrown tumours, dating from a serious illness, is a case in point. The shrinking of large fibroid tumours of the uterus, after boring and internal bruising of their structure, as employed by Mr. Baker Brown, seems to me to be consequent upon the production of internal cicatrices, which so far interfere with their freedom of blood-supply as to initiate the wasting process. The subcutaneous crushing and breaking up of tumours (*Cyclopædia of Surgery*, vol. iv, p. 403), employed by French surgeons in some cases with success, is another instance in which internal—i.e., subcutaneous—destruction of a portion of a tumour has induced the wasting of the whole. One case is thus described (*Cyclopædia of Surgery*, loc. cit.): "I saw Thierry treat in this manner a lipoma of the mastoid region in a Paris banker, who was anxious that the operation should leave behind no visible cicatrix. The tumour was no larger than a small pullet's egg, yet it required for its crushing three sittings, at intervals of three weeks. Resolution took place at last, and, after a lapse of six months, the tumour had completely disappeared, but Thierry entirely failed in another case, etc."

In all these instances we see much to encourage us to inquire whether we may not so plant—if I may use the word—cicatrices around or in the interior of tumours which we cannot, or prefer not to extirpate, as to determine the balance of nutrition against growth, and in favour of atrophy. May we not hope, like the French surgeon, to be able to cause the disappearance of tumours without the production of visible cicatrix?

In conclusion, let me say that, should any readers of this paper have met with facts which seem to confirm the views here expressed, I shall esteem it a favour if they will kindly communicate them.

ON LOCOMOTOR ATAXY.*

By J. LOCKHART CLARKE, M.D., F.R.S., etc.

THE effects on the voluntary control of muscular movements produced by dividing the posterior roots of the spinal nerves, as shown in the experiments of Claude Bernard, Brown-Séquard, Leyden, and others, are of great importance in reference to the muscular incoordination of locomotor ataxy. (*Leçons sur la Physiologie et la Pathologie du Système Nerveux*, 1858, tom. i, p. 261.)

In one of his experiments on a frog, Claude Bernard cut the posterior roots of the nerves that supply both the anterior and posterior limbs. When placed in water, the animal remained quite motionless. If it was excited by pricking its head, which retained sensibility, it executed certain movements with all its limbs; but these movements were of a disorderly character, and not at all in that harmony with each other which is necessary for coordinate movements, such as swimming. In order to ascertain whether these disorderly movements were due to cutaneous anæsthesia, or to the loss of muscular sensibility, Bernard stripped off the skin from all the limbs of another frog, in which the nerves had not been divided. When placed in water, the animal moved with all its usual agility, and swam in its accustomed manner. Moreover, Bernard divided the cutaneous nerves of a sparrowhawk and of a dog, without producing any disorder of the voluntary movements.

On another occasion, he cut the posterior roots of the lumbar nerves on the right side in a young dog. The left leg executed perfectly the movements of progression; while the right leg, which was rendered insensible by the operation, was dragged along after it, and a little raised and somewhat bent. In another dog, he divided two or three of the posterior roots on the right side in the lumbar region. The animal was no longer able to support itself on the corresponding leg, which was in a state of semiflexion. Nevertheless, movements of an apparently involuntary character occurred in that leg whenever the animal moved. Moreover, we are told that the limb, which was completely deprived of its sensibility, was reduced to extreme weakness.

The experiments of Rosenthal and Leyden, with regard to the effects of division of the posterior roots on the voluntary movements, tend to the same general conclusions as those of Claude Bernard. There are two experiments, however, that deserve especial notice. In one of these, three posterior roots were divided in the lumbar region on the right side, and only one root on the left side. The right leg became paralysed, while the left leg retained its power. When the animal was placed in water, the left leg assumed its usual position, while the right leg hung extended and inert. In another frog, two posterior roots were divided on the left side only, in the lumbar region. When the corresponding limb was forcibly extended, it remained in that position,

unless the opposite limb was excited. The act of jumping was awkwardly performed, because the limb that was deprived of its sensibility was not drawn up.

Let us now turn to some similar experiments performed by Brown-Séquard. In 1855, this distinguished physiologist published an account of the effects observed in several cases after dividing a variable number of the posterior roots of the spinal nerves. (*"Recherches Expérimentales sur les Voies de Transmission des Impressions Sensitives," etc., Mémoires de la Société de Biologie*, 1855, p. 31 et seq.) The most important of these experiments are the following.

EXPERIMENT I.—In a guinea-pig, rabbit, or dog, he cut all the posterior roots of the five or six last dorsal and two first lumbar nerves on the right side. After allowing the animal to rest for some time, he found that the power of voluntary motion was diminished in a remarkable degree in the right hind leg, but that its sensibility was exalted, while, on the contrary, it was diminished in a very evident manner in the left hind leg. Moreover, he found that the blood-vessels were rather dilated in all parts of the body behind, and on the same side as, the section of the nerves; and that the temperature of those parts was raised from one to three degrees above that of the corresponding parts of the opposite side. These are the same effects as he had long since found after section of one lateral half of the spinal cord in the dorsal region. Immediately after section of the nerves, the phenomena just described are strongly marked; so that sometimes the paralysis of voluntary movement seems to be complete; but, after some hours, these effects diminish.

EXPERIMENT II.—When the same roots were cut on the opposite side, sensibility and voluntary movement were retained only in a slight degree in the posterior extremities. At the end of eight or ten hours, the animal could execute tolerably strong voluntary movements, but was still unable to support itself on its hinder extremities.

EXPERIMENT III.—All the posterior roots were cut from the caudal extremity to the superior part of the lumbar region. After the operation, it was found that the sharpest irritation of the posterior roots, or of the posterior columns of the cord from the extremity to about the middle of the lumbar region, did not appear to cause any pain; but, starting from the latter point, sensibility began to be manifested, and continued to increase in degree in proportion as the irritation approached the part of the cord which retained its nerves entire.

EXPERIMENT IV.—In a new-born puppy, after the carotids were tied, the spinal cord was divided across close to the medulla oblongata. The cord was then laid bare through nearly the whole extent of the dorsal, and through the upper part of the lumbar, region. After having satisfied himself that the reflex function was still active, and that pinching the skin of one of the limbs was followed by movements in all four, Brown-Séquard cut the posterior roots of the eight last dorsal and two first lumbar nerves. After the operation, he found that pinching the skin of the anterior limbs excited reflex movements in those limbs only, and that irritation of the hind limbs was followed by movements in those limbs alone. Artificial respiration was then practised. The reflex function was exalted; but the movements were still limited to either the anterior or the posterior limbs, according as the one or the other was excited. There was, therefore, no longer any transmission of nerve-force from the posterior to the anterior extremities, or vice versa; although the spinal cord itself between the lumbar and cervical enlargements remained uninjured.

Of the remarkable phenomena resulting from these experiments, Brown-Séquard could then see only one explanation, which I shall not lose time in stating, since he has, I think, most properly abandoned it for another, which rests upon a better foundation. This latter explanation is given in an interesting paper in the *Journal de Physiologie* for 1861, in which the most probable cause of the phenomena resulting from the experiments is stated to be an alteration in the nutrition of the spinal cord, in consequence of irritation of the sensory or excitomotor nerves. (*"Remarques sur quelques Points de la Physiologie de la Moelle Epinière et du Cerveau," Journal de Physiologie*, tom. iv, 1861, p. 584 et seq.) That irritation of the posterior roots does really produce this effect, appears to be satisfactorily proved. For instance, crushing with forceps those of a single pair on one side is followed by the same effects as division of the corresponding lateral half of the spinal cord; so that irritation of the centripetal nerve-fibres weakens the action of the corresponding lateral half of the cord at the place of insertion of roots that are irritated.

If the posterior roots of a pair of dorsal nerves be irritated on each side, anæsthesia and paralysis of the posterior extremities immediately follow; and, if several pairs of nerves be excited, the same kind of effects increase in proportion to the number of nerves and the degree of irritation. Moreover, it was found that the functions of the cord, as a centre of reflex action, and as a conductor of voluntary movements and

* Continued from page 345 of number for September 25th.

sensitive impressions, diminish in proportion to the degree of irritation of the sensory roots.*

We have already seen that section of the posterior roots abolishes the tone or tension of the muscles to which they distribute fibres. (M. T. G. Brondgeest, *Archiv für die Holländischen Beiträge zur Natur und Heilkunde*, vol. ii, 1859 and 1860.) Whether this loss of muscular tone be due to the curious changes which, according to Brown-Séquard, take place in the condition of the spinal cord on section of the posterior roots, is not determined; but it certainly must add to the paralyzing effects of those changes. In the experiments above described, *all* the fibres of the posterior roots were divided, and therefore *all* the muscles to which they were distributed were paralysed or relaxed; but in locomotor ataxy, except in the very advanced stage, only *some* of the fibres of those roots are injured, and in a very irregular manner; so that some of the muscles would appear to escape, while others are paralysed or relaxed. The way in which this unequal tension of the different muscles produces incoordination of voluntary movements, I have already pointed out.†

DEATH FROM ACCUMULATION OF HAIR IN THE STOMACH OF A WOMAN.

By PALEMON BEST, M.B.Lond., Louth.

THE patient was C. C., a labourer's daughter, aged 30 years. On my first visit to her, in April 1869, she complained of severe pain in the stomach, vomiting and diarrhoea, great prostration, and suppression of catamenia. She said that these symptoms had lasted for several years, and were due to a tumour in her stomach. The patient was tall, thin, with an anxious expression of countenance, and marked pallor of the skin and mucous membranes. The tongue was small, red, with white fur at the back and middle; pulse 80, weak, regular; respirations 18, regular. There was no oedema of the feet. The urine was copious, with no albumen.

When she was undressed and laid on her back, with her knees raised, a tumour was readily felt, mainly occupying the epigastric region. The tumour seemed rather globular in shape, was smooth and hard, but not painful to touch; it was very moveable, more so on the left than on the right side, and could be pushed upwards and to the left for a considerable distance under the rib-cartilages of the left hypochondrium. It did not accompany the movements of the diaphragm in respiration. The lower margin of the tumour extended obliquely from the right hypochondrium downwards across the epigastric and umbilical to the upper portion of the left lumbar region. This margin, although rounded, was very clearly defined, and sometimes seemed to have a notch about its middle; at other times, this notch could not be felt. The upper margin of the tumour was not perceptible on palpation; and percussion did not separate the dulness it occasioned from that caused by the surrounding organs—the liver, heart, and spleen. Although the tumour appeared close to the anterior surface of the abdominal cavity, the passage of flatus along the stomach between it and the tumour could be seen and felt, as in the descending colon in cases of enlargement of the left kidney. Long continued pressure and movement of the tumour induced pain and vomiting.

The vomited matter consisted of solid food and dark coloured sour fluid; it only very rarely contained streaks of blood after much straining. The pain in the stomach, which passed to the back between the shoulders, was much increased after taking food; and this act had nearly always induced both pain and vomiting during the last six or seven years. The vomited food and liquid never fermented. The number of stools in twenty-four hours varied from two or three to seven or eight; they were dark coloured, watery, not very offensive, and about equal in quantity. She had never suffered from constipation lasting several days and followed by profuse diarrhoea. The quality of food had no influence on the vomiting or purging.

She had never had jaundice. She had enjoyed good health until twelve years of age, when she suffered from a severe attack of ague. The catamenia appeared at the age of 15. About this time (1854), she began to suffer from pain in the stomach and vomiting and diarrhoea, but was able to keep her situation as servant, or attend to domestic duties at home, until about six years ago. Since this time, she had been constantly under medical treatment, and unable to undertake any employment. Mr. W. C. Calthrop, who had attended her at various times since 1856, at this time (January 1863) detected a hard nodular tumour at the epigastrium. She stated that the tumour was then very

much smaller than now (1869), and could be pushed entirely out of reach beneath the rib-cartilages of the left hypochondrium. Mr. Calthrop believed that she had "scirrhus" of the stomach. She continued under his care, the tumour increasing in size, and other symptoms not materially changed, until 1865, when she sought parochial relief, and became a patient of the district medical officer whom I succeeded in April last. She remained under this gentleman's care from 1865 to April 1869, except during a short interval which she spent in the Hospital for Diseases of Women or the Surgical Home in London, her symptoms undergoing no other change than increase in her debility and in the size of the tumour.

Such were the leading features of the case on my taking it; and I was fairly puzzled by them—at one time thinking that there was an enlargement of the left lobe of the liver; at another time believing that it might be some unusual form of splenic enlargement, which, by pressure on, or adhesion and perhaps projection into, the stomach, caused the distressing vomiting and pain. The hardness of the tumour was probably too great for it to be colloid cancer of the stomach; and the age of the patient, the moveability of the tumour, and its long duration, were other objections to this view, and, *à fortiori*, to its being either scirrhus or encephaloid cancer.

The treatment at first pursued was that suggested by my partner, Mr. Bogg, who visited the case for me before I had seen it, and considered that the tumour was due to an accumulation of feces in the transverse colon. Although I differed from this opinion, free purgation by mouth and by enemata was employed at intervals during three weeks, but without effecting any improvement. Bismuth and soda and hydrocyanic acid also failed in improving the patient's condition. Citrate of iron was then tried, and a decided improvement followed its use; the patient felt much stronger; her spirits improved; and in June the catamenia, which had been absent for four years and a half, reappeared, and were again seen at the proper time in July and August. The vomiting continued about the same; the diarrhoea was much less troublesome; but the size of the tumour remained the same. In the autumn, she frequently walked distances of about half a mile and back. On September 7th, when stooping to glean in a harvest-field, she was seized with violent pain in the abdomen, and became so faint that she was obliged to be carried home. Peritonitis immediately set in. Opium, hot fomentations, and turpentine stupes, were used; and she took brandy, milk, eggs, beef-tea, etc. Temporary improvement occurred, but was followed by renewed attacks; and, after much suffering, she died on October 25th, 1869.

Permission to make a limited examination of the body having been obtained, this was done on October 27th by Mr. Calthrop and myself. The peritoneal coverings of the stomach and intestines were glued together by recent lymph. The peritoneum contained about ten ounces of dark coloured fluid, composed probably in part of brandy. On lifting the stomach to discover the tumour, it was found that this lay within the stomach, which I opened by a long horizontal incision along its anterior wall, from the cardiac nearly to the pyloric extremity. A black tumour covered with hair was removed from the stomach and oesophagus.

It formed an accurate mould of these organs, and somewhat resembled a very small black swan. (It has been photographed.) The exact weight of the tumour was thirty ounces (avoirdupois). On section, it

* For further information, see the paper from which these statements are taken.

† It would be an interesting experiment to divide in an irregular manner *only some* of the fibres composing each posterior root of the nerves.