against disease as compared with another. Undoubtedly, the statistics of a single town are instructive; far more instruction, however, will be gained from the compared statistics of various and many towns. In addition, also, it would tend to throw light upon the causes of diseases, on the means of their prevention, and the right understanding of their etiology, the greatest reward of the worker in the wide expanse of "the science of medicine."

ON BICHLORIDE OF METHYLENE.

By BENJAMIN W. RICHARDSON, M.D., F.R.S.

THE recent notes in the JOURNAL on bichloride of methylene call for one or two explanatory sentences bearing on the action of that substance. I shall say what has to be said best by stating briefly the physical character of the bichloride, and the relation it bears to chloroform and other allied anæsthetics.

Chloroform and bichloride of methylene belong to one family of chemical organic bodies, in which family, carbon, as the base, combines with hydrogen and chlorine in different proportions. If one opens the hand, extends the thumb, and lets the extended thumb represent the carbon, all the other combinations of the group can be shown with the four fingers. Let the thumb represent carbon; extend the first finger, and let it represent hydrogen; keep the second, third, and fourth fingers together, and let them represent chlorine: the hand describes the composition of chloroform, which is composed of C. H. Cl Cl Cl. Bring up the second finger to the first, and let it now represent hydrogen, and the hand describes the composition of bichloride of methylene, which is composed of C. H. H. Cl Cl. Bring up the third finger to the second and first, and let it also represent hydrogen, and the hand describes chloride of methyl, which is composed of C. H. H. H. Cl. Bring the fourth finger to the other fingers, and let it, with the others, represent hydrogen, and the hand describes marsh gas, C. H. H. H. H, or, as it is called chemically, hydride of methyl. Or, let the fingers altogether represent chlorine, and the hand describes the tetrachloride of carbon, which is composed of C. Cl Cl Cl Cl, the hydrogen of marsh-gas being entirely substituted by chlorine. In these compounds, we have a complete chemical anæsthetic family group. Every member of the group produces sleep, and insensibility to pain, when inhaled by the living animal. In this respect, all the members of the family resemble each other in one leading characteristic, but they differ in detail; the difference turning on difference of physical property, and this, again, turning on the proportion of chlorine.

Let us compare, from this point of view, chloroform with bichloride of methylene. Chloroform, we have seen, is composed of C. H. Cl Cl Cl, and bichloride of methylene is composed of C. H. H. Cl Cl. Thus, chloroform has one more equivalent of chlorine than bichloride of methylene, and bichloride of methylene has one more equivalent of hydrogen than chloroform—in fact, in it, an equivalent of hydrogen is substituted for an equivalent of chlorine in the chloroform.

This difference causes essential alterations in the physical property. The combining weight of chlorine is as thirty-five to one of hydrogen; the difference causes an alteration in the specific gravity, the vapour density, and the boiling point of the two fluids. For example, the vapour density of chloroform, taking hydrogen as unity, is 59.75; of the bichloride of methylene, 42.5. The boiling point of chloroform is 142 deg. Fahr., and that of bichloride of methylene (according to the researches of Mr. Perkins, who has corrected the earlier observers), 104 deg. Fahr.

In turn, the difference of physical property modifies action. The lower boiling point of the bichloride causes it, when exposed to the air, to diffuse with greater rapidity than chloroform. I have seen it diffuse so rapidly that the condensed water from the breath of the patient became frozen in the inhaler. For this same reason, there is also a quicker diffusion of the vapour in the lung, and a quicker action on the blood. Again, for the same reason, there is quicker elimination of the substance from the body, with a less expenditure of the animal force, than occurs with the heavier chloroform.

In other respects, bichloride of methylene resembles in action the remaining chlorides; *i.e.*, it causes, sometimes, muscular excitement, and, occasionally, vomiting. These symptoms are more or less severely produced by all the representatives of the series; the effects varying with the amount of the chlorine. The chloride of methyl produces very little excitement, and no vomiting. Chloroform produces great excitement, and vomiting frequently; the latter symptom being often active and prolonged. Bichloride of methylene holds a place, in these respects, between the two.

When we correctly understand these points of difference between bichloride of methylene and chloroform, we see, with a little more reflection, what difference of symptoms will result from their inhalation, and what differences of method are required in their administration. If we diffuse equal portions of chloroform and of bichloride in an equal space of air, at the same temperature, each in a closed chamber, and place two animals of the same kind, weight, and condition, one in each chamber, the animal inhaling the vapour of the bichloride is narcotised more quickly than the animal in chloroform, and with less excitement; being narcotised, it will remain, while in the same atmosphere, as deeply narcotised as the animal in the chloroform, but it will die later; further, if it be taken out when it has ceased to breathe, it will more readily recover under artificial respiration than the animal which has ceased to breathe in the chloroform. If, however, two animals be exposed to the same quantity of the two narcotising agents, at the same temperature, not in a confined chamber, but in an open space, there will be difference of action. The animal breathing the bichloride of methylene, unless the dose be very large, will be excited, and will possibly not pass into the third degree of narcotism at all; while the animal breathing the chloroform will be narcotised to the third degree. The reason of this difference is obvious enough, and is, that the lighter bichloride has been lost by diffusion in the air, more rapidly than the heavier chloroform. Again, if the two animals be narcotised to the same degree, one with the bichloride, the other with chloroform, and the narcotics be withdrawn at the same moment of time, the animal narcotised with the bichloride will recover more quickly than the animal narcotised with chloroform, and the return to the natural temperature, from the reduced temperature to which both will have been brought, will also be more quickly effected.

These facts explain all the anomalies which those who exhibit bichloride of methylene to the human subject have so often observed; they explain, moreover, that the anomalies depend, not on the substance, but on the mode in which the substance is administered. Bichloride of methylene, owing to its rapid diffusion, can be made to produce good anæsthesia for short operations within the minute; an operation lasting two minutes may be performed, and, the vapour being withdrawn, recovery may be perfect in five minutes. We can scarcely want anything more in the direction of rapidity of action. Bichloride of methylene can be made to produce deep anæsthesia in four minutes; and, when a long operation has to be performed, the sleep can be sustained, without a struggle, for an hour (I am speaking from experience); recovery, when the vapour is withdrawn, being complete in from ten to twenty minutes. We can scarcely want anything more in the direction of prolonged action.

To obtain these two different results, there must be a principle in the method of administration of the bichloride. To produce the rapid effect, the agent must be given as Mr. Rendle gives it; *i.e.*, its vapour must be confined in a small, simple, hollow cylinder, which adapts itself to the mouth and nostrils at one end, and is perforated with holes, for admission of air, at the other; the cylinder being lined within with loose demette. The diffusion of a drachm of the fluid is sufficient for very short operations when applied in this way. To produce the prolonged effect, the same cylinder may be used, but air must be more freely admitted by holding the inhaler more loosely over the mouth and nostrils; and, when the insensibility is complete, the administration must be continued in full, for a minute or two, unless some urgent symptom should interfere, until the organism is well charged with the narcotic. A sufficient charge once introduced, the sleep may be sustained by the simple process of readministration, carrying the effect each time a little longer than the period when the anæsthesia is perfect and the voluntary muscles are quiet.

The question of the age of the patient has been adduced as having a bearing upon the action of bichloride of methylene and of chloroform, the idea being that chloroform is best suited for the young. There is really no ground whatever for the fancy, except this: that, as the respiration of children is quicker than that of adults, they come more quickly under the influence of the vapour of the bichloride, because they inspire more of it. Hence, to babes and young children, the vapour, even for short operations, should be administered more diluted with air than to adults. Happily, if in young children or other young animals quickly narcotised with bichloride of methylene, the respiration be for a moment checked, prompt artificial respiration, gentle but prompt, suffices to restore animation with a certainty that is remarkable. I have restored suspended respiration in a young animal under bichloride of methylene, two and even three times after respiration had ceased.

A practice has been suggested of first putting the patient to sleep with the light bichloride, and of keeping up any required prolonged effect with chloroform. I have already said enough to shew that this practice is unnecessary. I would add, now, that it is excessively dan-

gerous, and I predict, without any hesitation, that fatal accidents must inevitably follow upon it. It is far better to use chloroform from the beginning of an operation to the end of it, than to begin with a lighter anæsthetic, push that to the degree of annihilating sensibility, and then to cap the proceeding by adding a dose of the heavier agent. A horse at its fullest speed in a race might possibly reach the goal successfully if, by some sleight of hand, the jockey on its back could be suddenly weighted; but, as a rule, the horse, already well-nigh exhausted, would begin to flag and would go down under the additional burthen.

I have written these lines about the bichloride without any desire to praise it because it is an introduction of my own. It is naturally gratifying to observe that a theory of action, based on knowledge of physical laws, proves correct in practice, as the theory relative to the action of bichloride of methylene has proved; but men who are absorbed in the study of natural phenomena, as I am, soon die daily to their work, and when once they have placed a thought or practice before the world, treat it as belonging entirely to the dead. To me, bichloride of methylene is the same as to others; and, although I know it has many advantages, I am so far open to its faults, that if, to-morrow, I could find an equally manageable and effective volatile anæsthetic, out of the chlorine family altogether, I should throw over the bichloride without a moment's hesitation, feeling the act a duty. At the same time, while it retains its place in practice, it is just to remove any prejudices against the bichloride of methylene, resting on imperfect perception of its properties and action.

POSTSCRIPT.—The above paper was sent to the JOURNAL in September last; and since that time a case of death has occurred at the Charing Cross Hospital during the administration of the bichloride of methylene. The facts of that case, ably and faithfully reported by Mr. Peter Marshall, suggest to my mind no modification of what I have written, since the administration of the narcotic and the proposed operation in the case referred to were planned almost against hope, to save a man who was sinking from exhaustion, and whose breathing was mechanically obstructed by cancerous growth. I do not see that any general anæsthetic can ever be discovered to meet successfully a case of this extreme character. It is a case in treating which science can only be relieved from the chance of discredit by the firm refusal of her ministers to accept in her name, under the blessed, even though unreasonable, impulse of human kindness, an useless responsibility. In this position, we can put ourselves only in the place of the sufferer, greeting the impulse as the noblest gift that we could ask for or receive, and accepting ourselves (as the patient in the case of which we are speaking did) the responsibility of the attempt made to save us from a death inevitable, imminent, and, in its threatened natural aspect, far more cruel and terrible than if it came from the effort to avert it.

CASES OF HERPES CONTAGIOSUS (?).

By J. BRENDON CURGENVEN, M.R.C.S., etc.

MATTHEW PAINE, aged 22, a coachman, married, of sober habits, inhabiting a newly built stable in a healthy situation, was in good health until October 13th, when he felt his throat sore; he had shivering, and severe vomiting of water and mucus, which was described by him to be "as green as grass".

Oct. 14th. I first saw him. He complained of great muscular soreness over the trunk and limbs. His skin was hot; face flushed; pulse over 90; tongue red and clean. There was a white patch of herpes on each tonsil; redness and swelling of the fauces; and the submaxillary glands near were swollen and tender. Deglutition was difficult and painful.

Oct. 16th (fourth day). The tonsils and the mucous membrane of the pharynx were more swollen, and there was greater difficulty in deglutition. The pimples on the tonsils had coalesced into uniform white patches, yet they were not more extended in circumference. Other patches had appeared on the uvula and the upper and under surfaces of the tongue, which organ continued red.

Oct. 17th (fifth day). He slept very little last night, being constantly disturbed by difficulty of breathing and accumulation of large quantities of thick tenacious mucus in the throat; and towards morning he became delirious. He discharged, by retching, about six ounces of thick jelly-like mucus, that adhered firmly to the bottom of the basin. The inflammation—possibly the herpetic eruption—had extended down the œsophagus and into the larynx; for he was now quite hoarse, his voice being not above a whisper. There had appeared, since the previous day, clusters of herpetic vesicles at the corners of the mouth, at the orifices of the nostrils, on the eyelids near the inner canthi, on the left

side of the scrotum and prepuce, with single vesicles scattered over the thighs, forehead, and backs of the hands. He had partial loss of power in his hands and wrists; he could not close his hands, nor straighten his wrists. The temperature was high; pulse 98; bowels regular.

Oct. 19th. He had a better night, but continued to bring up a quantity of thick mucus after each dose of the medicine (chlorate of potash and hydrochloric acid). There was a patch of vesicles on each side of the neck, immediately over the tonsils: these, about a dozen in each patch, had not coalesced, but the vesicles of all the other patches had, forming one scab for each group; that on the scrotum rubbed off, leaving a raw surface. His hands and wrists had recovered their power, which might or might not be due to a blister applied between his shoulders on the previous night.—His wife complained of lassitude, slight shivering, and sore throat. She had a patch of herpes on the left tonsil; the tongue was slightly coated; pulse 84; skin hot.

Oct. 20th. The husband's symptoms were improving; he could swallow better, and took a little bread in his beef-tea. The discharge of mucus was less.—The wife had a patch coming on the right tonsil.—His mother, who had sat up two nights with him, and had been with him more or less from the commencement of his illness, had a patch on her right tonsil.

Oct. 21st. The husband was better; temperature by thermometer, 100.6.—His wife vomited, yesterday and to-day, a watery fluid. She had muscular soreness, heat of skin, pulse 92, and aching of the hands.—His mother had muscular soreness and aching pains in the hands. A patch had appeared on the left tonsil. Tongue slightly coated; pulse 90.

Oct. 22nd. The wife had several red spots on her hands last night ("herpes sine vesiculis," Wilson), only two or three of which on the left hand gave rise to vesicles.—The disintegrated mucous membrane over the spots of herpes on the husband's tonsils and tongue had been cast off, leaving ulcers with well defined and healing margins.

Oct. 26th (fourteenth day). The husband was still hoarse. The wife and mother were almost well, their throats being only slightly sore; the herpes was gone. The husband's voice gradually returned during the next few days. The cuticle is peeling from the hands and feet.

Oct. 27th. The husband's brother was attacked with shivering, sore throat, and fever. He had a group of white pimples on each tonsil. Tongue coated; pulse 102; skin hot and perspiring.

Oct. 28th. The pimples were less perceptible; pulse 120. He had cramp in his feet occasionally.

Oct. 29th. The eruption on the tonsils was gone; the tongue was coated and swollen. He had a troublesome cough last night; no chest-symptoms. He had an inflamed gland in the right axilla.

Nov. 1st. The symptoms were, copious perspiration, occasional cough, and cramp in the feet. Pulse 84.

Second Group of Cases.—Oct. 22nd. On visiting a family that I was attending, I found the housemaid with a sore throat, which, on examination, presented a patch of herpes on the left tonsil, and two or three pimples on the right. She had had shivering and muscular soreness, but no sickness. The cook had just recovered from a sore throat that commenced seven days previously. The two servants slept in the same bed.

Oct. 23rd. Last night, the mistress was taken with shivering and sore throat. The tonsils and fauces appeared red and swollen, but there were no pimples.

Oct. 24th. She was sick twice last night. There were two pimples on the left tonsil.

Oct. 25th. She felt much better; the pimples were gone, and the redness was less.—The patches on the housemaid's tonsils had gone, and she felt better.

REMARKS.—It is evident that these cases are cases of herpes in which the whole system is involved. In this, they differ from those local and milder forms with which every one is well acquainted. The disease differs also, in that it is contagious, as far as these two groups of cases permit an opinion. But herpes may be said to be a local neurosis, and cannot appear as a general febrile neurosis. I would ask, Do we not meet with hemicrania and other kindred local neuroses? and can we say that these are in no way produced by a similar disturbance of nerve-force as those general febrile neuroses, ague and remittent fever?

Herpes has been divided by Willan, Hebra, and Wilson, into herpes labialis, nasalis, and præputialis; herpes zoster, iris, and circinatus. I can find no mention by these authors of any form of the disease like the one I have described, in which all the species of the first class were combined in one case. Hebra remarks, that "an herpetic affection of the skin of the face, the red parts of the lips, and the mucous membrane of the mouth, sometimes occurs in perfectly healthy subjects, being then the only morbid condition which is to be detected; but, in