had taught me, and then I began to see that in most instances that displacement is merely a symptom. I continued the friend of Velpeau to the day of his death, but he never got much further in uterine pathology; and, when I broached the subject, used to shake his head, and say in his amiable, jocose way, "Don't let us talk about it; you are a deserter from my flag; you have gone over to the enemy, and deserve punishment". Velpeau's influence, however, in this question, has been great and pernicious on the Continent.

At home, in 1843 I believe, our lamented colleague and friend, Sir James Simpson, published in the *Dublin Medical Journal* a paper on uterine displacements, in which he adopted pretty nearly the same views as Velpeau—those of what I call "the mechanical school". A year or two later, in 1845, he called on me in London, and said that he had procured my recently published work on uterine inflammation, had read it twice through carefully, on his way to town, and, on his return home, would examine into all the questions I raised. I subsequently heard from him that he had found my descriptions of disease clinically correct. The influence, however, of former views remained powerful in his mind throughout his life; and I found his practice uncertain in this respect. Sometimes he had cleared up those of his cases which I subsequently saw, as regards inflammatory conditions, and sometimes he had not, seeing nothing but the displacement, even when decided lesions existed. I have never before alluded to the opinions of my departed friends: I now do so with all esteem and affection; but they are gone from us; the doctrines which they held are become a part of the history of the profession, and as such they may be alluded to and criticised without disrespect to their memory.

In conclusion, I would recapitulate the views I have endeavoured to express in the following terms.

1. I consider that, under the influence of mechanical doctrines pushed to an extreme, uterine displacements are by many too much studied *per se*, independently of the inflammatory and other lesions that complicate and often occasion them.

2. That the examinations made to ascertain the existence of inflammatory complications are often not made with sufficient care and minuteness, as evidenced by the fact that I constantly see cases in practice in which inflammatory lesions have been neglected entirely, and in which the secondary displacements have been alone studied and treated.

3. That inflammatory lesions are often the principal cause of uterine displacement through the enlargement and increased weight of the uterus or of a portion of its tissues which it occasions.

4. That when such inflammatory conditions do exist, as a rule they should be treated and cured, and then time should be given to Nature to absorb and reduce hypertrophied and engorged tissues before mechanical means of treatment are resorted to.

5. That the relief from the sensation of bearing down which pessaries and bandages give is no real criterion of their being the proper means to use, such relief being often felt when there are inflammatory lesions present, which their presence aggravates.

6. The above statements must not be considered in any way to imply that I do not recognise other causes of displacement of a non-inflammatory nature, such as laxity of ligaments and soft parts, wide pelvis, laceration of perinæum, severe shocks, etc.

ON INTERMITTENT TETANY.

By C. CURRIE RITCHIE, M.D.,
Physician to the Hulme Dispensary, Manchester.

It is remarkable that this affection should have so long escaped recognition by the English profession, especially as it must be by no means uncommon. No mention is made of it in any English treatise on systematic medicine that I have seen, nor does it appear in the *Nomenclature of Diseases* issued by the Royal College of Physicians.

It is probable that spasmodic contractions of muscles have long been observed and classed with other affections of a convulsive character, but it was not till Dance published his "Observations on a form of Intermittent Tetanus", in 1831, that special attention was directed to the well-marked group of phenomena constituting the disease now known as intermittent tetany. Trousseau, in his *Lectures on Clinical Medicine* (New Sydenham Society's translation, Lecture xiii), gives a most admirable account, historical and clinical, of this affection, which first received the name "tetany" from Dr. Lucien Corvisart, in 1852.

As professional interest in the subject has been revived by the publication of a case with clinical remarks by Dr. Wilks (*BRITISH MEDICAL JOURNAL*, 11th June, 1870), I have ventured to record the following cases which have come under my observation within the last two years.

CASE I.—Mrs. M., aged 28, sent for me on 28th August, 1868. She had three children, and was about the seventh month of her fourth pregnancy. She had suckled her youngest child for sixteen months, during the latter five of which she had suffered from profuse menorrhagia. Four months ago, she began to have severe headaches and pain darting down the back, accompanied by painful contractions of the hands and feet, which prevented her from attending to her household work. These attacks came on at first every three or four days, each lasting about two hours; latterly, the attacks had become more frequent. Ten days ago, the weather being very cold and wet, diarrhoea came on, which had weakened her very much. On receiving some unpleasant news, the morning I saw her (28th August), she had "fainted"; immediately afterwards, however, on regaining consciousness, she found her "whole body twitching" so that she was unable to sit quietly, or to use her limbs. She had to be undressed and put to bed. When I saw her about 11 A.M., she was sitting propped up in bed, her expression evincing great anxiety. There was twitching of the muscles of the face, and drawing together of the lips, interfering with articulation; the fingers were rigidly fixed in a semiflexed position, with the thumb forcibly drawn across the palm; the wrists were semiflexed. The toes were also incurved, the dorsum of the foot being highly arched, and the muscles of the calf hard and tense. The fingers and toes could not be straightened; she had a feeling of numbness in them. Her mental powers were unimpaired; pulse 108, temperature 99.2° Fahr. I put her under the influence of chloroform, and was then able to straighten the fingers and toes; they again became rigidly contracted when the anæsthetic was withdrawn. She was ordered ten grains of bromide of potassium, to be taken every half hour; and at my evening visit (7 P.M.), I found the contractions less marked, and the pain diminished. I ordered the bromide to be taken three times a day. The contractions gradually ceased; the patient remained free from them after the evening of the 31st August. She was confined on November 2nd. As she felt weak, I recommended her not to nurse her child, and ordered her citrate of quinine and iron, with cod-liver oil. When I last saw her (August 1869), she told me that she had had no return of the contractions, and now enjoyed perfect health.

CASE II.—Mrs. A., aged 34, was delivered of her sixth child on March 27th, 1870; the labour was natural, with the exception of slight atony of the uterus. She made a good recovery, and remained well till the end of April, when she began to suffer from severe headache after each act of suckling her child. A few weeks afterwards, she began to have "flickering" feelings all over her body, which amounted to a kind of numbness in her hands. Her sense of touch became impaired, so that she felt as if she constantly wore a pair of thick woollen gloves. She stated that "she had let several articles of crockery slip out of her fingers" from this cause. I saw her on June 3rd. For several days she had been suffering from painful contractions of the hands and feet immediately on getting out of bed every morning. At 4 A.M. on June 3rd, she was seized with severe pain in the back, and cramps of the limbs; the feet were arched. I saw her at 10 A.M.; her condition was as follows. She had a sallow, cachectic appearance; there was slight muscular twitching of the face and eyelids; speech not affected; mind clear. The fingers were semiflexed and perfectly rigid; the thumbs drawn across the palms, abduction and extension causing great pain; muscles of the forearm hard and tense; upper arm natural, with the exception of slight contraction of the biceps. The muscles of the calf were so tense that they felt to the touch like pieces of wood. The heel was drawn up and the toes incurved. The pain in the legs and feet was very great, and was accompanied by a tingling sensation. Previously to my seeing the patient, continuous friction of the legs and feet had been kept up for about three-quarters of an hour, without any intermission of the contractions. Immersion of the limbs in cold water had no effect; cold water bandages were then applied, which somewhat relaxed the spasms, and afforded relief to the patient. She sent for me again, however, about 2 P.M., as the contractions were as bad as ever. Ether spray was then applied to the feet, which almost immediately stopped the contractions; they were readily reinduced on compressing the limb below the knee with both hands, but ceased on reapplying the spray. At 8 P.M., there had been no return of the cramps; half a drachm of chloral hydrate was then administered. She had a good night's rest, but had two slight attacks next morning, June 4th; to continue the chloral three times a day. On June 8th, she was quite free from the contractions and pain; a mixture containing the ammonio-citrate of iron with infusion of calumba was substituted for the chloral. I saw her on July 15th, previously to her leaving Manchester; she was much improved in appearance, and felt perfectly well.

CASE III.—M. C., a strumous-looking boy of 13, used to have "working fits", according to his mother's statement, when he was about three or four years old; his face and mouth were drawn to one side;

his hands twitched and were sometimes so firmly set that it was very difficult to separate the fingers. His mother does not remember whether his feet twitched, but on several occasions he asked her to take off his boots and rub his feet. These attacks lasted over three or four months, each attack lasting from a few minutes to two or three hours. Some months after this, he had scarlatina, followed by general dropsy, from which he recovered very slowly. His mother only remembers his having had one recurrence of the "cramps" shortly after his recovery from scarlatina. After this, he remained perfectly free from them, till a year ago, when he fainted on seeing his father evacuate the fluid from a blister on his foot with a darning needle; on regaining consciousness, he had one of his attacks of twitching. He came under my care as an out-patient at the Hulme Dispensary, on 21st June, 1870, suffering from severe coronal headache, worst in the morning. His appetite was good; his bowels had acted regularly twice a day for a month. He was ordered quinine, and was improving slowly under it, till the morning of the 19th July, when he came to see me at the Dispensary. His headache, he said, was worse than ever, and he complained of severe pain at the epigastric region. He seemed to be suffering so much, that I was proposing to his mother to give him a subcutaneous injection of morphia, when he fainted. He remained unconscious for about a minute; he then had painful contractions of the fingers which he said "felt numb and dead". The thumbs were firmly drawn across the palms, and the fingers were rigid and semiflexed. There was slight twitching of the muscles of the face. Speech was thicker than natural, but there was no great compression of the lips. He had feelings of cramp in the legs, and there was twitching of the feet. The twitchings were relieved by immersion in cold water. I put him on twenty-grain doses of bromide of potassium three times a day. He has had no return of these attacks up till to-day (August 1st), and his headache is much better than it has been for three months. He is still under treatment.

I have met with other two cases of this affection, but unfortunately have no notes of them taken at the time. One was the case of a boy four years of age, who came under my care as a dispensary patient. His symptoms corresponded closely to the history of the attacks in M. C. (Case III), about the same age, and yielded readily to small frequently repeated doses of bromide of potassium. The other case was that of a young woman of nineteen, who had suffered from menorrhagia for several months previous to her being attacked with painful contractions of the fingers and toes. In this case, the symptoms though distinct, were not severe; the treatment was, therefore, mainly directed to the regulation of the menstrual flow, and the spasms gradually passed off.

This affection is so rarely fatal, that opportunities of inquiring into its pathology and morbid anatomy are extremely few; indeed, in all Trousseau's experience, he met with only one fatal case. There can be little doubt from the nature of the disease, however, that it is essentially a disorder of function. Trousseau states that the alleged predisposing causes of this curious affection are nursing, menstruation, pregnancy, and the puerperal state—with perhaps the pathological conditions complicating dentition. Cold he considered not only an "exciting cause, but sufficient by itself to bring on" this disease; and so frequently did it occur, under the observation of Trousseau, in nursing women, and as apparently the immediate result of cold, that he at one time called it "rheumatic contraction of nurses." He also states that chronic diarrhoea is the most active exciting cause, and that after the first attack of these contractions, they may be reinduced by emotion, or by "compressing the affected parts, either in the direction of their principal nerve-trunks, or over their blood-vessels, so as to impede the venous or arterial circulation." This last observation I was able to verify in Case II; but in most of the English cases, not only has there been no diarrhoea, but in several cases the bowels have been confined for some time before the attacks. The diagnosis depends chiefly on the intermittent character of the tonic spasms, with the peculiar contraction of the hand and foot, and incurvation of the thumbs and toes.

I was much interested in the parallel drawn by Dr. Moxon in his valuable paper on this subject (*Guy's Hospital Reports*, Series III, Vol. xv), between the symptoms of this affection, and those produced by the use of ergoted rye; and in the two cases I met with after reading his observations, I directed inquiries to this point, but could not elicit that rye-bread had been at any time eaten by either patient.

With regard to treatment, Trousseau recommends blood-letting, and afterwards quinine, with small doses of opium and belladonna as adjuncts. Dr. Broadbent adopted the belladonna treatment in two cases (*BRITISH MEDICAL JOURNAL*, August 6th) in February of this year, with the result of the immediate cessation of the attack, which returned several times on the drug being withheld. In another of his

cases, the attacks ceased after the administration of the ammoniated tincture of valerian.

Dr. Moxon's case, and also three cases reported by Dr. Haddon (*Edinburgh Medical Journal*, August 1870), as well as three of my own cases, were treated with the bromide of potassium, and the contractions disappeared in all. But, as Dr. Wilks very justly remarks, the disease has a spontaneous tendency to subside; and in one of my cases this was amply verified. It is probable, therefore, that no specific effects can be attributed to any of these remedies; and that more is to be hoped for from the regulation of disorders of function, when such are present, and a generally tonic system of treatment.

During the paroxysms, chloroform inhalations are of great service, but, unfortunately, the contractions usually return when the inhalation is stopped. Trousseau pointed out the fact that cold sometimes stops the contractions, when applied to the affected parts, even although it has a powerful influence on the production of the affection. I tried this in the three cases which I have reported in full. The hands and feet were immersed in cold water without effect in two cases—in one instance, the application of cold water bandages relieved for a time. Having determined to try the effect of ether-spray in my next case, I did so in Case II, with the result of a complete cessation of the attacks. Whether the effect was physical or psychical, I cannot tell; I would venture to suggest, however, that it might be worth a trial in such cases in future.

CASE OF PARACENTESIS VESICÆ.

By W. DRAPER, M.R.C.S., York.

THE comparative infrequency with which we are called upon to tap the bladder, and the peculiar circumstances indicating the performance of the operation in the present case, may make the following notes of interest.

The subject of the operation was a married man, of rather robust constitution, aged about 65. Some years since he consulted my partner, Dr. Procter, in consequence of difficulty in passing his urine. On examination, it was found that a large-sized catheter could be freely passed into the urethra for about an inch and a half, but beyond that point not even could a No. 1 be introduced.

As the obstruction was not of a spasmodic character, and as there was no history of stricture from disease, a diagnosis of malformation of the urethra was made: this opinion was strengthened by the patient's statement, "that he had never been able to pass his water in a full stream". As the dysuria was only slight, no further notice was then taken of the case. Gradually, however, as the patient advanced in years, his urination became more difficult; and at the time to which the present report refers, another examination discovered a considerable enlargement of the prostate gland. Catheterism was again attempted, but with precisely the same results as formerly: the smallest instrument could not be introduced beyond the point before mentioned. The prostatic enlargement continued until micturition was entirely obstructed. With the view of overcoming the malformation, I passed a small director down the urethra as far as the obstruction, and attempted to dilate the parts by means of a long narrow knife, but the operation was ineffectual. The symptoms and consequences of retention rapidly became more urgent, and the most available means of relief seemed to be paracentesis of the bladder. The opinion of another experienced surgeon (Mr. Keyworth) was taken, which fully concurred with that already formed. I accordingly operated by making a small primary incision just above the symphysis pubis, and then introducing a long curved trocar and cannula into the distended bladder. A large quantity of highly ammoniacal urine was drawn off, to the intense relief of the patient, and an elastic catheter passed through the cannula and retained in the bladder, the cannula being withdrawn. From this time up to that of the patient's death, which occurred *purely from exhaustion* a week after the tapping, he continued in comparative comfort, not a sign of peritonitis or of any other evil consequence of the operation being evidenced. Thus the operation as a palliative measure—the only one for which it was performed—may be considered as eminently successful; and the results would quite justify its repetition in a similar case. The pubic region was selected as the *locus operandi* in preference to the rectum, in consequence of the highly infiltrated state of the tissues surrounding that part, and the difficulty there would have been in acquiring and maintaining the necessary position in a highly sensitive, ponderous, and helpless man. The operation of perineal section was inadmissible for the same reasons; and lastly, as the chances of cure were just as remote in one case as in another, the operation least irksome to the patient was adopted.