water cannot be added to the list) as the recognised causes of fever have been, and are, in operation here to an almost unequalled extent is not to be denied; and it would be difficult to decide which enjoys the pre-eminence as an epidemic producing agency. Separately, each would play an important part in the deterioration of our labouring population, but on their united action their baneful consequences are incalculable. One result is the production of a race impaired in mind, morals, and physical constitution, possessing organisations so enfeebled and susceptible, that a noxious germ deposited therein, finds a soil prepared for its reception admirably suited for its development and propagation.

That relapsing fever is *contagious* it is not necessary to adduce other proof than that already on record. Our experience as regards the house-surgeons, nurses, and patients, ninety-two per cent. of whom contracted the disease directly from others, confirms what has already been written on the subject. This, unfortunately, is lost sight of—is almost ignored—in the measures which are deemed prophylactic of epidemics.

While sincerely sympathising with every effort put forth to promote the wellbeing, and to improve the *physique*, of our lower orders, such as extended sanitary measures, the education and improvement of the masses, and the suppression of the vice of drunkenness, I have, for many years, maintained that, concurrently with these means, energetic measures should be resorted to for the prevention of epidemics by prompt action when the first signs of such are manifested; thus to stamp them out when in their inchoate state. That this desirable consummation is practicable, I firmly believe, if a general system of registration of disease were established, if the sick were promptly removed for treatment, and if suitable arrangements were made for the reception of the healthy inmates of an infected dwelling while undergoing disinfection and purification. When these measures are wisely carried out, we may hope that, though sporadic cases of infectious disease might occur from time to time, they would not assume an epidemic form, carrying in their wake death and misery.

OBSERVATIONS ON THE TRUE NATURE OF THE SO-CALLED "SKIN-GRAFTING".

By DAVID PAGE, M.B.,

President of the Royal Medical Society of Edinburgh.

ALTHOUGH the experiments of M. Reverdin on the transplantation of skin to ulcerated surfaces, so as to hasten the healing process, have been repeated in many instances with accredited satisfactory results, nothing beyond the mere clinical details of these experiments have been advanced in explanation of the physiological changes involved. From careful observations conducted throughout a case of successful transplantation, I have come to the conclusion that these changes and their results are altogether different from those which previous information on this subject would lead one to infer. When a breach of surface fails to heal by primary union, the efforts which are next made to accomplish healing by granulating action depend for success upon the combined facilities for contraction and cicatrization that are afforded by the surrounding healthy textures. The callous ulcer is an example on the point, for in it the removal, by appropriate means, of the solid oedema and hardened cuticle is quickly followed by a reappearance of vigorous granulating action upon a surface till then unhealthy and inactive.

So far, then, does healing by granulation depend upon the co-operation of neighbouring parts; and, when once it is fairly established, the formation of a cicatrix over the breach that contraction fails ultimately to enclose, becomes a matter of time. To shorten the period required, and, if possible, to insure an integumentary cicatrix—that is to say, one more or less identical with the true skin—are considerations of interest, for which the novel method "of skin-grafting" has been supposed to afford the conditions. To ascertain the exact nature of this method, an opportunity occurred in the case of a man, 38 years of age, suffering from an extensive callous ulcer of the leg, who was admitted to the wards of Dr. Gillespie, senior surgeon of the Royal Infirmary, in the month of July last. Dr. Gillespie has kindly allowed me, as his house-surgeon, to carry on in the present instance these observations. The ulcer exhibited at that time a greyish coloured and irregular surface, little more than moistened with a foetid transudation, and measured six inches in length; being only prevented from completely surrounding the leg by a narrow isthmus of sound skin, about two inches in breadth, behind. After the solid oedema of the leg and foot and the hard white edges of the ulcer had been removed by the usual means, the surface became covered with vigorous granulations; and, under simple dressings and rest, contraction and cicatrization proceeded actively, so that by the middle of October the ulcer was reduced to a narrow irregular patch,

lying in the midst of newly formed cicatrix. During the next three weeks, healing action seemed to be in abeyance, even under frequent changes in the mode of treatment; and it became evident that the opposition to further progress lay in the resistance of the surrounding cicatrix.

On the 4th November, I removed two portions of skin from the front of the upper arm, the patches—each a quarter of an inch in diameter—being simply raised up in dressing forceps and cut off with a pair of scissors. They were then placed upon the healthy granulating surface of the ulcer, and there retained by strips of isinglass-plaster. The dressings were not changed until the second day, and then the pieces of transplanted skin were found in position, seemingly adherent, and their margins exhibiting the transparent and purple tint of the delicate cicatrix at the edges of the ulcer. Two days later, on the 8th November, when the dressings were next removed, an increase of suppuration had occurred, and one of the pieces came away in the discharges; the other still adhered, but the cuticle had desquamated from its surface, and, on being removed, displayed a delicate and pink pellicle below. The edge of this pellicle was carefully raised at one part, and proved to be colourless and transparent, so that the vascular appearance was in reality due to the granulations upon which it lay. And then, upon examination, with the aid of the microscope, the film was seen to consist of young epithelial cells. The ulcer was examined again on the 10th November, by which time the remaining patch had increased to double the original size, while two days later still it had at one point reached the circumference, so that the ulcer now exhibited a cicatrix extending, by means of the transplanted portion, into the very centre of the surface.

The identity of this latter pellicle with the one spontaneously proceeding from the edges of the ulcer was established by the amount of resistance that both offered when detached, and by microscopic examination, showing them to consist of young epithelial cells. Further, when the healing of the ulcer was completed at the end of November, the cicatrix artificially induced, and that formed naturally elsewhere, were indistinguishable one from the other, and under the microscope proved to be of the same fibro-cellular structure.

In this way, then, the cicatrization of an obstinate but healthy ulcer was effected; but beyond this nothing more, for observation showed the identity of the central and peripheral cicatrices, which means also no nearer approach to the condition of true skin, but merely a more rapid growth of a tissue of very low vitality, devoid of the functions of the true skin, and easily destroyed by causes that would not affect the latter; for indeed, in the case under observation, the ulcer had existed for twelve years, during which time it had alternately healed and broken out from the weakness of the cicatrix; and, from what has now been accomplished, I anticipate no better security against a similar condition.

I am convinced that the so-called "skin-grafting" consists, in truth, not of a transplantation of true skin, but of epithelium, exactly analogous to what is occasionally observed in certain ulcers, especially in those associated with a syphilitic cachexia, where patches of delicate cuticle appear spontaneously on the surface and spread towards the circumference, but to be carefully discriminated from similar phenomena in ulcers resulting from extensive burns, for in these the apparent regeneration of skin is due to the fact that little insulated spots of true skin having escaped destruction, reappear after the first effects of the injury have been removed. These act as centres for healing action in a very different way. That the action of the transplanted portions is not merely that of affording a point of attachment (*point d'appui*) for spontaneous cicatrization, was satisfactorily proved by a series of experiments, in which pieces of sheep's-skin, caoutchouc, and other substances, were applied to the surfaces of healthy ulcers, and then retained in the same way as the portions of skin. But they only acted as foreign bodies, and on removal I found that a destruction of the granulations and depression of surface had resulted from their pressure.

Besides the cases in which skin-grafting has proved successful, have been those in which the granulating action was healthy; for if it were weak or absent, the patches refused to adhere. The temporary disappearance of some of the transplanted pieces, noticed by at least one surgeon, is, I think, due to the varying thickness of the young epithelial layer removed.

In conclusion, I am led to believe by these results of observation that, beyond a somewhat greater rapidity of cicatrization, especially where contraction of surrounding textures is resisted either by the situation or the extent of the breach of the surface, this method of skin-grafting is of limited application; and, as its results lead to no regeneration of the true skin, but merely the formation of a cicatrix in every respect identical with that formed spontaneously in the natural process of cure, it is an operation which cannot rank with plastic operations proper, and which is not likely to occupy a permanent position in minor surgery.