may exist long before the percussion of the chest reveals undue resonance, or auscultation shows any lengthening of the expiratory murmur. Might we not here do something by early notice of subjective symptoms?

In nervous diseases, the application of the remarks I have made are endless. The early symptoms of ataxy and of general softening of the spinal cord are often such as are subjective only; feelings of weight, or want of equilibrium, or of restlessness, which go for nothing when the patient is under the stimulus of the physician's presence, and can walk firmly, and stand on one leg with his eyes shut; but which, nevertheless, when neglected, creep on until a malady is obviously present which our science may arrest, but can rarely cure.

The early subjective symptoms of brain-disease are too numerous for detail. I will only allude to a few examples, which I am sure you can multiply indefinitely. The "general paralysis of the insane" often passes unnoticed until it is highly marked; and yet at an early period some apparently trifling change in the feelings of the patient, some exaggeration of self-importance, or slight egotism of manner, might have given the alarm, when found in conjunction with certain slight and occasional physical weaknesses which have to be looked for to be found.

The early symptoms of "softening of the brain" are often complained of and disregarded, or set down to dyspepsia or fatigue, simply because they are subjective. A feeling of bewilderment, or of want of mental grasp or grip; an inability to take in new notions; a listlessness or carelessness, with sometimes a sense of well-being, are among the most significant of the signs of cerebral decay. Might we not here often arrest and even postpone indefinitely the progress of disease?

In epilepsy, it often happens that long before the attacks have developed into even the minor seizures of *le petit mal*, the patient has been conscious of some strange and sudden mental condition which he cannot describe, but which is significant enough. Here, and in allied cases, could we not do much by an early recognition of the possible disease? In all these cases I think we could. A scrutinising attention to the minutest departure from health, in mind, or emotion, or sensation, would, I believe, help us greatly in the estimation and prevention of disease; and we must ever remember that in the practice of our noble profession we have to deal with man as a whole, to examine him and to treat him as such; that by neglecting any part of the information which he can give us with regard to his sufferings or his wants, we shut out from ourselves one great source of information which we can ill afford to lose; and that by putting aside that kind of knowledge of his state, which can come to us only through the workings of his own mind, we rob ourselves of those elements for a correct diagnosis, without a due establishment of which all our attempts at treatment must be either empirical or fruitless.

MODERN MEDICAL DOCTRINES.

Extract from the President's Address, delivered at the Conjoint Annual Meeting of the East Anglian and Cambridge and Huntingdon Branches, at Norwich, June 30th, 1871.

By PETER EADE, M.D.Lond., M.R.C.P.,
Physician to the Norfolk and Norwich Hospital, etc.

[AFTER thanking the members of the Branches for the compliment paid to him in his election as President, and offering them a welcome to Norwich, Dr. Eade gave a brief sketch of the history of the city, referring especially to its honourable position in the pursuit of the sciences and arts, and especially in the annals of medicine. He then proceeded to give a review of the progress of medicine during the year.]

With regard to the year that has passed, I would venture to express the opinion that, since our last anniversary, the science of medicine and its allied branches of knowledge have continued to progress. We may not have had such a valuable boon as chloral provided for us, nor such a great theory propounded as that of the safety-giving influence of carbolic acid after operations; but still I think that those who have perused our JOURNAL and the general medical literature of the country, will agree with me that there has been no check to the steady persevering work of our medical inquirers, and also that in this work various members of the British Medical Association have borne their full share. Even if nothing else had been done, the discovery of the inoculability, not only of portions of the integument, but even of the epithelial scales—of the very dust of our skins—upon the raw surface of another individual, would be almost enough to mark a distinct era in our advancing knowledge. Coming as it does upon the heels of Villemin's discovery of the inoculability of tubercle and other organised matter, there is no doubt it is helping to clear the ground for our thorough understanding of the pathology of pyæmia, of all cellular

and many contagious diseases, and—may I not say?—of phthisis itself. For it is very clear that, if we can see that isolated particles or cell-dust from the skin can live on a fresh surface and there grow and increase, there is no difficulty in believing that similar particles shed from the surface of the pulmonary mucous membrane can be inhaled by a second person, become transplanted on to his mucous membrane, and constitute there the starting point of a fresh and similar disease. I am of course aware that this possibility is by no means demonstrated, and that tuberculous particles differ considerably from ordinary nucleated epithelial scales; also, that there is much negative evidence to be adduced against this view. But when we remember that, whatever their microscopic characters, tuberculous matters have been proved to be capable of reproduction, and also call to mind the many instances which are constantly occurring of untainted individuals nursing the consumptive and themselves quickly becoming victims of the disease, there seems reason at least for believing that this is not unlikely to be shown hereafter as a true fact of this disease, and that the opinion long entertained of the occasional contagiousness of pulmonary phthisis is founded in truth. It is not long since the communicability of typhoid fever and cholera was vehemently contested. We now know (thanks, I believe, mainly to the researches of two eminent members of this Association—Dr. W. Budd, and the late Dr. Snow) that both these diseases are communicable by the reception into the system of particles—possibly mere poisoned epithelium—discharged from the mucous surfaces of diseased individuals.

It is really most lamentable that the true clinical history of such a disease as consumption, which destroys one-sixth or one-seventh of the whole population, and which is open to the observation of every one of us, should still be wrapped in such impenetrable darkness. The discovery of its inoculability into lower animals would seem to be the one ray of possible light which has lately been shed upon it—for the distinction of phthisis into several varieties, and the more accurate recognition of its various pathological forms, has as yet done little for its elucidation, and has not even enabled the best observers to agree as to the interpretation of the best known facts. In illustration of this, and of the consequent need for more work in this direction, hear what a reviewer of Niemeyer's recent *Lectures on Phthisis*, writing in the *Edinburgh Medical Journal* for December last, says. "There is no work", he writes, "on the treatment of phthisis in the English language so advanced in its pathology. It leaves the crude theories of Laennec and his followers far in the rear, and, by showing the essential dependence of tubercle on preceding inflammatory processes, it also shows, etc." Now contrast this with the views expressed by one of the most able and experienced physicians of our own country in the special field of consumptive disease. No longer ago than April last, Dr. Cotton wrote in our own JOURNAL to this effect. "Phthisis is, in my opinion, one of the diseases of organic degeneration, consisting essentially in the development or deposition of a certain lowly organised and variously constituted material, which we call tubercle. In the great majority of consumptive cases, I believe that such deposition is *not due to inflammatory action*—any inflammation, whether pulmonary, bronchial, or pleuritic, which may accompany it, being of a secondary kind." Dr. Cotton goes on to say that many cases of phthisis do begin with some form of pulmonary inflammation; but his general opinion on the subject I have quoted above in his own words.

Three or four years ago, at the meeting of our Association at Oxford, I read a paper in which I ventured to propound the opinion that a lowered vitality—nervous and anatomical vitality—of the pulmonary tissues was an essential element of the development of tubercular growths; and that, as tubercle itself was a low form of cell or nuclear growth, so it flourished only at best in lung-tissue, whose vital resistance was lowered by abstraction—either directly or reflexly—of nerve-power, so as to render it an appropriate soil for the development of this lower type of growth. All increase of knowledge since this time seems to have rather strengthened this view; and certainly a capability of previously healthy individuals for receiving and permitting the growth of inhaled tubercle-germs upon their pulmonary mucous membranes, when exhausted by nursing and anxiety, would not militate against it. Whether the result be actually induced in this disease, to quote the words of Mr. Paget, in his recent lecture on Dissection-Wounds, "by mere diminution of a normal power of resisting changes," or (as Dr. Carpenter has shown to be much more probable) by the production in the fatigued organs of some material on which morbid poisons may multiply and flourish, seems uncertain; but the tenor of this passage evidently is to show the belief of both these eminent physiologists and pathologists in the power of implantation and growth of living particles on or in lung-tissues, which they could not invade when these were in possession of their normal vitality. Other examples of this effect of lowered innervation in permitting the springing up of a lowered type

of growth are well seen in the hypertrophy of connective tissue, which accompanies the progress of the disease known as Duchenne's paralysis; in the formation of pus in parts where nerves are compressed or interfered with by tumours or injuries; and in the development of cancer elements in lymph exuded after an injury, and so placed beyond the reach of the necessary nervous control.

The great tendency of medical thought has of late unquestionably been in the direction of nerve-pathology and nerve-therapeutics. The nerve theory has in great measure—and doubtless rightly—superseded the humoral theory; and, as a result of the increased attention directed to the subject, the number of disorders found to be referable to some disease or derangement of nerve or nerve-centre, goes on ever increasing. The more, too, the ganglionic or sympathetic system is studied, the greater is found to be the number of ailments which are referable to a diseased condition of its elements. Take as a single example the great abdominal ganglionic centres, and see what a multitude of disorders can now be referred to their morbid conditions, some of which but a very short while ago were not only without a locality, but even without a name; Addison's disease; other forms of melasma; anæmia idiopathica; and probably hypochondriasis and some forms of jaundice; besides various reflected disorders, such as certain cases of mania, probably some examples of delirium tremens, and other familiar diseases and functional disorders, most of them characterised by epigastric sinking or other uneasiness.

But by far the greatest change of opinion has occurred as to the number and variety of symptoms and disorders which are now looked upon as due to causes acting at a distant part of the system, but producing their effects far from their source, through the medium of connecting nerves and nerve-tissues. Much, very much, has now been done in this direction. The work which was begun by Prochaska, continued and developed by Marshall Hall and Brown-Séquard, has indeed borne great fruit; but I venture to think that no limit can at present be set to the possible results of further investigations and inquiries in this direction. Not only have many hitherto unexplained affections been satisfactorily explained on this principle, but there seems reason to believe that the true *modus operandi* of the causes of many of our most common acute diseases may be—as some of our best thinkers have lately tried to show to be probable—through some agency of a reflex nature.

Take the simplest of all—the case of a *common cold*, with its resultant chill, and sore throat, and catarrh. It may indeed be true, as has long been held, that some resulting inefficiency of cutaneous action may allow of accumulation in the blood of perspiratory matters, and so give rise to some of the phenomena witnessed; but it is very plain that all the facts of this disease are by no means to be so explained, and that some further *rationale* of their action is required. Even if we leave out of the question the strange circumstance that the sore throat, or catarrh, or bronchitis, is frequently produced by merely getting the feet damp, or sitting for a few moments in a small current of air, or by exposure to other very partial cause, this old perspiratory theory—which was taught in our student days, I presume, to very many of us now present—fails to afford an intelligible explanation of the true nature of a simple cold.

But, if we allow a reflex explanation for a common cold, must we not at once agree to the further hypothesis, that all the so-called mucous inflammations of the respiratory tract—laryngitis, bronchitis, pneumonia—are or may be of the same nature and character, though differing in situation and intensity, and that they are in fact but varied results of a reflex paralysis—of a removal of the nerve-power of the vaso-motor nerves governing the vessels of the “inflamed” parts, and allowing the relaxation, congestion, and subsequent exudation of *liquor sanguinis*, in which this process of inflammation consists?

An investigation of the reason why removal of the vital heat, or vital force or nerve-power, from a given distant part, by moisture or current cold, should produce a catarrh, a cynanche, a bronchitis, a pneumonia, a rheumatic fever, would seem to be a subject worthy of the closest attention, and likely largely to increase our knowledge not only of these diseases themselves, but also of the direction in which nerve-force is apt to travel from various parts of the surface to other parts. In other words, it might lay bare the knowledge of the existence of great and definite nervous arcs or channels, of which at present we can only have a bare suspicion.

A common cold is the commonest, apparently the simplest, and certainly the most easily studied, of all these inflammations of the respiratory mucous membranes (the possibly partial reflex-paralyses. Might not the concentration of our inquiries upon the intrinsic nature of this disease unfold to us the secret history of the more serious chest-diseases, and point out to us not only their essential nature but the means of cure?

We all know and believe in the common adage, that we cannot cure a cold. Are we not fast coming also to the conclusion that we cannot cure a pneumonia, and is there any reasonable doubt that the tumid, red, congested condition, which we see in “inflamed” or catarrhal nostrils or throat, represents the exact condition of the mucous membranes lining the air-tubes in bronchitis or the lung-tissue in pneumonia? In this respect I think that practice is outrunning theory; for, whilst the essential nature of pneumonia is yet undetermined, the almost universal practice now in its treatment, in addition to the warmth of a bed, is the application of external warmth to the inflamed part by means of constantly renewed hot poultices. Possibly we might go further and do better, were we to apply this artificial warmth to the exact part from which vital heat or nerve-force was removed by the cold or wet which caused the illness. It seems reasonable to suppose that the same nervous arc or circuit through which the nerve-force was abstracted might often be the best channel for its restoration.

Assuming also that these internal inflammations are but forms of reflex paralysis, do we not see at once a reason why so-called remedies are often so useless for their check or cure, and why nothing but a certain period of rest and care, a period sufficient for the re-development of the normal nervous power of the diseased part, suffices for restoration to health? and why also rest, and chiefly physiological rest (as has recently been pointed out by Dr. Gull), is often almost the *only thing* that is needed for such restoration?

But, after all, I would ask, what is this doctrine of physiological rest, but stating in other words the need of time for the restoration of inflamed, *i.e.*, exhausted, or semi-paralysed parts? The “six weeks” that are said to be good for rheumatic fever; the four or five days that are known to be essential before a cold or a pneumonia can get “the turn” towards recovery; do but express the fact that this period of time is that in which the nerves of the part are able to recover from the exhausted condition which precedes, and allows the rise of, that particular form of disease. We may possibly neutralise the lactic acid of the blood in the one case, or the bilious or perspiratory poisons in the other, and so help to make the recovery more rapid; but in the absence of such or other complications, the rest and the vital nourishment by properly regulated warmth or temperature would seem to be the essentials of the treatment.

Before I conclude my address, there is yet one other thing worthy, I think, of mention in connection with this subject of nervous disorders, and also of some value in a practical point of view; and that is, that in considering disease and disease-causes, in this light of reflected causation, it may be worth while to remember that, as the bodies of the lower animals are not only divided into small sections, but also into the larger segments or divisions of head, thorax, and abdomen, so it is probable that some similar marked physiological division of *our* bodies also exists in the nervous sense, and that, though no anatomical line of demarcation is perceptible in the spinal cord, yet probably a definite physiological division exists defining the point of separation and of junction of the nerves appropriated to these three great segments of our bodies. Certain it is, at all events, that irritation reflected along the cord produces special symptoms which may in practice be distinctly classified in this sense, and of which we may perhaps get our best illustration if we call to mind the various symptoms produced by a *diseased or disordered uterus*, and observe how readily they may be grouped with reference to these three great segments of the body:—*e.g.*, the frontal headache, the vertigo, the puerperal convulsions or apoplexy, the mania, puerperal or otherwise, referable to the cephalic segment; the globus hystericus, the sighing and yawning, the intercostal neuralgia, the palpitation, the paralysis of upper limbs, referable to the thoracic segment; the nausea and vomiting, the abdominal pains and cramps, the pulsating aorta, and possibly the albuminous urine of pregnancy, referable to the abdominal segment. I add the albuminuria, for I, for one, am by no means satisfied with the explanation usually given of uterine pressure being the cause of this symptom, and would venture to express the opinion, founded on some observation, that not only albuminous urine, but also some of the graver seizures which are met with at the puerperal period, such as the comatose and apoplectic attacks (belonging to the cephalic segment), and the pleuritic and pneumonic attacks (referred to the thoracic segment of the body), are but examples of exhaustion of the nerve-force of the pelvic nerves producing as a result exhaustion by reflexion in these distant parts and organs.

A SON of Mr. Watkin Williams, General Secretary of the British Medical Association, has lately been elected by competition to a valuable foundation scholarship at Winchester College, being placed fourth in a list of twelve successful competitors. There were in all 111 candidates.