THE CIRCULATION IN THE ACARDIAC FŒTUS.

By GEORGE JOHNSON, M.D., F.R.C.P.,
Physician to King's College Hospital.

It is satisfactory to me to find that the correspondence between my Reviewer and myself, published some months ago, had sufficient interest for Dr. J. W. Ogle to induce him to write an elaborate paper on the subject of the circulation in the acardiac fœtus (BRITISH MEDICAL JOURNAL, Oct. 28th). I had looked up all the authorities now quoted by Dr. Ogle before I commenced the discussion; but I did not refer to them during the correspondence, because the question whether an acardiac fœtus had survived its birth could be decided without such reference; and, further, it appeared to me that the speculations of Dr. Houston and Dr. Marshall Hall only tend to throw a cloud of doubt and obscurity over a question which Sir Astley Cooper's dissection had rendered clear and intelligible. I will endeavour as briefly as possible to state the facts of the case.

There is no authentic instance on record of an acardiac fœtus unassociated with a cardiac twin. Sir Astley Cooper's dissection and drawings show a direct communication between the two umbilical arteries of the perfect fœtus and the single umbilical artery of its acardiac companion. It is obvious, then, that the blood must of necessity be forced by the heart of the perfect child along the umbilical artery of the acardiac fœtus, as it was forced *per saltum* from the placental end of the divided cord in the case of twins cited by Dr. M. Hall. It is true that this involves, as Dr. Marshall Hall says, an inversion of the order of the circulation in the cord of the acardiac fœtus, the blood passing into that fœtus by the umbilical artery, and out by the vein; but this is a necessary result of the fact that *the heart which propels the blood through both fœtus is outside the body of the imperfect fœtus*.

The movement of the blood and the direction of its current are determined by the heart. *The blood moves from the heart through the arteries; it returns to the heart by the veins.* This is a law without exception. It is manifest that Dr. M. Hall's hypothesis of "lateral action" involves a reversal of this law, and makes the blood move from the heart through the umbilical vein of the acardiac fœtus, and return towards the heart by the umbilical artery. This is an incredible hypothesis, inconsistent with anatomical facts and with hydraulic principles. The only part of the circulation in which the ordinary course of the arterial and venous currents is actually reversed is in the umbilical cord of the acardiac fœtus; and this reversal, I repeat, is necessitated by the fact that, one heart having to serve for two fœtuses, the central organ of the circulation is external to the imperfect fœtus.

Dr. Houston's suggestion that, notwithstanding the free communication between the vessels of the two fœtuses, the heart of the one has no influence upon the circulation of the other, but that the placenta has some mysterious power of attracting and repelling the blood, is obviously an untenable hypothesis. Such speculations as those of Dr. M. Hall and Dr. Houston, in opposition to the facts so clearly demonstrated by Sir A. Cooper, remind one of the curious hypothesis of Riolaus, of whom Harvey says, that "he would have the blood to make its way into the left ventricle through the septum of the heart by certain invisible and unknown passages, rather than through those ample and abundantly pervious channels, the pulmonary vessels, furnished with valves, opposing all reflux or regurgitation." (Harvey's *Works*, Sydenham Society's translation, p. 99.)

Finally, I would suggest that our present knowledge of the functions of the vaso-motor nerves and the muscular arterioles necessitates a thorough revision and modification of the theory of a capillary force and a *vis à fronte*. The case of an acardiac fœtus obviously affords no support to that theory. On the contrary, the invariable vascular connexion between such a fœtus and a fully developed twin is one amongst a multitude of facts tending to show that, in the circulation of the higher animals, while other influences regulate the supply and distribution of blood, the heart is the main and essential motor power.

OBSERVATIONS ON THE RADICAL CURE OF RETROFLEXION OF THE UTERUS.

By PERCY BOULTON, M.D., etc.,
Physician to the Samaritan Hospital for Diseases of Women.

DURING the last four or five years it has fallen to my lot to treat a large number of cases of retroflexion of the uterus; and, after many trials of various stems and pessaries, I have arrived at a plan which seems in every way satisfactory. In the JOURNAL of September 23rd I

see a report of Dr. Beatty's paper, which I regret I did not hear at Plymouth; and I wish to record my experience without further delay.

I would commence by endorsing all the advice given by Dr. Beatty as to the preliminary treatment in order to reduce inflammation, congestion, ulceration, etc., before introducing any stem. When the uterus is in a proper condition for receiving the stem, the question is, What kind is the most suitable? Dr. Beatty uses one of Sir James Simpson's, which requires "a flat boxwood pessary in the vagina" to keep it *in situ*. I object to this pessary for the following reasons.

1. The wearing of a stem invariably causes an increased flow of catamenial and other discharges; and such a pessary must retain these in the vagina.

2. It must effectually prevent the passage of the weak astringent wash which Dr. Beatty recommends to be used daily.

3. No examination can be made without removing the pessary; and, "as the bulb of the stem is to rest and move on the smooth surface of the boxwood pessary, and thus allow the uterus to change its position as the woman is horizontal or perpendicular," I think it more than likely that the uterus would become retroverted, and the stem slip out.

4. In most cases a pessary is unnecessary.

My plan is to introduce a self-retaining stem, thus doing away with the necessity of anything in the vagina. I have used various kinds, made of vulcanite or German silver. Those of Drs. Sims, Routh, Bantock, and Greenhalgh, have been preferred; and of these the latter is by far the best. The advantages are, that it is self-retaining, cleanly, being made of German silver, and allowing free escape of discharges; it is easily introduced; and the spring is stronger than the vulcanite ones, and therefore less likely to slip or get out of order.

The first fault I have always recognised. The stem, as ordinarily used after the operation of hysterotomy, is too large to be used in most cases of retroflexion; and I have, therefore, been in the habit of introducing stems identical in shape, but only two inches and a half in length, including the rounded head; and passing the whole within the os uteri. At times the edges of the head have caused pain to the wearer for a short time; but in most cases it has appeared to fit quite naturally within the widened cervix; and the absence of anything in the vagina has always seemed a great advantage. I have used this stem more than any other, and never had any difficulty with it till August last, when the following case came under my care.

Miss E. D. was suffering from melancholia, which was distinctly of a hysterical character. On examination, I found the uterus retroflexed, the fundus forming a tumour between the rectum and cervix. On August 3rd, I passed the sound two inches and a half, straightening the uterus; and then I inserted one of Dr. Greenhalgh's spring stems two inches and a half in length. The patient was anxious to go home to the country; and, as her mental condition rendered it advisable that she should be under constant surveillance, I consented to her doing so, cautioning her to keep very quiet. On September 14th, she came to London to have the stem removed. She had been wonderfully well, and had felt no inconvenience. When I examined the uterus, it appeared perfectly straight; but the stem had gone. I concluded that it had slipped out when at stool, and had not been noticed, as several of the vulcanite ones have done with other cases of mine. I, however, passed a sound gently, and discovered that the stem was in the uterus, but had worked up, the head being nearly an inch from the os externum. I had not much difficulty in extracting it with a copper sound bent into a short hook at the end, which I passed at one side of the stem, and then turned and drew it away.

This case is instructive. The pointed extremities of the open stem must have passed up the cornua leading to the Fallopian tubes; and such a contingency is worth remembering in the future; for, although in this instance the stem was extracted, it might not always be easy to do so.

Not wishing to give up the stem that has hitherto served me so well, I have had a sheath made for the head, with neck and shoulders attached; and this does away with the only faults I have yet discovered. The smooth oval sheath over the head, reduced in size, protects the cervix from the sharp edges which used to cause pain to the wearer; while the shoulders prevent it from working higher up into the uterus, and render it easy to extract. This adaptation of mine makes the stem somewhat resemble Dr. Bantock's vulcanite one; but it is infinitely preferable, and leaves, I think, nothing to be desired.

In most cases of retroflexion, a self-retaining stem is all that is necessary. I have had cases, however, where, owing to increased size and weight of the uterus, from congestion or other causes, or from difficulty of defecation, there has been partial prolapse. In these I always use a Hodge's pessary after replacing the womb and introducing the stem. This takes the strain off the uterine ligaments, and answers every pur-