discharge was kept up from the skin, with so happy a result that, fifteen months later, I attended her in a confinement, and she has lived six years since.

A young woman, aged 19, presented herself to me, with extensive tubercular disease of both lungs, which had been running a constantly progressive course for more than two years. She was extremely emaciated, had profuse perspirations, and her legs were cedematous. Her case was one of advanced tubercular bronchitis. I put a seton of six silks in her chest, and at the end of a fortnight the symptoms began to abate. She wore the seton three months, by which time her cough had nearly ceased, and her expectoration was *nil*. She then pressed me to remove the seton, and, upon inquiring the reason of her urgency, she informed me that her banns had been twice asked, and she intended to be married the following week. She *was* married to a worthless fellow, who used her very badly; and, after an interval of eighteen months, I was called on to attend her in an attack of acute tubercular pneumonia, under which she soon sank.

I will not weary you with the details of cases. My memoranda supply me with the records of thirty-five cases of confirmed phthisis in various stages, in thirty-two of which marked benefit was derived from the employment of setons, while in three the seton was removed without benefit. In none of these cases had I reason to regret the introduction of the seton: emaciation and debility were not increased by the discharge from the skin, but, on the contrary, their progress was arrested *pari passu* with the diminished discharge from the lungs.

Passing to another variety of strumous disease—*strumous ophthalmia*—an issue or seton in the temple is the most effectual remedy, and often the only means of stopping the progress of the disease. Probably in the present day setons and issues are used more often in the treatment of strumous ophthalmia than in that of any other disease; and I take it that it is so because their remedial power can, in the eye, be actually seen and appreciated.

It is not only in the true strumous ophthalmia, attended with corneal ulceration, that setons are of service—they are equally so in all those minor affections of the eye and its appendages which take their origin in a strumous diathesis.

For the relief of an eye-affection, the seton is usually placed in the temple; but a serious drawback to its use in this situation is the scar which it leaves. I find that, placed in the arm or any convenient part of the body, it has an equally good effect.

Some ten years ago, a girl was under my treatment for many months in consequence of frequently repeated attacks of strumous corneitis. Having failed to eradicate the constitutional tendency by medicines, I put a seton in each temple. The result was a complete cessation of the disease; but, having to earn her bread as a nursemaid, she was naturally anxious to get rid of a disfigurement which prevented her from taking a situation, and, contrary to my advice, removed the setons at the end of a month. Not many weeks afterwards, exposure to cold wind brought back the disease, which again assumed a remittent character. Being in a situation at a distance, it was only when compelled to return home that she again became my patient. I then introduced a seton of six silks into her arm, and kept it discharging the whole winter, during which time she was entirely free from the eye-affection, and has never had a serious return since.

A young gentleman seldom passed a month or six weeks for several years without an attack of strumous conjunctivitis. Being first at a private school, then at Eton, and latterly in London, he had been subjected to various plans of treatment. The attacks yielded sometimes to leeching, sometimes to blistering, and at other times to sedative applications; but exposure to cold wind or damp air always induced a recurrence. At last I persuaded his friends to allow me to put a seton in his arm. The suppuration was so severe at first that I had great difficulty in inducing him to retain it; he did so, however, and by degrees the fungoid granulations which had sprung up round the points of insertion disappeared, and a healthy suppuration was established. From the time the seton was inserted the ophthalmic attacks ceased; during the three months he wore it they did not once recur; and for three years since he has had but the most trifling threatenings of his old malady.

Two years ago, a younger brother of the same family was threatened with the same affection. I immediately put a seton of one silk in his temple and stopped the disease, and he has had no recurrence.

A third form of strumous disease, in which I have used setons with great advantage, is tubercular meningitis. My attention was first directed to the beneficial effect of a discharging surface in this disease by a case which occurred to me, in company with the late Dr. Symonds, soon after I commenced practice. A child three years of age, apparently in the last stage of hydrocephalus, was accidentally scalded on the chest, and a large suppurating surface was the result. Dr. Symonds

was fully persuaded that the case was hopeless, and was equally convinced, when the child recovered, that the extensive suppuration turned the balance in its favour. Since then I have treated a considerable number of cases of this disease by the introduction of a seton in the poll, and am fully persuaded that, when the attack is sufficiently prolonged, the establishment of a free discharge offers the best hope of cure.

There is yet another form of struma in which I have used setons with advantage—viz., chronic glandular enlargements. I have found obstinate enlargement of the axillary glands yield to a seton in the arm, or below the clavicle. A seton above the clavicle or in the arm will facilitate the removal of enlargement of the cervical glands; and not long since a most obstinate enlargement of the inguinal glands, which for many months resisted all treatment, eventually yielded to a large seton in the arm, kept discharging for eight weeks. Lastly, I may mention chronic strumous ulcers as curable, when other means fail, by a discharge established at a distant part of the body.

I have avoided extending this paper by the narration of cases, and have omitted the mention of any concomitant treatment. I would not have it supposed on this account that I advocate the use of setons to the exclusion of other remedies. I apprehend the part the seton plays is that of evacuant—distilling, as it were, from the blood the morbid material. It remains to check the reproduction of this material by catalytic remedies.

My apology for calling the attention of this meeting to a subject which is probably so familiar to all present, is the very good result which I have experienced in my own practice from its employment.

A SUCCESSFUL METHOD OF TREATING CERTAIN CASES OF DYSMENORRŒA AND STERILITY.*

By PROTHEROE SMITH, M.D.,

Senior Physician to the Hospital for Women, London.

To treat of the subject of dysmenorrhœa generally, the time to which I am restricted would be insufficient. I shall, therefore, now only speak of that form of the malady which is the result of abnormal constriction of the uterine canal, and which also frequently presents a mechanical obstacle to pregnancy. With this condition, known as obstructive dysmenorrhœa, we are all more or less acquainted, since it is of frequent occurrence amongst the unmarried and the sterile.

The lumbo-abdominal distress, the bearing down and forcing uterine effort immediately preceding the menstrual flux, and the experience of relief when it is established, with the constitutional debility which follows, with consequent neuralgic headaches and pains in the hypogastrium, sufficiently characterise the disease; while chronic and long standing, pains in the *bas ventre*, especially on the left side, continual backache, tenaceous leucorrhœal discharge, with an aggravation of constitutional disturbances, mark the progress of the malady. The diagnosis, however, is complete when, by touch and uterine sound, the constriction of the canal is proved. This condition was described by Dr. Mackintosh, of Edinburgh, in 1844, who proposed as a remedy the use of bougies, so as gradually to effect the dilatation of the straitened passage. This plan of treatment was frequently attended with success, and was, about the same time, modified and improved by the late Sir James Simpson, who used metallic stems two inches and a half long, having a button or bulbous end, by which these "short sounds" were retained and secured. Such means, at times, seemed to answer; but, unless pregnancy soon followed, the constricted parts of the uterine canal resumed their abnormal condition, and called for a repetition of the treatment.

In Dr. Black's excellent edition of the *Selected Obstetrical Works of Sir J. Y. Simpson* (vol. i, p. 679) published this year, is the following paragraph.

"At a meeting of the Edinburgh Obstetrical Society, March 10th, 1847, Dr. Simpson stated that he had now been in the habit for three or four years past of performing the operation of incision of the cervix uteri for obstructive dysmenorrhœa. He first described the operation to the Medico-Chirurgical Society in 1844, and it had latterly been adopted by Dr. Rigby, Dr. Protheroe Smith, Dr. Oldham, and other accoucheurs in London and elsewhere."

The instrument used by Sir James was a kind of *lithotome caché*, by which he divided the strictured part, the incision commencing at the union of the cervix with the body of the uterus, and passing more and more into the substance of the cervix as the instrument was withdrawn, and so dividing its lower edge.

In the year following, I had a double hysterotome constructed in

* Read in the Midwifery Section at the Annual Meeting of the British Medical Association in Plymouth, August 1871.

Paris, with two blades cutting laterally; and Dr. Greenhalgh subsequently contrived a very ingenious instrument of this description, by which he proposed to gauge accurately the extent of the incisions on both sides. Dr. Black, however, observes that—"In consequence of the gratifying results sometimes produced by incision of the cervix uteri in obstructive dysmenorrhœa and sterility, that operation became an extremely favourite one with Simpson. Certain risks connected with it, however, and, in particular, its liability to be followed by pelvic inflammation, inclined him ultimately to a rigorous selection of cases, and to the enjoining of recumbency for several days after the performance of the operation."

Stimulated by Sir J. Simpson's favourable opinion, for some time I followed this mode of treatment; but the results, as regards the removal of dysmenorrhœa and sterility, were not such as to convince me of the advantage of its general adoption. On the contrary, in several instances, severe metrocervicitis occurred, together with considerable hæmorrhage, requiring the plug to control it. In others, the painful menstruation was unrelieved, and the patient remained sterile. My objection, however, to the use of the hysterotome was confirmed by observing that, in several instances, it had aggravated the evil; for not only did the patient continue barren, but the original constriction was increased, and, with this straitened state of the uterine canal, the consequent distress was augmented. The pathological condition I discovered by the use of sea-tangle tents. The first case which yielded this information to me was one in which Sir J. Simpson had performed the operation some years previously. For a time, after recovery from its immediate effects, relief from the distressing symptoms was experienced by the patient; but after a short time the dysmenorrhœal sufferings returned, and, as they gradually increased in severity, she was led to consult me. Finding that the uterine sound could be passed only with extreme difficulty through the os internum uteri, I introduced a small sea-tangle tent, which, on the following day, I had considerable difficulty in withdrawing. It was only after continued traction, kept up steadily for nearly twenty minutes, that this was accomplished. When removed from the grasp of the cervix, its form explained the difficulty. At an inch and a half from its proximate end, the tent was not altered in form from its size before it was introduced; whilst above and below this narrow waist, it had swollen to its utmost extent; from this compressed part ran a narrow groove on each side, evidently showing that, at the constricted os internum, as well as longitudinally on each side, there existed unyielding tissue, not only at the points of the original stricture, but also at lines corresponding with the wounds made by the hysterotome, which led me to the conclusion implied in these remarks—viz., that the deep grooves and the longitudinal furrows were the impression of hard cicatrices, the effects of the previous operation. Seeing that the result of incising the cervix, if successful in enlarging the canal, must be to destroy the perfectness of its structure to the extent of the tissue divided by the hysterotome, or that, if rendered unsuccessful by the healing of the wound, it might leave the patient worse than before, from the hardened cicatrix which might follow the operation, I soon adopted a course which I have pursued for many years with a success which encourages me to hope that its announcement will be acceptable to obstetric practitioners generally.

In my lectures, at the commencement of this year, on Flexions, Torsions, and Displacements of the Uterus, at the Hospital for Women, I observed that, "finding sterility and dysmenorrhœa were often benefited by dilating the cervix uteri by the bougie, after the manner advocated by Dr. Mackintosh, and with a view of improving upon the plan suggested, I got Messrs. Fergusson, instrument-makers to St. Bartholomew's Hospital, to make me in 1841 the instrument I now exhibit, after the model of Heurteloup's lithotrite, by which the extent and direction of the uterine cavity was easily measured, and a constriction in any part of the passage as readily overcome." With this uterine dilator, I conceived it might be practicable to dilate permanently the constricted os internum, and afterwards, when necessary, to give the normal shape to the os tincæ by dividing it laterally at the commissures of the labia uteri *per speculum*. I have now adopted this mode of treatment for more than a quarter of a century; and although, in some instances, I have failed to cure, yet I have succeeded in so many others, that I feel warranted in advocating its adoption in the selected cases in which it is eligible. To diagnose such, it has been my custom to ascertain from the history of each case, as well as by a careful physical examination of the uterus, that it is one simply of stricture of the os internum, and narrowing of the cervical canal and mouth. But should there be any inflammatory condition, whether of the mucous lining or of the deeper tissues of the uterus, or any thickening or induration of these parts, it is essential to success, however tedious the process may be, to remove such morbid conditions before adopting the *extension forcée*, by which the stricture is to be overcome.

The plan which I then pursue is, first to prepare the patient by a purgative dose, and by abstinence from local excitement, and from alcoholic drinks, or much animal food. When hyperæmia exists, I scarify laterally the labia uteri repeatedly at the commissures of the labia, by which the vascularity of the organ is reduced, and the shape of the os tincæ, when constricted, is improved. After accustoming the uterine canal to bear a metal bougie, which should be repeatedly and daily introduced, and increased in size until that of a No. 10 catheter can be borne without any pain, then the uterine dilator may safely be employed. It will be seen that this instrument consists of two short blades, two inches and a half long, the inner being continuous with the sliding shaft, with which it is nearly at a right angle, having at its proximate end a screw worked by a nut so as to mark precisely, by an index on the handle, the extent of dilatation employed. This should be used at first cautiously about every second day, always ceasing to screw as soon as pain is experienced. This is immediately relieved by a turn or two of the screw the reverse way.

It will be found, in a short time, that the uterus becomes accustomed to the dilatation, when it may be employed to a greater extent; and in the course of a few days or weeks, as the case may be, a forced dilatation to the extent of an inch or an inch and a half may be used with impunity. After this it will only be necessary to use the dilator daily for two or three days, and afterwards at longer intervals, to keep the parts open till they permanently heal in the state of distension effected by the operation.

Should any congestion or inflammation result, scarifying at the commissures of the labia will relieve by free bleeding, whilst, at the same time, the os tincæ is made to assume a more open, and, therefore, a more normal shape. I prefer, generally, to effect this by the repeated use of a small scimeter-shaped knife, as I find that by so doing the risk of inflammation is diminished, and it prevents cohesion of the cut sides of the labia uteri, both which accidents occasionally attend the operation when performed at once by a hysterotome. When preternatural shortness of the uterus, from original malformation, exists, the operation is contraindicated; also when stricture depends on endometritis; when, otherwise, metritis or metrocervicitis is present; when there are fibroid tumours causing inflammatory adhesions; when dysmenorrhœa is characterised by deciduous membranes; when there are conical hypertrophy and elongation of the cervix; when there is globular enlargement of the anterior labium uteri, embraced by the posterior lip in the form of a crescentic membrane; and when displacements and dislocations of the organ complicate the case—these and all other organic diseases which may attend this malady should be removed prior to the adoption of forcible extension by the dilator.

Time would fail me, were I to attempt to illustrate these remarks by a review of all the cases in which I have employed this mode of treatment. It must here suffice to remark that, I have seen this practice followed by complete success in many instances, from amongst which I would quote the following extreme, but successful cases of dysmenorrhœa and sterility, treated by forcible extension and incision of the labia uteri.

CASE I.—A lady, aged 41, after sixteen years of married life without pregnancy, has since given birth to two children.

CASE II.—A lady, aged 37, her husband hemiplegic from his youth, married thirteen years without family, and now has a living child.

CASE III.—A patient, aged 39, married twelve years, and barren, became the mother of a living child.

CASE IV.—The wife of a clergyman, aged 28, having been sterile for six years after marriage, was delivered of a living child.

CASE V.—A resident in a distant colony, married fifteen years, and barren, aged 38, became pregnant, but miscarried, in consequence of over-exertion at the second month of utero-gestation.

CASE VI.—A delicate lady, aged 29, married nine years, but never pregnant, became so, and had a living child, followed by a second two years after.

I select these few cases from a large number, because in all there existed more or less dysmenorrhœal symptoms, and three of them were complicated by considerable structural disease, which had first to be removed before the plan of treatment I now advocate was adopted.

In this paper I have endeavoured, as far as possible, to confine my observations to facts capable of demonstration; and I hope the experience of other members of this Association may confirm my own conclusion that forcible distension of the constricted os internum and incision of the os externum only, in the manner described, in cases of obstructive dysmenorrhœa and sterility, is preferable to the operation by the hysterotome. The former is attended with less risk; and, with this impunity, it offers greater success, with the further advantage of leaving the uterine tissue, excepting at its extreme mouth, without lesion.