TWO CASES OF PUERPERAL FEVER:

WHICH RECOVERED UNDER THE REPEATED LOCAL USE OF A STRONG SOLUTION OF CARBOLIC ACID.*

By PETER EADE, M.D.Lond., F.R.C.P.,

Senior Physician to the Norfolk and Norwich Hospital, etc.

CASE I.—Mrs. C., aged 42, was seen (with Dr. Beverley) on October 13th, 1878. She had been confined of her first child nine days before; had felt ill on the following day; and had ever since been becoming gradually worse, with chills, sweating, continued fever, with a temperature of from 102° to $103\frac{1}{2}^{\circ}$; and, latterly, with diarrhoea and a tumid abdomen. There had been a slight secretion of milk, and a small lochial discharge, which was faint and disagreeable smelling. The catheter had been required for the relief of the bladder. She had been very sleepless. On examination, the uterus was felt large, and high up in the abdomen, rather more on the left than on the right side; and, on vaginal examination, its os, cervix, and lower part were seen to be large, swollen, and of a peculiar dirty aspect. No direct cause for the illness was known; but the drains were found to be in a faulty condition. The diagnosis now made was puerperal fever, or puerperal septicæmia; and the patient was ordered to take a teaspoonful of oil of turpentine every three hours; whilst we applied glycerine of carbolic acid freely to the outside and to the whole canal of the cervix uteri.

On the following day, she appeared to be a little better. She had taken five doses of turpentine without sickness, and without further purging. The pulse was 120; the temperature 103° . The milk was entirely gone. The lochia were rather freer, and less offensive. The uterus was less high in the abdomen; its os and cervix were sloughy-looking, and covered with bloody discharge. The uterine cervix and its canal, quite up to the cavity of the uterus, were again well smeared with the strong solution of carbolic acid, and the turpentine was continued at longer intervals.

Next day, although the turpentine was disagreeing, and had to be discontinued, she was evidently better; but the local application of the carbolic acid was continued night and morning; and, on the following day (October 16th), the os and cervix uteri, though still retaining the peculiar sodden and ash-coloured appearance which had been noticed, were evidently cleaner; and, at several spots, florid granulations were appearing in the midst of this dirty tissue.

From this time, she steadily improved. The carbolic acid was applied for several days longer; and, though her convalescence was retarded by a tiresome and irritating eruption of miliaria on the chest and upper body, yet, in about a month, she was reported to be perfectly recovered.

CASE II.—Mrs. F., aged 24, was seen (with Mr. Muriel) on October 27th, 1880, four days after her confinement with her first child. The labour had been natural, except for a slight laceration of the perinæum. At this time, the symptoms were pain in the abdomen, which was swollen and conical in form; dorsal decubitus; apathetic manner; a sweating skin; absence of milk from the breasts; purging of liquid pea-soupy stools; and scanty, mucous, and slightly offensive lochia. The pulse was 120; the temperature 102.4° . The uterus could be felt large, soft, and tender, five inches above the pubes. There had been retention of urine since delivery. On vaginal examination, the os and cervix were seen to be soft, flabby, and swollen; the os very patulous. The vagina also was swollen, and, like the large os and cervix, presented a peculiar patchy appearance—i.e., with sloughy grey and red spots or patches of mucous membrane intermingled. A dirty ropy discoloured mucus was oozing from the cervical canal. The patient had been taking opium freely for her pain; she had taken also several drachm-doses of oil of turpentine, and had been freely and frequently syringed out with a strong watery solution of carbolic acid; but she was evidently rapidly getting worse.

The whole outside and inside of the cervix of the uterus, and even far up into the uterine cavity, as well as the upper part of the vagina, were now freely mopped over and smeared with a solution of equal parts of carbolic acid and glycerine; the opium and turpentine, and carbolic lotions, were ordered to be continued; and poultices were applied to the abdomen.

On the following day (October 28th), she declared herself to be distinctly better, and to have less pain. The pulse was still 120, and the temperature 104° . The swelling of the abdomen and of the uterus were slightly less. The bowels were still relaxed. There was more of the

dirty vaginal discharge. The cervix uteri, examined, was found to be rather less grey or ash-coloured; but still large, swollen, pappy, and with a patent os. In spots where the carbolic acid had been freely applied, some florid red granulation, inclined to bleed, was showing itself. The whole upper part of the vagina, and outside of the cervix, as well as the cervical canal, far up into the uterus, were again freely brushed over with glycerine of carbolic acid, and the same application was repeated in the evening.

October 29th. She had passed a better night, and had slept for two hours at a time. The abdomen was smaller. There was scarcely any diarrhoea. The pulse and temperature were as on the previous day. She had passed urine without the catheter. She said she felt better. The uterine discharge was less. It was of more natural quality, and less offensive. The os and cervix uteri still looked of a dirty grey colour; but were markedly less flabby, more contracted, and firmer.

Some of the vaginal discharge was now collected, and examined under the microscope. It was dirty and purulent, with a mixture of viscid stringy mucus, and showed an abundance of pus-globules, chiefly large, many large granulation-cells, elongated fibre-cells, and vaginal epithelium. No bacteria could be discovered. The strong solution of carbolic acid was repeated night and morning.

On the following morning (the third of the treatment), the pulse had dropped to 92, and the temperature to normal; and the other symptoms were improved, though there was still some diarrhoea. The acid was applied locally for the last time on the evening of this day, as she was beginning to object to the introduction of the speculum, and the uterus was becoming almost healthy in appearance; but the syringing was continued a little longer.

In reference to causation, it should be mentioned that a case of puerperal septicæmia had occurred immediately before, in the practice of the medical attendant of this patient, the subject of which had died in the ordinary way.

REMARKS.—I have, in the last two years, seen five cases of "puerperal fever", besides one case of acute scarlet fever, immediately after delivery. Of the five, one was a case of septicæmia, where the stress of the disease fell chiefly upon the lungs. Two were cases where inflammation of the body of the uterus, and of the peritoneum, appeared to be the chief local phenomena; and the other two were the cases related above, where, in addition to other symptoms, there were marked vaginal and cervical septic changes. Of these cases, the scarlatinal patient recovered; and of the other five, the three first died; whilst the other two, which were thoroughly and repeatedly dressed locally with strong carbolic acid, recovered.

My first definite ideas as to the possible efficacy of treating the lower and exposed portions of the human uterus, in at least the communicated cases of "puerperal fever", with a very strong solution of carbolic acid, were derived from some communications embodied in a pamphlet (probably well-known to members of this Society) by Mr. Woods of Merton, on the subject of *Fever or Straining in Ewes after Lambing*. This disease of "straining" was a very fatal one; and had been attributed to bad lambing yards; i.e., to yards which had become foul and tainted—in this respect being exactly analogous to the tainted wards of crowded or infected lying-in hospitals. The term "straining" is stated by the eminent veterinary authority, Mr. Fleming, to include metritis, metro-peritonitis, and parturient fever—terms which express the most usual pathological conditions in human puerperal fever.

The treatment of this disease, recommended in Mr. Woods' valuable work, is simply the smearing of the uterus and vagina of the ewe with carbolic acid mixed with oil, in the proportion of one part to seven; and the most satisfactory results are stated to have been obtained from the practice. Moreover, quite recently Mr. Woods has informed me, that whereas formerly a very large proportion of the ewes attacked died, now that carbolic oil is freely applied to the hands of the shepherd-accoucheurs, and to the vagina and uteri of the sheep, death is the exception.

Such results as these have, of course, attracted much attention; and this veterinary experience has, I believe, induced much thought as to the applicability of the principle involved to the human parturient subject. The carbolic acid has, however, I believe, in women been chiefly used as an injection, and this only in a dilute form. For example: Dr. Matthews Duncan has lately recommended, for washing out the uterus, a solution of one in thirty parts of water; and Professor Stadfeldt of Copenhagen, who has written on this subject, has only used a three per cent. solution. The plan adopted in my cases was based upon the principle, not only of washing away foul secretions and decomposing matters, but of attempting, by the application of the strong—almost caustic—solution of the acid, to destroy the infected and still infecting local surfaces, with their multiplying germs, if such there were; and I

* Read before the Norwich Medico-Chirurgical Society.

have not seen any record of this more heroic practice having been hitherto employed.

The first of these cases occurred as long ago as October 1878, and I then hesitated to report it; because I thought it better, for various reasons, to wait for the confirmation, or the contrary, of the advantage of the local use of these strong solutions of carbolic acid to the uterus in "puerperal fever". Now, however, the satisfactory result of this second case makes me hesitate no longer to report the two.

It may not be a waste of time here briefly to recall the present views as to the group of diseases classed under the generic name of puerperal fever. Dr. Matthews Duncan, in his admirable paper, recently read before the Birmingham Medical Society, points out that, as now understood, puerperal fever is not a simple entity, but that a case to which we give this name may be one, either, *a*, of traumatic inflammation, with general fever; *b*, of poisoning of the blood by absorption of putrid matters or discharges (sapremia); or, *c*, of infection of the blood by living germs or micrococci (septicæmia, pyæmia); the distinction, however, between these varieties of postparturient fever appears not always to be very well marked. And Dr. Playfair says, it is probable that two distinct groups may be formed—namely, those in which the patient is poisoned by her own septic matters; and those in which the septic poison is conveyed from without.

Let me further quote from Dr. Playfair's work on *Midwifery*, as to the pathological appearances of the vagina and uterus, in septicæmic cases, at the site of absorption. He says: "These are described by pathologists as identical in their character with the inflammatory oedema which occurs in connection with phlegmonous erysipelas." These seem to consist of swelling, oedema, diphtheritic-looking exudations, and softening of the vagina, of the cervix, and of the membranes of the uterus, as well as of the uterus itself—conditions which were present in both the cases now related, as observed during their progress.

The fact being allowed that the profession generally is fully alive to the great advantages of antiseptic midwifery, at least in doubtful conditions, and to the special utility of carbolic acid, the question arises, What is the best mode of employing this agent? I would venture to suggest that both the use of intra-uterine dilute injections, and of the local application of the strong acid, may each be best in varying cases. Where the septicæmia arises from decomposing placenta, membrane, or clot, within the uterus, it would seem probable that a full and continued stream of the dilute disinfecting fluid may be most useful; but, where we have a specifically poisoned surface to deal with, in the inoculated cases, that there the more potent application may be the best, and indeed the only reliable, mode of treatment. And this is indeed my own opinion, derived from general observation, as well as from the daily observed influence for good of the acid upon the sloughy tissue. In my cases, the acid unquestionably caused a diminution of swelling, a lessening of the oedema, and an improvement of colour, in the parts to which it was applied; and knowing as we do its destructive influence upon germ-life, and the equally destructive effect of these germs, when living and multiplying, upon the tissues which they invade, we may fairly consider such a theory reasonable, and probably true.

In any case, the two patients lived, whilst others died; and, with the confessed unreliability of leechings, turpentine, iron, quinine, and every other remedy, and in view of the terrible and intractable nature of puerperal fever, I think that any light, however small or vague, thrown upon the best method of treatment by the recovery of the patients, is not only earnestly to be welcomed, but to be pursued, as possibly leading to a more definite knowledge. I have good hope that the path here indicated may be in some respects the true one.

SOME CASES TESTING THE ANTISEPTIC METHOD.

By ALBERT WILSON, M.B., Leytonstone, N.E.

THE conflict existing between the opponents and the supporters of antiseptic surgery has done a great deal to forward the views of the latter. Whilst scientific evidence supports the theory, and careful trial proves the practice, nothing but the experience of various surgeons can be brought against it. It is surprising in this advanced age to hear surgeons condemning such a system, which requires so great and scrupulous care for success, merely because it has not succeeded in their hands. They forget their own fallibility, whilst overlooking the infallible and immutable laws which govern the universe, even in the minute and invisible. At the same time that these men condemn this system, they unconsciously testify to the truth of the principle, by their fear to do operations which they see antiseptic surgeons perform without hesitation or anxiety, and by their delay till the last extremity to operate in

other cases, as chronic abscesses, knowing full well that, from the day they operate, the patient will descend to a slow but certain death. We, who contend for that grand system of antiseptics, which is one of the greatest boons to humanity that surgery ever conferred, do not fight for one method so much as for the principle. If a new method were introduced, provided it were as perfect as that now employed, and more convenient, then it would be adopted in preference; and I venture to say by none more readily than by the founder of the antiseptic system himself.

Fully aware of its difficulties and drawbacks, Professor Lister has spent years of untiring zeal and energy, not only in perfecting, but also in simplifying it. He has now obtained for healing wounds much the same conditions as if they were subcutaneous. What more can be desired than this to justify the statement that the system is nearly perfect?

Many object to practise antiseptics because it is an "awful nuisance" to carry a spray about, and waste of time is caused by the spray and attention to other details. Doubtless, these are inconveniences, but they ought not to be obstacles, considering the brilliant and certain success which will be the reward for such extra energy and patience, not to speak of the gratification afforded by saving life or preventing unnecessary suffering.

I will briefly relate some of my results while opposed not only by the inconveniences met with in private practice, but also by extremely unfavourable hygienic conditions, such as I think could hardly be exceeded.

Whilst resident at the Cowgate Dispensary, in Edinburgh, I did several operations which thoroughly tested the efficacy of the antiseptic treatment.

CASE I.—A woman, aged 23, had an exostosis growing from the internal condyle of the femur. I removed it on January 26th, 1879. In chiselling off the tumour, I removed a piece of the external shell of the femur. The woman was in very destitute circumstances, she and two others inhabiting a small room up one of the narrow closes or wynds of the Cowgate. A dingy window with four broken panes looked out on a small stable-yard, where they were loading manure within five yards of the house. The broken panes were stopped up with garments, and the operation was performed in an atmosphere most unfavourable. The wound was stitched with catgut, and a drainage-tube inserted. A gauze dressing was applied, and a posterior splint to keep the knee at rest.

Next day, the temperature and pulse were normal, and the patient was quite comfortable during the whole time of treatment. The wound was dressed on the 28th. There was no inflammatory blush. It was dressed again on March 4th and 8th, when the wound was united except where the drainage-tube, which was removed last time, had been. It was subsequently dressed on the 14th, 21st, and April 2nd, when it was quite healed.

CASE II.—A woman, aged 23, suffered from mammary abscess. It was opened on November 13th, 1878, and about three ounces of pus evacuated. It was dressed on the 15th. There was no inflammatory blush. The cavity was filled with blood-clot, but the drainage-tube prevented any tension. Only serum could be pressed out of the wound; no pus. The breast was full, with diffuse hardness, though there was not the slightest pain.

The wound was dressed on the 18th and on the 22nd. The drainage-tube was now removed. The general swelling of the breast had gone, and the hardness now existed as a lump in the gland-substance. The sinus was very slow in healing. The breast was dressed on the 29th, December 6th, 11th, 20th, and 26th, on which last date it was quite healed. There remained a very hard lump running from the cicatrix to the gland-substance, which, but for the history, might have aroused a suspicion of scirrhus. After the first dressing, the patient went about her usual employment, and never found any pain or inconvenience from the breast. I never saw her except on the dates above mentioned. This case is rather a contrast to the ordinary course of mammary abscesses; and I think, in it and similar cases, antiseptics, though complicated with a spray and what some call tedious details, amply repay the surgeon by the fewness of the dressings.

I treated several other mammary abscesses, with similar results—absence of constitutional or local disturbance. I think that this, among the poorest and lowest class of society, shows how easy it is to carry out antiseptics in private practice.

CASE III.—Another case, which tests most conclusively the efficacy of the antiseptic treatment, was one of empyema of seven weeks' duration, which came under my care whilst acting temporarily at one of the children's hospitals.

A little girl, aged 4, having all the symptoms of empyema on the right side, was admitted on January 12th, 1880. The dullness