are to be distinguished at a great distance; he must for that purpose possess a good visual acuteness, and be neither myopic nor hypermetropic, since an imperfect one or an ametropic defect may prevent him from distinguishing with the naked eye (and the use of correcting glasses must be interdicted in the case of a certain class of workmen, such as engine-men and stokers) the colours of the signals, except at distances less than those prescribed as necessary.

This truth has been admitted by the Medical Congress at Amsterdam, in the Ophthalmological Department, where the first article of their regulations is thus worded: "1. The experiments made on this subject on the lines of the railroad, prove that, to distinguish signals at the required distance, it is necessary to possess a sight perfectly normal."

The concomitant examination of the refraction and of the acuteness of sight is, therefore, indispensable; and the commission recently appointed by the Belgian Government, called "Committee of Daltonism", has not hesitated in considering it as the necessary complement to that of chromatopsical acuteness. This double experiment can easily be made by means of the Loiseau optometer, which has been already mentioned.

There are two classes of subjects to be thus examined: the one including men actually in the service, and desirous of being maintained; the other class is that of subjects wishing to enter the service. Both are equally interested in concealing their imperfections, and, in certain cases, pretend to qualities they do not possess. The Loiseau optometer (special model), modified according to the questions that are to be asked, and the conditions required in different administrations, will in certain cases be found of the greatest utility.

I did not intend to enter now on the question of Daltonism. It may, however, be useful to say a few words of the centric scotoma, which affects the eye for the most part when under the influence of intoxication, whether alcoholic or any other. The relative centric scotoma, so named because it surrounds the point of fixation (central), and because it often happens that the visual sensation is only altered in its relative chromatic properties (Nuel) with reference to the direct sight, may not be sufficiently proved by the wool-test; these may, however, be seen by the peripheral parts of the retina, which may still be healthy, in passing in front of them before exciting the direct point of vision. When subjects so affected look at a distance straight before them, it is the central point of the retina which must perceive the observed signals, and that point is exactly the one whose chromatopsic unfitness has been shown. Thence arises the danger of confusion in the interpretation of signals. To avoid this cause of error, we have adapted to the Loiseau optometer a system of discs in coloured glass. These can be slid in front of the apparatus in the place of the sign-board, which must be previously removed. Instead of the ocular lens, there is placed a thick opaque diaphragm perforated by a stenopaic aperture. The subject should be able to distinguish the colours of the disc in looking through that opening of the size of a pin's head, thus admitting only the intervention of the central vision.

BENZOATE OF SODA IN THE TREATMENT OF ACUTE RHEUMATISM.

By DAVID MACEWAN, M.D.,

Surgeon to the Dundee Royal Infirmary.

BENZOLIC acid is closely related to salicylic acid in chemical composition, and its physiological effects are in some respects similar; it is, therefore, a reasonable question for investigation, whether, like the latter, it possesses antirheumatic virtues. With the view of determining this point, I commenced to prescribe it nearly a year ago, in the form of benzoate of soda, for the treatment of acute rheumatism. On the first occasion on which I employed it, the relief of pain and subsidence of fever were so immediate, and the recovery was so rapid and complete, that I had no hesitation in adopting the same treatment in subsequent cases. The results have been favourable, as the following brief notes of some cases will show. The temperatures given were taken either about midday or early in the afternoon.

CASE I.—Jane S., aged 19. I saw her on May 19th, 1879, and was told that she had been suffering from slight pains in the joints for four or five days, but was able to go about till the 18th, when she had a rigor, and had to take to bed. The joints affected were the right shoulder, wrists, knees, and ankles. Pulse 120; temperature 102°. The tongue was coated. She was perspiring freely. The cardiac sounds were normal. She was ordered to have fifteen grains of benzoate of soda every three hours.

May 20th. She had passed an uneasy night. The pain in the shoulder

was gone; the right elbow had become affected. [Pulse 108; temperature 100.8°.

May 21st. She felt greatly relieved. Both wrists and one knee were still slightly affected; the other joints were free from pain, but somewhat stiff. Pulse 96; temperature 99°.

May 22nd. She had no pain except in one knee. Pulse 74; temperature 99°.

May 23rd. She was completely free from pain. The pulse and temperature were normal. She made a good recovery.

CASE II.—Jane T., aged 21. I saw her for the first time on December 10th, 1879. Four years ago, she had an attack of rheumatic fever, which continued three months. She had had shifting pains in her joints for the past five days. She became worse yesterday, and had to keep her bed. The joints affected were the left shoulder, wrists, elbows, ankles, the left knee, and some of the metacarpal joints of the left hand. Pulse 104; temperature 101.2°. There was a loud systolic bruit, clearly due to chronic mitral disease. She was ordered fifteen grains of benzoate of soda every three hours.

December 11th. The condition of the joints was much the same. Pulse 96; temperature 100.6°. She was perspiring very freely.

December 12th. The pains were gone from the wrists and elbows; the other joints were still swollen and very painful. Pulse 96; temperature 103°. She did not complain of præcordial pain or uneasiness. She was ordered to have twenty grains of the medicine every three hours.

December 13th. The joints were nearly free from pain. Pulse 78; temperature 99°.

December 14th. There was no pain. The pulse and temperature were normal. Recovery was uninterrupted.

CASE III.—Wm. H. became ill about the beginning of January 1880, and was treated for a fortnight by salicylate of soda, but without much benefit. On January 14th, the elbows, shoulders, and knees were very painful and swollen. The tongue was coated with a white fur. He had profuse perspiration. Pulse 102; temperature 102.8°. The cardiac sounds were normal. He was ordered twenty grains of benzoate of soda every two hours. For three days the temperature ranged from 101° to 102°, and the joints continued more or less affected; the symptoms then gradually subsided; and in about a week from the commencement of the treatment he felt quite well, and was able to be out of bed.

CASE IV.—Ann H., aged 27, had been ill for five days when first seen on January 24th. The wrists, knees, and ankles were painful, tender, and swollen; pulse 108; temperature 101.2°. The heart sounds were normal. She was ordered twenty grains of benzoate of soda every three hours.

January 25th. Pulse 108; temperature 101°. The state of the joints was not much altered.

January 26th. The joints were not so painful; the pain in the ankles was quite gone; pulse 96; temperature 99.2°.

January 27th. Pulse 84; temperature 100°. The painful state of the knees was gone; the wrists were still affected.

January 28th. Pulse 88; temperature 100°. She still had pain in the wrists.

January 29th. The joints were free from pain. Pulse 88; temperature 99°.

January 30th. She was quite well.

CASE V.—Mrs. F., aged 50, was laid up with rheumatic fever for three months, thirteen years ago. The present attack began on February 13th. When seen for the first time on February 16th, she complained of pains in both hips, in one knee, and both ankles; the slightest movement caused great pain. Pulse 104; temperature 101.8°. There was mitral disease of old standing. She was ordered twenty grains of benzoate of soda every two hours.

February 17th. Pulse 96; temperature 100°. The pains were greatly relieved.

February 18th. Pulse 96; temperature 99°. She had no pain except in the left knee, which was still slightly swollen.

February 19th. She was quite well. The medicine was stopped.

She continued well till February 21st, when she was seized with pain and swelling in the left wrist; pulse 100; temperature 100°. The medicine was resumed in the former doses, and next day the painful state of the joint had subsided, and her pulse and temperature were normal. She made a good recovery.

In addition to the above, I have treated several other cases, of which I have no notes, in the same way, and with equally good results.

The cases here narrated, are, of course, too few in number to warrant any confident conclusions; but they are sufficient to show that there is a strong probability of benzoate of soda being a remedy of considerable efficacy in the treatment of rheumatic fever. In all of them, the symptoms passed off in periods varying from three

days to a week after the commencement of the medicine; in none did cardiac complications arise during the course of the treatment; and, I think, the convalescence was more rapid than in cases I have seen treated by the salicylate of soda.

In Case v, there was a slight relapse, probably owing to the medicine having been too early discontinued; for, on its being resumed, the pain and fever speedily subsided.

Benzoate of soda possesses this advantage, that it does not give rise to the nausea and depression, or the unpleasant head phenomena, which the salicylate frequently produces. It is most convenient to prescribe it in the form of a mixture, and it may be given in doses of fifteen or twenty grains every two or three hours. It should also be continued in diminished doses for twenty-four or forty-eight hours after the rheumatic symptoms have disappeared.

DELIRIUM FOLLOWING THE TREATMENT OF ACUTE RHEUMATISM BY SALICYLIC ACID.

By T. D. ACLAND, M.B.Oxon.

In the BRITISH MEDICAL JOURNAL for January 29th, there are reports of six cases of acute rheumatism treated with salicylic acid and accompanied by delirium. In his remarks on one of these cases, Dr. Bastian put before his hearers the most recent views which have been expressed as to the causation of the delirium; and gave, as his opinion, that Dr. Murchison was probably wrong in attributing the symptoms to uræmia. In the year 1877, through the kindness of Drs. Bristowe and Ord, the opportunity was given me of making some observations on this point, which incline me to believe that, when the delirium is the result of the treatment, and not a natural sequence of the disease, uræmia may play an active part in its causation.

The method in which the inquiry was conducted was as follows: The urine of patients who were being treated with salicylate of soda was collected for each successive period of twenty-four hours. The percentage of urea was estimated from a sample of the whole (the mean of three observations being, as a rule, taken); and then, by a simple calculation, the total amount of urea excreted in the day was determined. The result of these inquiries made in four cases—one of which lasted for twenty-one days—was to lead me to the following conclusions.

1. As a rule, the treatment was followed by a decline in the temperature.

2. The decline in the temperature was generally coincident with or followed by a marked diminution in the excretion of nitrogen, which, except in very minute quantities, is wholly excreted as urea (Dr. S. H. West, *Proc. Roy. Soc.*, 1880).

3. The lowering of the temperature, diminution of pain, and lessened excretion of urea, were nearly coincident.

In two of the cases, these results were well marked; in the one, treatment was discontinued twice, with the result each time of increased pain, pyrexia, and total amount of urea excreted *per diem*; in the other, in which it was unfortunately impossible to obtain accurate observations, treatment was discontinued five times, with the result, on each occasion, of similar changes in the temperature and urea excretion. In both cases, as soon as the salicylic treatment was resumed, the pain was relieved, the temperature fell, and the amount of urea passed in the twenty-four hours was much lessened. Do the facts justify a supposition that, supposing the temperature had not fallen and the salicylic acid had still been continued, delirium would have resulted? Further researches alone can decide; and the difficulty of obtaining all the urine which is passed when a patient is unconscious is so great that as yet my efforts to do so have failed.

Five cases in which albuminuria has been present have come under my notice. In all of them it has passed away during the continuance of the salicylic treatment; though, in one case, there was a very considerable amount of albumen present when treatment was commenced. Should these statements be borne out by further inquiries, they may have an important bearing on the vexed question of the causation of delirium by salicylic acid; since, if they are true, there may possibly be uræmia due to lessened discharge, or perhaps lessened formation of soluble nitrogenous excreta, without the occurrence of nephritis; and it would seem probable that, in most cases, the albuminuria is only a part of the febrile process, and not due to the drug.

The results which have been observed in the treatment of enteric fever with salicylic acid seem to favour the above views; since the drug, in many recorded cases, has precipitated the appearance of the "typhoid" symptoms, which are generally ascribed to uræmia.

From the nature of the investigation, it has been difficult to obtain complete records of any large number of cases. Some accident has,

in many of them, broken the chain of observations, and made them incomplete; but those which have been recorded (*St. Thomas's Hospital Reports*, 1878), may, I venture to hope, lead others interested in the subject to continue the inquiry.

ACUTE PULMONARY TUBERCULOSIS.

By J. P. H. BOILEAU, M.D.,

Assistant Professor of Pathology, Army Medical School, Netley.

THE rarity with which this disease is followed by recovery is very clearly evidenced by the remarks of Dr. McCall Anderson, in the BRITISH MEDICAL JOURNAL of February 19th. I have treated two cases of this disease. In the first, all doubts as to diagnosis were cleared up by a *post mortem* examination; in the second, recovery took place, and it is now more than nine years since my patient was attacked. The following extract is from my case-book.

CASE I.—A. B., aged 29 (service eleven years), consulted me at Barbadoes on July 20th, 1871. He was suffering from pulmonary distress, cough, and hæmoptysis. He was placed on the sick-list; and I found that, although the cough had not been severe until very lately, he had been annoyed by it for several weeks past, and that it became aggravated after a hard ride into the country, which he took on the 14th instant.

There appears to have been an hereditary tendency to phthisis in his family. In the autumn of 1865, at Malta, I attended him for hæmoptysis, of which he had several attacks. He was not, however, then on the sick-list; and, during the six years that had since elapsed, he had enjoyed excellent health.

On examination, I found pulse (lying) 88; respiration 24; temperature (axillary) 101. He complained of wandering pain and heaviness about the front of the chest, headache, dyspnoea, and constipation. I found the respiratory sounds very feeble throughout the lungs, a general slight dulness, but no positive stethoscopic sign of pleurisy, pneumonia, or bronchitis. In the evening, his temperature had risen to 103°; pulse 100; respirations 25.

Subsequent examinations gave the following results.

Morning Observations.				Evening Observations.			
	Temp.	Pulse (lying).	Resp.		Temp.	Pulse (lying).	Resp.
July 23	101	90	28	July 23	100.8	88	28
" 24	100	84	28	" 24	100.2	100	30
" 25	99.5	84	28	" 25	101	93	31
" 26	101	88	28	" 26	100.5	96	30
" 27	100	88	28	" 27	100	93	30
" 28	100	84	28	" 28	100.8	84	28
" 29	100.6	84	28	" 29	—	—	—
" 30	100.1	84	30	" 30	101	—	—
" 31	100	96	30				
Average from 22nd inclusive..	100.3	87	28	Average from 22nd inclusive..	100.9	93	29

The sputa were scanty, and viscid, yellowish, and opalescent; latterly presenting milky-white spots and bloody streaks; no rust or frost. The cough was, therefore, more or less dry; it was often paroxysmal, and easily excited. The skin, which was delicate, emitted a peculiar odour; it was always moist. He became subject to flushes of heat, which could be seen in his face. I noticed an obscure crepitus at the base of the right lung; this gradually increased in loudness and extent. Three days after my first examination, I found dry crepitation, anteriorly and superiorly in the same lung; this became exceedingly well marked towards the apex, with dulness; vocal resonance and fremitus being morbidly increased. He lost flesh daily, and gradually became weaker. There came to be a difference of 30 between his standing and his lying pulse. Having made up my mind that he was the subject of acute tuberculosis, I submitted his case to the principal medical officer of the station, and recommended that he should be sent away from the island as speedily as possible. He embarked for England on August 10th, in a very unpromising state; but he gradually got better and stronger, he resumed his military duties, and he has done many a hard day's work since. He is now again serving abroad. I believe that the sea-voyage home, taken during the onset of the disease, saved his life; and I felt at the time that every hour he continued in the island was militating dreadfully against his recovery, so we got him out of the place with all possible speed.

WE understand that the Director-General of the Army Medical Department will not vacate his appointment on the expiration of his term of office in April next, and that he will continue to occupy his present position for another year.