which differed slightly from those described by Owen and Reinhardt, and which Professor Flower named *Pseudorca meridionalis*; but, since more skulls of the same species have been obtained, it is evident that they all belong to the one species, *Pseudorca crassidens*. A smaller, but closely allied form, is the *Orcella*, found in some of the rivers of India and the Bay of Bengal.

The *Beluga*, or white whale, is a purely arctic species, distinguished by being of a whitish tallow colour, and having no dorsal fin. The cervical vertebrae are not ankylosed together; the head is narrow; and the rostrum is small. The teeth are few, and of small size. A specimen of this whale was caught about two years ago, close to Dunrobin Castle, in Sutherlandshire, and presented by the Duke of Sutherland to the museum. It has a curious dislocation of the cervical vertebrae, which has probably been the result of an accident. The atlas has been dislocated laterally from the condyles of the skull, leaving only a small passage for the spinal cord. The animal lived for some time after the accident, as the vertebra has become ankylosed into its new position.

THE DIRECT METHOD OF ARTIFICIAL RESPIRATION FOR THE TREATMENT OF THE DROWNED, STILL-BIRTH, ETC.

By BENJAMIN HOWARD, A.M., M.D., F.R.C.S. ED.

I.

THE first sentence in the four-page Shortest Instructions of the Royal Humane Society for the treatment of the drowned is, "Send immediately for medical assistance."

The history of all the Society's receiving houses in London, however, even with their unparalleled facilities in this respect, furnishes not a solitary instance in which such assistance has been obtained in time to be of use. This, added to a nearly similar experience in the Island City of New York, and, so far as I have been able to ascertain, of other capitals also, painfully, but most emphatically, proves that only so far as non-medical men shall be made competent for the task, can the resuscitation of even a small proportion of the rescued be hoped for.

This conclusion led me, some years ago, to add to every set of rescue-apparatus within many miles of New York City, the printed instructions of the Royal Humane Society, and the National Lifeboat Institution, and to have them specially explained by familiar lectures to the harbour police. Another conclusion, much more disheartening, and which, to my astonishment, was forced upon me by various examinations of these men, many months afterwards, was that, whatever the value of the methods in question in medical hands, to these men, and to others like them, they were simply confusing and practically worthless. A third conclusion, the responsibility of which I could not evade, was personally to make a thorough investigation of the whole question. The outcome of this, arrived at step by step, year by year, was what is now known as the Direct Method of Artificial Respiration, embodied in a small half-page of simple instructions, reading as follows.

HOW TO RESTORE THE DROWNED.

Fig. 1.—Pressing and Draining Water from Lungs and Stomach.

"1. Instantly turn patient downwards, with a large firm roll of clothing under stomach and chest. Place one of his arms under his forehead, so as

* Being a report, with additions, of a lecture delivered before the Annual Meeting of the British Medical Association in Manchester.

to keep his mouth off the ground. Press with all your weight two or three times, for four or five seconds each time, upon patient's back, so that the water is pressed out of lungs and stomach, and drains freely out of mouth. Then :

Fig. 2.—The Bellows-Blowing Movement for pressing Foul Air out, and drawing pure Air into the Lungs.

"2. Quickly turn patient, face upwards, with roll of clothing under back, just below shoulder blades, and make the head hang back as low as possible. Place patient's hands above his head. Kneel with patient's hips between your knees, and fix your elbows firmly against your hips. Now—grasping lower part of patient's naked chest—squeeze his two sides together, pressing *gradually* forward with all your weight, for about three seconds, until your mouth is nearly over mouth of patient; then, with a push, *suddenly* jerk yourself back. Rest about three seconds;—then, begin again, repeating these bellows-blowing movements with perfect regularity so that foul air may be pressed out, and pure air be drawn into lungs, about eight or ten times a minute, for, at least, one hour or until patient breathes naturally.

"Note.—The above directions must be used on the spot, the first instant the patient is taken from the water. A moment's delay—and success may be hopeless. Prevent crowding around patient; plenty of fresh air is important. Be careful not to interrupt the first short natural breaths. If they be long apart, carefully continue between them the bellows-blowing movements as before. After breathing is regular, let patient be rubbed dry:—wrap in warm blankets—take hot spirits and water in small occasional doses, and then be left to rest and sleep."

While in England, as in America, the average layman usually at a glance seizes the principle, and on a careful reading at once becomes competent sufficiently well to practise the direct method, professional experts, on a closer examination, will find in it some important facts unknown before, assuring in the treatment of the drowned, and of apnoea from any cause, an effectiveness otherwise impossible.

As I proceed, I think it will appear, though perhaps not in the order I may now indicate, how in contrast to the instructions of the Royal Humane Society, and also of the National Lifeboat Institution—

1. The direct method alone provides for a thorough preliminary ejection of fluids from the stomach and thorax.

2. This method alone makes the drainage of the pharynx constant, precluding failure from lodgments there, or suction thence into the trachea.

3. In this method, the usually impracticable task of opening the mouth and pulling forward the tongue is more than superseded by simple position.

4. This same position also secures, and in the only way possible, the instant elevation of the epiglottis.

5. The compression by this method is the most complete, and capable of the most delicate adaptations.

6. It is the only method which can be practised by one person, and which by the same person can easily be continued as long as it may be of use.

Having myself corroborated, both by experiments and in practice, the conclusions of the first Resuscitation Committee of the Royal Medical and Chirurgical Society respecting the swallowing and inspiration of water, either the one or the other, or both, according to certain conditions of the submersion, the direct method, in contrast to the method curiously enough recommended by that same committee, begins by emptying the water-logged thorax, relieving the distended stomach, releasing the otherwise immovable diaphragm, and thus making an inspiration possible.

Having further corroborated the results of the experiments by the same committee respecting the "enormous force", as they state it, with which fluids, and even solid bodies, may, by the suction of an inspiration, be drawn into the trachea, in contrast to the position most favourable to these accidents, so strangely recommended by this same committee, the direct method maintains drainage of the entire pharynx from the beginning to the end of the operation.

In the position for first ejection the roll of clothing, by its size, makes the thorax and stomach the highest points, the nostrils and mouth the lowest; the latter being kept free of the ground by a wrist or forearm of the patient beneath his forehead. The displaced fluids rushing forward and downward cleanse the upper air-passages, which by position are then completely drained.

On the release of this first compression for ejection, the rebound may, at the very outset, secure an inspiration—the one aim of the whole process.

The facts in this method, to which, without reserve, I referred as new and important, and which relate to the head-down position of the supine patient, may be found in detail in No. 6, vol. viii, of the *Proceedings of the Royal Medical and Chirurgical Society*. They are briefly as follows.

In the position of utmost extension of the head and neck, as I call it, and which simply by the elevation of the thorax, as directed in the supine position, is made moderately possible, the line of gravitation of the tongue has its terminal point shifted from the posterior wall of the pharynx to the hard palate, at or about its junction with the soft palate. The entire posterior wall of the pharynx is shifted backward; its anterior wall is shifted forward; thus, its antero-posterior diameter, as much as is possible, is throughout increased; while, by the shifting upward and backward of the nares, their entrance is brought more directly over and in a line with the course of the pharynx. The larynx being pulled downward and forward by the sterno-thyroidei muscles, and fixed there, the extensive motion upward and backward of the lower jaw puts upon the stretch the genio-hyoidei, mylo-hyoidei, and anterior bellies of the digastric muscles, causing the hyoid bone, and, by means of the hyo-epiglottic ligament, the epiglottis, to share together the motion of the jaw. Thus, the epiglottis is instantly made vertical. The thyroid insertion of the palato-pharyngei muscles being brought downward and forward by the sterno-thyroidei and fixed, the palato-pharyngei muscles are put upon the stretch their whole length by the extensive movement upward and backward of the head; and thus the posterior pillars of the fauces, the arches of the palate, and the velum palati, into which latter membrane these muscles are inserted, are all pulled downward and forward; they are thus made tense and kept so. The velum being thus stretched some distance in front of the back of the pharynx, a postoral air-way is secured, from the nares to the glottis, from which the tongue is doubly excluded.

Hitherto the propositions for obtaining an air-way, without the existence of which nothing except failure can be assured, have, even in the best selected circumstances, been practicable only in part. In No. 21, vol. i, of the *Lancet*, I have stated the fact, and shown the reasons why it can be no otherwise, that for the elevation of the epiglottis, the universal practice of pulling forward the tongue, however skillfully it be done, avails nothing, in spite of the fact that, even with the aid of jaw-prisers and gags, and tongue-forceps and assistants, and all the appliances of the operating theatre, the opening of the clenched jaws and the seizing of the tongue, which is deemed but a preliminary to the raising of the epiglottis, is oftentimes, as any practised chloroformist will admit, one of the most difficult tasks he is called to perform.

In the face of this experience of experts, the following instructions to the accidental bystander by the Royal Humane Society, if the matter were a less serious one, would be simply comical.

In these instructions, after directions to send for medical assistance, the solitary and unlearned comrade of the drowned man at the water's edge is told to "open the mouth, draw forward the patient's tongue and keep it forward", the only suggestion as to how, in either case, he is to do it being, "apply an elastic band over the tongue and under the chin".

It may be superfluous to say that, after much inquiry, I have been unable to find anybody who had ever even heard of a drowning case in which by a layman these instructions had been carried out—and for the best of reasons.

With the jaws inseparably clenched, the slippery tongue receded into the pharynx, the epiglottis fallen back—only as either one or more of these conditions may be accidentally incomplete, can exchange of air, or recovery by any mere thoracic procedure whatever, be possible. Just here appears conspicuously one of the merits of the direct method. The bystander has simply to place the patient in the position as directed, and instantly the free postoral air-way is substituted for an impassable oral way in the manner described, and respiration is at once possible.

When, as sometimes occurs, syncope is a principal or incidental feature in the case, this position is, of course, for that also, the best and quickest remedy. Mucus, too, and other accumulations which otherwise might flow or be sucked into the trachea, are thus during respiration, as I have before remarked, constantly drained away.

Having now an open and widened air-way, we come to the movements for respiration, in which manual compression of the thorax and displacement alternate with inevitable rebound, corresponding inrush of air, and replacement.

The precise manner in compression which I prefer is to place the thumbs along the anterior free borders of the costal cartilages, so that the fingers naturally fall into the lower intercostal spaces. The force then applied to the ribs, is, through the medium of the thoracic parietes, so distributed, as, without shock or violence, to bend the ribs upon themselves in the line of their own axes, the operator feeling his way as to how little or much he is doing, or may do, and regulating his course accordingly. More particularly, he is thus able on the first return of short and imperfect gasps to observe, aid, and deepen these, and, in various ways impossible by the Humane Society's or other methods, to adapt his movements to the varying need of the moment.

The direct method is as easy to do as it is to understand. The kneeling position, even if not habitual with the operator, is not a tiring one. The part to be pressed is raised to his hands, and his weight is the chief force employed, while there are frequent intervals of complete rest. The rate of the respirations, which, it will be noticed, is one-third to one-half slower than in previous methods, gives, even then, nearly three times more respirations a minute than will, as can be easily proved, suffice without discomfort for an ordinary healthy man in repose. The eight-or-ten rate, while it supplies more oxygen than the asphyxiated or partially recovered patient can for some time assimilate, allows of better diffusion, minimises superfluous disturbance, and promotes ease in its continuance. On such an inconvenient spot as a boat, as also on a feather bed, etc., where the other methods are impracticable, I have myself, as have others, effectively continued this method entirely unaided for more than an hour.

In all these particulars also, this method is in complete contrast to the other methods. The illustrations of the Marshall Hall method include three or four operators; for the Sylvester method two at least are necessary.

For the semi-rotation of a perhaps heavy and virtually dead body, at the rate of seventy-five times every five minutes, according to the instructions for the Marshall Hall method, no one man would be strong enough; and no four accidental bystanders would be trained enough to do it in unison. Without discussing the otherwise possible merits or demerits of this method, it is, for the treatment of the drowned, simply as nearly as possible impracticable.

In performance, the Sylvester method is both difficult and very fatiguing. The holder of the tongue is directly in the way of the operator. The compression desired is attempted upon a part almost beyond the reach of the operator; is made transversely to the ribs and downwards, instead of in the line of their axes, and inwards and upwards; while between the hands of the operator and the part to be pressed are interposed the arms of the patient. The maximum of labour thus so ingeniously adapted to a minimum of effect, is such that a few minutes' faithful performance of it quite exhausts the strength of even a skilful and duly assisted operator. The first and fatal check which, as I said, I myself received in the use of this method in connection with the Metropolitan Police of New York, however, and arising, I think, in part from the obscurity of relation between the motions as directed, and what non-medical persons know about the process of breathing, was owing, I afterwards found, to no local causes. Extending my personal investigations to almost every receiving house in Europe, I found that, with the exception of those on the Serpentine in London, even the *employé* experts, followed the instructions "draw the arms *gently* and steadily upwards", and "then turn down the patient's arms and press them *gently* for two seconds against the side of the chest" in such a *very gentle* way, that uniformly the whole process was simply a moving of the arms up and down *gently* in their sockets. This so completely missed the principle and the objective points of the method, that even the author of it would admit such performance to be a parody, which could promise no certain benefit.

In the amphitheatre, with operating surgeons, the whole question is a different one, the performance in that case usually being much better than the "instructions". Even here, however, the superior advantages of the direct method are scarcely less apparent.

With the head as low as is convenient from the beginning, should the anaesthetised patient give the slightest suspicion of apnoea, the instant head-down position, united with firm extension by pressing the chin upward and the forehead backward, corrects the condition in the simplest, quickest, and most intelligible way, superseding the present instrumental violence, and without disturbing the surgical operation in

* The italics are my own.

progress. As a rule, all threatening symptoms will thus be caused to cease. Should artificial respiration be required, just as we are always content to do with animals in the laboratory, let thoracic compression be made in the manner I have described, and we have artificial respiration in its simplest form. If, however, the operator prefers the supposed and, perhaps, positive advantage obtainable from arm-extension, the adaptability of the direct method is such, that this can be secured also. The extension may be made thoroughly, and direct compression also, thoroughly in rhythmical alternation. Even tracheotomy, electricity, friction, etc., might be added if desired, and the direct method proceed without interference.

The instructions of the Royal Humane Society were, I am aware, recommended by a committee of men the most competent and distinguished, perhaps, in the profession; but not one of whom, I believe, had the advantage of an actual experience in the treatment of drowned men. Their recommendations were based upon the results of experiments made with the aid of practised experts upon the selected and drained cadaver.

In every case, before the experiments were begun, tracheotomy was performed, and a tube was carefully attached to the trachea.

Thus all the more perplexing and, I may say, till now unsurmounted obstacles connected with the upper air-passages were jumped before starting, and the experiments were proceeded with under conditions as different as they could be made to be, from those encountered by physicians; while they entirely miss the first facts in the problem of the actual treatment of the drowned—the best instructions for the treatment of the drowned for the use of the largest number of uneducated laymen.

In trying to be brief I fear I have been obscure, especially respecting the facts which, at the beginning, I promised should be found to be novel and important, connected with the upper air-passages. I may, therefore, be allowed to say that each of these facts above mentioned has been repeatedly demonstrated by me to the acknowledged satisfaction, as well as surprise, of such distinguished authorities at St. Thomas's and Guy's Hospitals, and at the Royal College of Surgeons, as I think you would yourselves be most likely to select.

Unlike, I believe, any previous authors or their co-labourers on this subject, all that I have said or done in connection with it has been the offspring of arduous, practical, and extensive experience—an experience, I may add, which has involved no little sacrifice.

If I have been clear, I think you have now been able to understand the reasons for the distinctive positions and movements of the direct method, and to see the several advantages these secure in connection with ejection and drainage—the postoral air-way and the more delicately regulated and sufficient respiration, which, at the beginning, I said I hoped to be able to show.

That which I claim as the most distinguishing feature of the direct method, however, is, that its effectiveness is united with such simplicity as quickly to enable the *average layman* to be competent for the best treatment of the drowned, and that it thus supplies the only condition by which, according to our past experience, a larger saving of human life may in these cases be hoped for.

We will next consider the direct method in the treatment of still-birth.

THE HEALTH-OFFICERS OF THE ISLE OF WIGHT.—Dr. Ballard has made to the Local Government Board a very elaborate and lengthy report, on the sanitary condition and administration of the Isle of Wight. He records much inconvenience as arising from the multiplicity of local authorities in the island, and from the unsatisfactory conditions under which many of the health-officers hold their appointments. He finds much matter for criticism in the way in which the sanitary authorities do their work, especially as regards keeping pace with the growing necessities of villages, which are rapidly becoming towns. He regards, as very disadvantageous, the arrangements by which in the rural district, the nine district medical officers have been appointed *ex officio*, medical officers of health, and in the urban districts private medical practitioners have been separately appointed to the post. He thinks it of great importance that the services of a skilled medical officer of health, not occupied in private practice, should be available for each district; and in order to provide an adequate salary, he recommends that a combination of all the sanitary authorities in the island, should be made for the appointment of a single officer. The area and population are certainly large, but not larger than those of many existing combined districts where the appointment of a single medical officer of health has been attended with excellent results; and there seems no reason to doubt that an officer living in a central position, and devoting his whole time to sanitary work, would be able to exercise proper supervision over the whole island. If this desirable combination could be accomplished, it would be a great and obvious improvement on the present arrangements, which are admittedly imperfect and unsatisfactory.

THERAPEUTIC MEMORANDA.

EUCALYPTOL *versus* PHENOL.

DURING the prevalence of the last epidemic of small-pox in London, I called attention, in the *BRITISH MEDICAL JOURNAL* (February and May, 1871), and in the *Lancet* (April, 1872), to the dangers attending the use of phenol and its preparations as disinfectants, and as surgical dressings; and I advocated the substitution of heat and sulphurous acid for disinfecting purposes. I have had the satisfaction of seeing these latter processes—previously known, but little employed—firmly established in public favour; and I now have the further satisfaction of seeing another of my *protégés* proposed by Mr. Lister as a safe and efficient substitute for carbolic acid in surgical practice. I have long been convinced of the beneficial action of the aromatic emanations (not the absorbent powers) of the eucalyptus tree; and have endeavoured to introduce some of the harder varieties into this country, by distributing a large quantity of seeds in the most suitable districts. The French have studied the chemical nature, and the therapeutical action, of the eucalyptus globulus more thoroughly than we have done, probably because it is an important product of their country. The substance which we call the oil of eucalyptus is, or was very recently, called eucalyptol by the French chemists. The term is a more convenient one than ours; and, as it corresponds with our other antiseptics—phenol, thymol, etc.—we may accept it, without inquiring too curiously as to its chemical accuracy. The substance, indeed, is only the lighter and more volatile portion of the oil of eucalyptus, which distils over at a temperature below 175° C. (347° Fahr.); a second substance passes over at a temperature of 188° to 190° C. (360° to 374° Fahr.); while a third remains in the retort. On account of the slight solubility in water and its great volatility, Mr. Lister recommends that the eucalyptol should be mixed with damar gum; but it will probably be found that eucalyptus gum—or rather resin—from which the tree derives its popular name in Australia, will be found more suitable. The French have long been employing a combination of the kind, under the name of the *Alcoolature d'Eucalyptus*, a common tincture of the fresh leaves, containing a large amount of the essential oil and resin, some tannin, and probably a little chlorophyll, as it is of a dark olive colour. I have myself used the preparation as a surgical dressing, and in some forms of skin-diseases, without ill constitutional effects. I have found, however, that doses of twenty to thirty drops, taken internally, produce irregular action of the heart, and a feeling of prostration similar to that produced by an overdose of tobacco; so that eucalyptol may not be quite so safe as Mr. Lister has found it to be in his practice so far. The French chemists also prepare an *Eau distillée d'Eucalyptus*, which may be suitable for antiseptic surgery as "*l'huile essentiel s'en sépare difficilement*", and it can be concentrated by redistillation.

C. ROBERTS, Bolton Row, Mayfair.

INTERNAL USE OF IODOFORM.

I SEE in the *JOURNAL* of May 28th a communication from Dr. Crocker on the use of iodoform in the treatment of skin-diseases. I gather from it that he alludes to its use as an external application. It may, therefore, be of interest to mention that I have lately administered iodoform internally with complete success in an obstinate case of molluscum, where a prolonged course of arsenic and a variety of other remedies failed. It was given in one-grain doses three times a day. I found that the most agreeable way of giving the medicine was in the form of sugar-coated oval pills, procured from Messrs. Burroughs of Snow Hill. W. J. EAMES, M.K.Q.C.P., Fleet-Surgeon R.N.,
H.M.S. *Royal Adelaide*, Devonport.

HAMAMELIS VIRGINICA AS A LOCAL APPLICATION.

HAVING during the past few months given a somewhat extended trial to the new extract of *Hamamelis Virginica*, called "*hazeline*", I beg to place the following memoranda at the disposal of the readers of the *JOURNAL*.

As a local application in irritable and inflamed piles situated at the margin of the anus, where the remedy can be readily applied, I have never met with its equal. In most of the cases submitted to the treatment, the relief was immediate and permanent. My plan has been to have the part bathed in the solution three or four times a day, and a piece of lint dipped in it kept applied to the anus during the intervals. All urgent symptoms have passed away, as a rule, in from twelve to twenty-four hours.

In chronic and intractable ulcers of the varicose or eczematous de-