Six months after he had commenced this mode of living, I met him, and asked him how his sight then was. "Nearly well", he replied; and, to my surprise, on examining the eyes, I found that the nystagmus had almost entirely disappeared, the right iris answering freely and the left slightly to the stimulus of light. The left pupil was also less dilated. I may summarise the after-history of the case by saying, that I have seen this man frequently since, and that the eyes have continued to improve, so that, at the present time, he can read small print fairly well. I will not weary you by relating any more cases of the kind, of which I have several, but will merely make the following remarks in conclusion.

1. *Pathology.*—I have examined the optic disc with the ophthalmoscope, but do not feel sufficiently satisfied in my own mind to record my opinion here. I will just remark that the history of the above case shows that there is probably no grave permanent lesion of the retina which would lead to there being visible anatomical changes, else the rapid improvement of vision described would not have taken place.

2. *Cause of the Disease.*—The miners themselves give the credit of it to the insufficient light cast by the safety-lamp. This may be the case; but, since we do not find the disease mentioned as occurring in other classes of miners who work in almost as dark places, I am inclined to attribute it to the inhalation of the poisonous gases which are found in the coal, which probably alter the composition of the blood; and thus not only interfere with the nutrition of the eye, but also produce a local venous congestion. This question is of importance; for, if the complaint be produced by insufficient light, it is the duty of Government to encourage the invention of a lamp as safe as, and more effective than, the one in use at present. If, on the other hand, the disease originate in the cause I have suggested, better ventilation seems to be urgently required.

3. *Treatment.*—Little or no treatment appears to be necessary. In all the cases I have observed, very marked improvement took place on exposing the patients to free light and air; in many, complete cure was effected; in all, a great improvement was perceived.

NOTES OF A CASE OF INTRA-UTERINE AMPUTATION.

By R. J. ANDERSON, M.D.,

Demonstrator of Anatomy, Queen's College, Belfast.

CASES of intra-uterine amputation, although by no means common, yet present features of interest sometimes that have more direct relation to the cause of the accident. The following example has been thought worth recording, as an opportunity was afforded of dissecting the parts, so that the condition of the muscles after a period of many years was examined.

A female subject, aged 55, was brought to the anatomical rooms early in the session 1880-81. The left upper extremity presented no variation from the natural condition as far as the middle of the arm; below this point, a constriction existed; and in the region of the stricture the skin was darkly pigmented. A large cutaneous fold extended from a point six inches above the elbow to a point four inches below. It was two or three inches in breadth at the elbow-joint, and consisted of two portions of skin placed back to back. The distal phalanx of the thumb, and all the metacarpal phalanx except the base, were wanting. The proximal phalanx of the index finger was all that remained of that digit. The flexor profundus digitorum in the index finger was inserted a little above the flexor sublimis, near the distal extremity of the first phalanx. The flexor longus pollicis was inserted into the base of the metacarpal phalanx. The middle, ring, and little fingers were entire; the latter, however, was bent, the second and third phalanges being placed at right angles to one another. The tendon of the extensor communis digitorum to the index finger was attached to the distal extremity of the first phalanx. The extensor secundi internodii pollicis was attached to the remains of the proximal phalanx of the thumb. Nails were not present on the first and second fingers.

In the right upper extremity, the second phalanx of the thumb and a portion of the first phalanx had been removed. The last phalanx, and the middle, except the base of the index finger, the third, and three-fourths of the middle phalanx of the middle finger, were absent. Nails were absent on these; a nail was present on the ring finger, at its extreme end its distal phalanx had disappeared. The little finger was perfect. The flexor longus pollicis was inserted at the distal extremity of the first phalanx. The tendons of the flexors were inserted together into the base of the second phalanx of the index finger, that of the superficial flexor splitting. The tendons of the same muscles to the middle finger were inserted into the base of the second,

that of the flexor profundus being placed between the two parts of the sublimis. The common extensor tendon was inserted into the base of the second phalanx, and the arrangement in the case of the third and fourth was similar. The extensor primi internodii pollicis was inserted into the base, the extensor secundi internodii into the distal extremity of the first phalanx. A slip from the tendon of the extensor primi internodii passed to that of the latter muscle, and was inserted with it.

The right lower extremity presented few features of interest. An almost complete web extended between the fourth and fifth toes; a less marked cutaneous fold was present between the second and third.

The left lower extremity presented an interesting case of talipes varus. The toes had suffered as in the case of the hands. The inner aspect of the gastrocnemius was composed largely of fibrous tissue, and was much smaller than the outer. The other parts did not differ materially from the condition usually found in similar cases.

In a case referred to in the *Medico-Chirurgical Review* (vol. xix), a constriction existed above the elbow. The parts were atrophied and oedematous. The radial side of the wrist was bent, and the fingers contracted. In the case above recorded, the parts beneath and in the neighbourhood of the constriction were not much affected, showing that the constriction, if it were ever present, must have passed away. No constrictions were present in the lower extremities, as in Frickhöffer's case, where, with a talipes varus, a constriction occurred at the left knee-joint.

In volume xxxii of the same journal, an abstract of G. Braun's views is published, with cases. The abnormal growth of the amnion produces definite bands, that are carried to the back of the embryo, and the bands, extending from both sides, coalesce. These bands produce a very marked effect on the development of the embryo. In one case given by the above-mentioned observer, there was a fissured palate. The frontal and occipital bones were undeveloped, and fibrous bands extended across the cranium. In another, the nose and upper lip were fissured.

ELECTRO-DIAGNOSIS.*

By J. DIXON MANN, M.D., M.R.C.P.

THE rapid progress recently made in the pathology of diseases affecting the nervous system, by demonstrating the relation between organic lesions and the resulting train of symptoms, has greatly enlarged the field of diagnosis, and has developed the use of systematic methods of investigating the physical conditions to which such diseases give rise. In seeking to ascertain the essential nature of a neurosis, many methods are used to test the functional activity of the affected muscles and nerves; and, by comparing the results with those obtained from an examination of corresponding and healthy structures, to localise the causal lesion, and to gauge its intensity. In investigations of this kind, much information is derived from comparative observations of the immediate effects produced by electricity on nerve and muscle, in health and in disease. The diagnostic powers of electricity are largely used by those specially engaged in the treatment of diseases of the nervous system; but, owing to an exaggerated estimation of the technical skill required for its practice, electro-diagnosis has not yet received the general attention due to its importance. The time approaches when the diagnosis of many diseases of the nervous system, hitherto considered obscure, will be a matter of ordinary routine; every practitioner, therefore, should be conversant with all the recognised methods of investigation.

The application of the constant, or voltaic current, to a healthy muscle, produces contraction only at the opening and at the closure of the circuit—that is, at the moment the current begins, and at the moment it ceases to flow; whilst the current is flowing, the muscle is passive. Different degrees of current-strength are required to evoke muscular contraction, in accordance with the fourfold manner in which the contraction may be produced—namely: at the closure of the circuit, and at its opening, with either the positive or negative pole. Beginning with a mild current, and gradually increasing it, the first normal reaction which occurs is that produced by making contact with the negative electrode—cathodal closure contraction (C. C. C.). Next comes the contraction produced by breaking contact with the positive electrode—anodal opening contraction (A. O. C.). Almost synchronous with the last is the anodal closure contraction (A. C. C.); and, last of all, requiring a most powerful current, is the cathodal opening contraction (C. O. C.). When the current is very strong, the cathodal closure contraction assumes a tetanic character; if the current be still further augmented, the anodal closure contraction evinces a like tendency. The

* Read at the annual meeting of the Lancashire and Cheshire Branch.

diagnostic value of these reactions is derived from certain changes produced in their amount, order, and character, by different forms of motor paralysis. For the purposes of electro-diagnosis, paralyzes may be divided into two primary divisions: 1. The electrical reactions are unaltered; 2. The electrical reactions deviate from the normal. The last division is subdivided into: (a) cases where electro-irritability undergoes simple increase; (b) cases where electro-irritability undergoes simple decrease; (c) cases where qualitative, as well as quantitative, changes occur.

1. Paralyzes in which *electro-irritability is unaltered*. Cerebral: from apoplexy, embolism, or tumours. Spinal: from chronic myelitis. Peripheral: rheumatic, slight facial, and slight traumatic paralysis.

2. (a) Where there is *simple increase of electro-irritability*. (This deviation is of rare occurrence.) In the early stage of hemiplegia, in locomotor ataxy, and in the early stage of some peripheral paralyzes, *ex. gr.*, facial paralysis.

(b) Where there is *simple diminution of electro-irritability* (more frequent than the last). In bulbar paralysis (rare); in spinal paralysis, accompanied with muscular atrophy; in progressive muscular atrophy.

(c) Where there are *qualitative, as well as quantitative, alterations in electro-irritability*, called by Erb "the reaction of degeneration". This deviation is characterised by quantitative changes in the electro-irritability of nerves, and in the farado-irritability of muscles; and by both quantitative and qualitative changes in the volto-irritability of muscles. The following is an epitome of the changes produced by an acute attack of average intensity. About the third day after the onset of the paralysis, there is diminution in the electro-irritability (faradic and voltaic) of the nerve; about the twelfth day, it is entirely lost. In cases of recovery, nerve electro-irritability gradually returns. The farado-irritability of muscles also diminishes, and eventually disappears. The volto-irritability of muscles falls during the first week; but, in the second week, it undergoes a remarkable increase and qualitative change. The feeblest current, slowly interrupted, produces a sluggish contraction, differing from the rapid movement of healthy muscle; a state of muscular tonus, more or less marked, persists as long as the current flows; rapid interruptions, even with a powerful current, fail to produce any muscular reaction. The anodal closure contraction equals, and occasionally exceeds, cathodal closure contraction; and cathodal opening contraction tends to equal, or surpass, anodal opening contraction. After a variable period, the opening contractions disappear, and the volto-muscular irritability diminishes (in cases of non-recovery) progressively to extinction. The inverted order of the reactions persists to the end; the last contraction is that caused by anodal closure. In cases of recovery, when a certain stage of diminished irritability is reached, regeneration sets in, and is followed by a rise of volto-irritability towards the normal. The improvement in muscular volto-irritability, and the restoration of the other normal faradic and voltaic reactions, do not take place until after the return of motility. These changes in electrical reaction are due to degeneration of the intramuscular nerve-fibres, and to certain histological changes undergone by the minute muscular structures. The stage of degeneration may be deduced from the degree of abnormal reaction present; when the reactions tend to resume their normal order and character, regeneration has commenced. The following paralyzes are attended by the reaction of degeneration: severe traumatic paralyzes; paralyzes produced by pressure on a nerve, sufficiently severe as seriously to interfere with the performance of its functions; acute polio-myelitis anterior (infantile paralysis); hæmatomyelia; medullary paralysis; progressive muscular atrophy; and saturnine paralysis.

For the practice of ordinary electro-diagnosis, the disk electrodes, supplied with medical batteries, suffice; much more delicate reactions, however, may be obtained by the use of small olivary electrodes furnished with current-interruptors, by which the circuit can be closed and opened without changing the position of the electrodes. A pedal interruptor is a convenient accessory for producing slow interruptions with the induction apparatus. The following conditions are to be observed. The patient should be placed in a well diffused light; the limbs under observation should be arranged symmetrically, in an unconstrained position; and the patient should allow the muscles to be perfectly passive. If the paralysis be unilateral, care should be taken to apply the current equally to the two sides, both as regards strength and localisation.

As an illustration of the method employed in the practice of electro-diagnosis in its simplest form, take a case of infantile paralysis of one limb—say, the arm—a few weeks after the subsidence of the acute stage. Place a flat disk electrode, connected with one battery terminal, on the cervical spine, and fix it there. To the other battery terminal, by means of a double or branched conducting cord, attach two small electrodes, with current-interruptors; and, having selected a muscle, place

the electrodes on corresponding points on the two arms. See that the electrode covers are well moistened with water, acidulated with a few drops of acetic acid; and, if the skin be very dry, pass over the arms and neck a sponge squeezed out of hot water. Let an induced current first be used. Complete the circuit with the interruptor on the sound arm, and gradually increase the current-strength until a well-marked contraction is produced; now, try the effect on the paralysed limb: there is no response. If the current-strength be increased to the highest bearable limit, and the interruptions be made to succeed each other very slowly, still there is no manifestation of faradic irritability. Change the induced for the constant current; commence with the paralysed limb; and, by means of the battery commutator, make the electrodes applied to the arms alternately positive and negative; add cell after cell, opening and closing the circuit, with the interruptor, several times after each addition to the current-strength, until a slight muscular contraction is seen; this is first produced, probably, by closing the circuit with the anode, though in many cases it is not easy to decide the priority of A. C. C. over C. C. C. If the same current-strength be now applied to the healthy limb, no reaction is perceived. As previously stated, the volto-irritability of the paralysed muscles is exalted at this stage; a much weaker current suffices to evoke contractions in them than is required in the case of healthy muscles. If the current-strength be increased until the opening contractions are produced in the paralysed limb, it will be found that C. O. C. is coetaneous with, if it do not precede, A. O. C. A medium current, producing, with slow interruptions, well-marked contractions of the affected muscles, with rapidly succeeding interruptions, fails to produce any muscular reaction. When the paralysis is paraplegic, there is obviously no opportunity of comparing the diseased muscles with the corresponding muscles of the opposite limb; in such cases, experience is the only guide. The operator must, therefore, draw upon his memory for the requisite data wherewith to make comparison.

In cases of partial motor paralysis, especially when restricted to isolated muscles, a more minute investigation is required. The localisation of the electrodes must absolutely correspond on the healthy and on the diseased limbs; more than one examination will probably be necessary for the formation of a reliable opinion.

CÆSAREAN SECTION.

By ARTHUR PERIGAL, M.D., Barnet.

A. S., aged 20, unmarried, fell in labour with her first child at 9 A.M. on February 9th, 1879, at the union hospital. I saw her at 11 A.M., and found pains at long intervals, and weak; the head was very high up, the pelvis much contracted antero-posteriorly, and the sacral prominence markedly developed. Her general state was comfortable. At 10 P.M., I was called to her; the pains were then strong, but not frequent; the head had descended slightly; the cranial bones felt very hard. There was no bleeding. As no advance was made during the next hour, I applied the long forceps, thinking thus to overcome the obstacle; they were adjusted with difficulty, traction was used for an hour and a half, but no advance made. I now introduced my hand to endeavour to turn (which I should have done at first had the membranes not been so long broken), but the sacral prominence so jammed my hand, that I could not get my wrist past it; and, although I got hold of a foot, I was obliged to desist. I then asked Dr. Livingston (whose case originally it was) to help me. He, too, tried the forceps, then turning, but failed. We now evacuated the head, and applied the forceps and hook, but to no avail. The patient being a good deal exhausted, we gave a full opiate, and left her at 6 A.M. in charge of a qualified assistant.

At 11 A.M., we met in consultation with Mr. Carter, who suggested a final trial. The parts were hot and painful, the discharge offensive. She had slept; there was no delirium; pulse 120, weak. She had taken plenty of beef-tea and milk. After a fair trial, all hope of delivering *per vias naturales* was abandoned, and we decided on hysterotomy as giving the only, but slight, chance of saving her life.

The patient having been put under chloroform, and the bladder emptied, Dr. Livingston sponged the abdomen with carbolic acid water, and made an incision from the umbilicus of five inches and a half, cutting through the abdominal walls, and dividing the peritoneum on a director. The mesentery being gently pushed aside, and some dark serous fluid sponged away, the uterus was divided for about four inches and a half, and the child exposed. The child, which was very large, was easily removed; the placenta, though slightly adherent to the left wall, coming with it. I may add that its position was noted before the operation. The uterus at once contracted, no blood escaping into the abdominal cavity. Some fluid was sponged therefrom