other cases besides some of my own with success. There is also an admirable instrument on this plan invented by Dr. Meadows, which is deservedly in great request for this difficult class of cases; but I fear I must not trespass further on your time. If I have failed to establish the two propositions I set out with, I shall be pleased to think if I have succeeded in removing some of the unfounded prejudice with which the mechanical treatment of uterine displacements and flexions has been, to my mind, so unreasonably regarded.

If those who so strenuously oppose this therapeutic system had any preferable alternative to offer in its stead, there might be some less difficulty in understanding their position; but it is not so. A specimen of the treatment they recommend is embodied in the following quotation, with which I shall conclude. "Time is an important factor in uterine therapeutics. Patients" (suffering from uterine flexions, etc.) "should be kept for a length of time under observation, and should be seen and examined at intervals"! What benefit is to be derived from such a process of expectancy in the class of ailments which we have been considering, it is not easy to discover.

REMARKS

ON

THE RELATION OF DISEASES OF THE NASAL-PASSAGES AND NASO-PHARYNX TO AURAL AFFECTIONS.*

By THOMAS BARR, M.D.,

Aural Surgeon to the Western Infirmary, Glasgow.

THERE can be little doubt that an essential advance has been made, of late years, in the knowledge and treatment of those morbid states of the nasal and pharyngeal mucous membranes which give rise to disease of the ear. We have amongst us to-day two eminent *confirères*, who have contributed not a little to this advance in our knowledge. I refer to my friends, Dr. Guye of Amsterdam, and Dr. Loewenberg of Paris, whose work, in connection with the important subject of adenoid vegetations, has been of the greatest value.

Aural specialists have been of late more and more convinced of the great signification of nasal and pharyngeal diseases in their relation to affections of the ear. We recognise, as one of the well-established points of aural pathology, that this region is not only frequently the place of origin of ear-diseases, but that its morbid conditions have a most important influence on the course and issue of affections of the middle ear; and, therefore, in our modes of examination and in our therapeutic measures, attention to the naso-pharyngeal cavity is looked upon as of the very first importance. Even in cases where no further improvement is to be hoped for in the hearing power, we may often, through treatment of the nasal and pharyngeal mucous membrane, prevent the further advance of the evil.

The apparatus of the middle ear is influenced by very varied morbid conditions of the nasal and pharyngeal spaces. These conditions exert their influence upon the middle ear through the medium of the well-known passage of communication between the upper pharyngeal cavity and the tympanum—the Eustachian tube. By this channel the mucous surface of the middle ear, and that of the pharynx, are directly continuous; while, by the same channel, under certain conditions, free interchange of air takes place between the two cavities. Thus, by these two media of communication, mucous membrane and the air—pathological conditions of the naso-pharyngeal cavities—impress themselves upon the structural and functional integrity of the middle ear.

We know that affections of mucous membranes are very apt to propagate themselves by simple continuity. Hence, a large number of diseases of the tympanum and mastoid cells are caused by simple propagation, along the mucous surface of the Eustachian tube, of catarrhal or inflammatory processes, primarily affecting the lining of the pharyngeal cavity. And such diseases of the tympanum and mastoid cells, with their results, may exist even years after the primary cause in the pharyngeal mucous membrane has passed away. No one now disputes that congestion, swollen and hypertrophied conditions of the naso-pharyngeal mucous membrane, very frequently maintain a state of simple or purulent catarrh of the middle ear.

But important changes in the structure and function of the tympanic apparatus may arise from pathological conditions of the nasal passages and pharyngeal cavity, without any extension of inflammatory processes

along the mucous surface of the Eustachian tube to the tympanum. Here the mischief in the ear is brought about through mechanical interference with the lumen of the pharyngeal mouth of the tube, by the encroachment upon it of swollen, thickened, or hypertrophied tissue in the neighbourhood; or the injury to the ear may be caused by disturbance of that muscular mechanism by which the walls of the cartilaginous part of the tube are, in turn, separated or approximated. Such occlusion of the tube, going on for a length of time, will inevitably lead to permanent changes of structure, and irretrievable defect of function in the organ of hearing.

Upon the due permeability of the Eustachian tube depends the proper degree of tension of the tympanic membrane and the ossicular chain; and there is probably nothing which is so essential to the normal function of the organ of hearing, as the preservation of a due amount of tension in these parts.

It is not possible, in so short a time as it is proper that I should occupy your attention, to go over all the various ways in which morbid conditions of the mucous membrane, of the muscles, and of the nerves of the nasal passages, and naso-pharynx, may and do act upon the ear. I desire just to occupy a few minutes by adverting to a subject which has recently received great attention—not greater, however, than its importance deserves. I refer to the so-called adenoid vegetations in the upper pharynx. We are much indebted to Meyer of Copenhagen, and to Guye of Amsterdam, for their valuable labours in connection with this subject; and, recently, to Loewenberg of Paris, for his important treatise.

As you know, the mucous membrane of the naso-pharynx is very rich in mucous glands. But, besides these, Luschka has described a mass of glandular tissue, which he denominates the pharyngeal tonsil, found chiefly in the roof of the pharynx, but which also extends from the roof along the posterior and lateral walls into Rosenmüller's fossa and over the mouths of the Eustachian tubes. The so-called adenoid vegetations consist of an exuberant growth or hypertrophy of this adenoid tissue, and appear in the form of either tongue-shaped, or globular, or flat excrescences, chiefly on the postero-superior wall, from which, however, they not unfrequently extend to the posterior nares, where they interfere with the due permeability of the nasal passages. The pharyngeal mouths of the Eustachian tubes are often completely veiled by the mass of growth. Meyer found that $7\frac{1}{2}$ per cent. of persons who consulted him with affections of the ear suffered from this morbid state of the naso-pharynx. In connection with and dependent upon this condition, the ear is variously affected; sometimes we find purulent disease, and sometimes simple catarrh, resulting from or maintained by these morbid growths.

Disease in the ear may be caused by these vegetations in various ways: 1. By encroaching upon the pharyngeal mouth of the tube, and thus bringing about mechanical closure; 2. By the concomitant inflammation of the mucous membrane of the Eustachian tube, producing swelling or thickening, with more or less secretion; 3. By the extension of that inflammation to the mucous membrane of the tympanum, or even to the mastoid cells; 4. Through interference with the nasal breathing. When the nasal passages are markedly obstructed, a rarification of the air in the tympanic cavity is produced by every act of swallowing, just as in Toynbee's experiment, and the tympanic membrane with the ossicular chain is forced inwards to an abnormal degree. Or obstruction of the nasal passages may induce mischief in the ear, by causing sudden abnormal condensations of the air in the tympanum. Under the influence of vehement expiratory acts, as coughing, blowing the nose, sneezing, etc., the air, unable to find its way through the nasal passages, may pass through the Eustachian tubes with damaging effect upon the middle ear.

The treatment of these adenoid vegetations is of very great importance, not only for its effects upon the ear, but also in relation to the general health. Although the methods of examination and of treatment employed in diseases of the naso-pharynx do not properly belong to our subject, I would venture to say a word regarding one of the methods of examining the upper pharyngeal cavity, and one of the modes of treating these adenoid growths which I have found very useful. I refer to digital examination, and to the treatment of adenoid growths by the finger-nail. Since I have begun to explore more frequently the upper pharyngeal cavity with the index finger in persons having disease of the ear, I have found with surprising frequency a hypertrophied condition of the glandular tissue in the vault of the pharynx. By this mode of examination, we can, in most of the cases, gain a pretty clear notion of the condition of the upper surface of the soft palate, of the posterior edge of the nasal septum, of the posterior end of the inferior spongy bone, of Rosenmüller's fossa, and of the vault of the pharynx. I might just remark here that it is a mistake to judge of the condition of the upper pharyngeal space from the appearance of the back of the

* Introduction to a discussion in the Subsection of Otolary, at the Annual Meeting of the British Medical Association in Ryde, August 1881.

throat. The latter may appear almost quite normal, and yet the nasopharynx may be involved in very important morbid changes. The method of scraping away these morbid growths, especially the softer and flatter forms of the excrescences, by the nail of the index finger, has been introduced chiefly by Dr. Guye of Amsterdam. It is the least painful, and probably the safest mode of operation. It can be employed in children, who are much more frequently affected with these growths, when the use of instruments is quite impossible.

From want of time, I have necessarily omitted many points in connection with this interesting and important subject. I trust these will be taken up by some of my eminent *confrères* who have graced us with their presence.

FORTY-NINTH ANNUAL MEETING OF THE BRITISH MEDICAL ASSOCIATION.

Held in RYDE, Aug. 9th, 10th, 11th, and 12th, 1881.

PROCEEDINGS OF SECTIONS.

SECTION B.—SURGERY.

Wednesday, August 10th.

THE Section was opened by the President, W. MARTIN COATES, M.R.C.S., who delivered an address, which was published at p. 350 of the JOURNAL for August 27th. The President invited discussion upon his paper; and Mr. MACNAMARA (London) and Mr. REGINALD HARRISON (Liverpool) spoke in high approval of the practical nature of the operation, and of the sound principles on which it was based.

Antiseptic Trocar.—Dr. WARD COUSINS (Southsea) exhibited a new antiseptic trocar, the piston of which was made to work under a covering of India-rubber, and which the inventor had found a perfect guard against septic infection. In answer to Mr. J. B. Brierley, he stated that he had used it with complete success in cases of hydrothorax.

Dr. COUSINS also exhibited a new catheter-tube and urine-reservoir.

A New and Reliable Operation for the Cure of Web-Fingers, etc. By A. T. NORTON, F.R.C.S. (London).—The operation consists of cutting small flaps or tongues of skin from the posterior and anterior surfaces of the hand, with the bases of the tongues on a level with the heads of the metacarpal bones, and of attaching them together by their apices after cutting through the web. The flaps unite by the first intention, and thus any re-development of the web is entirely prevented. The points to be considered in the performance of the operation, in order to ensure a good result, are the following. 1. The flaps should be cut thick, so that their vascular supply may be complete, and the chance of sloughing reduced to a minimum. 2. They should be cut rather narrow, with judgment, otherwise they become compressed laterally and bulge upwards at the margins, instead of lying in adaptation to adjacent tissue. Such compression, of course, interferes with a free circulation of blood through the flaps. 3. The tissue between the knuckles should be cut back, or, if necessary, cut away, so that the apices of the flaps may lie well in contact with each other without tension. 4. The flaps must be of sufficient length to prevent tension when the suture is applied. 5. The apices of the flaps are very small in children, so that a very small needle should be selected to carry the suture, in order to avoid unnecessary injury of the flap. 6. The position of the flaps should be carefully arranged, so that the limit of the new web may be in the line with the natural web of the finger. If the flaps unite, no web can re-form. There is no reason whatever why union by the first intention should not take place, if attention be paid to the above recommendations; and thus one of the most troublesome and unsuccessful of operations is converted into one most simple and most certain of success.—The PRESIDENT and Mr. WATKIN WILLIAMS (Birmingham) spoke on the practical nature of the operation.—In answer to Dr. COUSINS, the author stated that he saw no reason why catgut sutures should not be used to connect the flaps, although, in the case related in the paper, he had used Chinese silk. It was of prime importance to cut the flaps thick, so as to secure their after-vitality.

On the Treatment of Stricture by Stretching. By REGINALD HARRISON, F.R.C.S. (Liverpool).—The author advocated the employment of stretching in certain cases of tight stricture of the urethra where retention had occurred and catheterism was difficult. In using the phrase "stricture-stretching" as best describing the object in view, he did not advocate a new instrument or a new method of treatment; it was to the application of well-recognised principles that he wished to

draw attention. Dr. Otis of New York had demonstrated the extent to which the urethra was capable of being dilated; and what was true of the healthy canal would be found to apply, though in a less degree, to the strictured one. The dilator employed by Mr. Harrison was Holt's, with some special adaptations which had been made by Messrs. Krohne and Sesemann. In the first place, there was attached to it a pilot bougie, which, in cases of tight or tortuous strictures, was invaluable. The number of dilating-rods was increased to eight, so that the process of stretching might be gradual; and, in order that there might be no jerking as the several rods were introduced, a spiral spring controlled the separation of the instrument, instead of a screw. The size of the dilator was No. 3 English gauge: the largest rod brought it up to rather over 12, and between these two extremes there were seven gradations. The patient being placed under ether, the dilator was passed fairly into the bladder, and then the several rods were slowly introduced until dilatation was completed. When this was done, a catheter was used, and all urine removed. A full-sized bougie was then passed every third or fourth day until the patient had learnt how to do it himself, when he must be enjoined to continue the practice. In conclusion, the author remarked that he was not recommending this as a panacea, nor as prejudicing other operations to which reference had been made, but as a safe and an efficient expedient for meeting immediate difficulties, and simplifying future treatment. If, in all cases of retention occurring with tight stricture, the use of this or a similar instrument followed upon catheterism, both patient and practitioner would be none the worse for the knowledge that, whatever spasm might temporarily do, at all events there was a way to the bladder, capable of admitting a No. 12 bougie, which had been made without any structural detriment.—Mr. MACNAMARA (London) thought that the operation could be applicable to only a few cases, and spoke strongly of the advantages of the old operation of gradual dilatation, the cures after which were often permanent.—Mr. HARRISON agreed in the main with Mr. Macnamara, but remarked that surgeons were often the creatures of circumstances; and he related the case in which he had first been led, almost by compulsion, to adopt this procedure.—Mr. RIVINGTON (London) had used Holt's instrument occasionally in the same "tender and loving way" as Mr. Harrison, and thought it a very valuable addition to our treatment of this disease.—Dr. ROYLE (Manchester) spoke; and Mr. STOKES (Dublin) endorsed entirely the views of Mr. Macnamara as to the benefits of gradual dilatation. He had seen disastrous consequences—no fewer than three deaths—from immediate dilatation, a method which was always followed by a rapid return of the stricture. He spoke also in high favour of Maisonneuve's operation by internal section, which he had adopted in strictures of the membranous portion, as well as of the anterior portions, of the urethra.—Dr. COUSINS and the PRESIDENT having spoken, Mr. HARRISON replied.

A Further Series of Cases of Immediate Cure of Inguinal Hernia. By W. DUNNETT SPANTON, M.R.C.S. (Hanley).—The author related nine additional cases in which his operation for immediate cure of hernia had been performed, making a total of thirty-four cases. Of these, thirty had been quite successful, and the remaining four much benefited. There had been no death. In some of the patients, no truss had been of the slightest use. A tendon or catgut ligature was employed in three of the cases, passed in a similar manner to the screw, and retained *in situ* until the parts became consolidated. The result in these cases was found to be on the whole less satisfactory than in those treated by the screw alone. In them, however, the beneficial influence of Listerism was most marked, and the author advised its use in every such operation in which an animal ligature was employed. The paper concluded with an appeal to surgeons to give the operation for radical cure a fair trial, and not to rest satisfied with recommending the mere use of trusses; more especially to urge parents, in the case of young children, to have them cured while young by some operation which had been proved to be both safe and effectual.—By the President's request, Mr. SPANTON described the operation, an account of which has been already published.—Mr. HARRISON (Liverpool) remarked that thirty out of the thirty-four cases had been spoken of as "more or less" effectually relieved, and inquired what the author meant by "less effectual", as the requirements of the operation seemed hardly met by such a result.—Mr. MACNAMARA (London) thought that the operation could hardly be compared to hare-lip, for this reason, that there was infinitely more risk; thirty-four had lived, but the thirty-fifth or thirty-sixth might die. He had himself seen three cases of death after Wood's operation, and several with much constitutional disturbance.—Dr. GRIFFITHS (Swansea) had seen satisfactory results from the operation; but in one case it was followed by a sharp attack of orchitis, which he regarded as a serious matter, and he would like to know in how many cases Mr. Spanton had met with the same result. For