and wrists, and the ankles, were distinctly bronzed, though not to a very great extent. The discoloration presented no definite boundary, but gradually faded into the healthy skin. The areolæ of the nipples and the garter-lines were not materially darkened in colour. There were two sinuses, discharging thick caseous pus, immediately above the right twelfth rib. Bronchial breathing, with moist sounds and bronchophony, was heard at both apices, with some dullness on percussion. The urine was very much like that met with in chyluria, being of a milky colour, and sweetish slightly ammoniacal smell, but was loaded with pus. I considered the case to be one of Addison's disease, with scrofulous kidney discharging through the loin. The physical examination also clearly showed the coexistence of phthisis. The patient was ordered to take a teaspoonful of syrup of iodide of iron, with a like quantity of vinum ferri, three times a day; while carbolio dressings were applied to the sinuses in the loins. He gradually grew weaker, and the sinuses continued to discharge large thick masses of pus, sometimes as large as the finger of an adult. He died of asthenia on July 2nd, 1880.

On July 3rd, I made a *post mortem* examination, but I was allowed by the friends of the patient to open the abdomen only, so that I am sorry I was not able to confirm the diagnosis of phthisis; but I feel confident, from the physical examination, that phthisis did exist. The body was exceedingly emaciated. There were two large sinuses in the right lumbar region, about the position of the kidney, through which a long piece of caseous matter, about the size of one's finger, projected. The skin of the neck, ankles, back of the hands, and wrists, was of a distinctly brownish colour; the nipples also were pigmented. The left kidney was healthy, but considerably hypertrophied. The left suprarenal capsule appeared to be quite healthy. The right kidney was totally destroyed, and converted into a large irregular caseous abscess, which communicated freely with the sinuses in the loin. The abscess extended upwards to the back of the right lobe of the liver, forming a cavity in that viscus about the size of an orange. The right suprarenal capsule was found intimately adherent to the upper and front wall of the abscess; it was of irregular outline, enlarged, and in section presented numerous cavities formed by broken-down caseous tubercle; in other parts, the structure was more fibrous and firmer. The liver and spleen were enlarged, and distinctly lardaceous.

REMARKS.—The above case seems to me to present several points of interest. It is very rare to find this disease affecting so young a patient. I remember seeing a girl, aged 16, who was the subject of Addison's disease, when I was clinical assistant at Guy's Hospital; but I have never heard of its occurring in so young a patient as the one I have just recorded. Again: both capsules are usually affected; but in this case the left appeared to be quite healthy, and this may be the explanation of the bronzing being only slightly marked. The hypertrophy of the left kidney was, of course, compensatory to the destruction of the right. One would like to know, if it were possible, which was the primary seat of the tubercle—whether the lung, kidney, or capsule; my own impression is that the capsule was involved secondarily to, and as an extension from, the kidney.

CANCER OF THE THIGH REMOVED PAINLESSLY, WITHOUT CHLOROFORM OR ETHER-SPRAY.

By JOHN F. LE PAGE, L.R.C.P.Ed., Durham.

J. J., AGED forty-three years, consulted me, on January 10th, 1879, respecting a tumour of his right thigh. On examination, I found a scirrhous, of the size of a small orange, on the anterior aspect. Ulceration had commenced several weeks previously, and extended over a surface measuring five and a half inches in circumference. It would appear that the nature of the tumour had not been clearly comprehended, for it had been poulticed for three or four weeks. Very extreme pain at times was complained of. I urged the importance of an early removal of the cancer; and, my patient being willing to submit to any mode of operating considered most advisable, I decided, having before had very good results from similar treatment, to destroy the malignant growth previous to its removal. And I did so, notwithstanding the opinion of many surgeons of considerable experience to the contrary, in the belief that some caustics exercise a specific action on the morbid structures of cancer; and that thus their action extends somewhat into surrounding tissues, infiltrated with cancer-cells or plasma, which are thus rendered at least unproductive, whilst the tissues themselves are not disorganised.

Although local anaesthesia was not used in the operation, I used, shortly before it, Richardson's ether-spray producer over the margins of the ulcerating surface, for a distance of about an inch from the edge. Pure nitric acid was then applied to the skin, destroying it for the breadth of half an inch all round. This preparation was neces-

sary, as chloride of zinc, which was to be used, acts very slowly on organised skin.

The cancer was situated about two inches below the point where the middle cutaneous nerve pierces the fascia lata. Immediately above the new margin, I injected five-eighths of a grain of morphia with a long subcutaneous needle. One part of chloride of zinc was then mixed into a paste, with two parts of flour and a little water, and thickly plastered over the whole surface. After six hours, the caustic was removed, and poultices applied. On the seventh day the eschar separated, and careful inspection with a lens revealed, between the healthy granulations, on all sides *debris*, which, on microscopic examination, was found to contain cancer-cells. I decided to again apply the chloride; and, after injecting morphia as before, plastered the caustic freely over the surfaces, continuing the application for three hours. Poultices were again applied, and, on the fifth day, the eschar was easily removed. After sponging the cavity, the beautiful glistening fascia lata was seen quite clean at the bottom, whilst the sides were covered with perfectly healthy granulations. A carbolio acid lotion was used, and the cavity rapidly filled up. In eight weeks, the healing process was completed. On December 2nd, 1881, my patient called on me, that I might see that the cure was altogether radical. In every respect, the leg is practically as good as its fellow, and there is abundant evidence that every vestige of the malignant growth was removed.

My object in recording this case is, not so much to laud the use of escharotics—although, without doubt, in many cases of malignant disease, they are very far superior to the knife—but to note the advantages of the mode of operating which I adopted. The great drawback to the use of caustics is, of course, the intense and protracted torture they inflict, for hours and even days—a torture to which the "stoutest heart" can hardly submit. In this case, the chloride of zinc was twice applied, in all for nine hours; and, neither during that period, nor subsequently, was any pain whatever experienced. My patient conversed freely during the hour I remained with him after the first application of the chloride; and on my returning, five hours later, he told me that he had thoroughly enjoyed a substantial dinner, and had smoked a cigar. This great desideratum, the immunity from pain, was attained by temporarily abrogating, through the direct application of morphia, the power to transmit centripetal impulses of the middle, and perhaps also of a branch of the external cutaneous nerve.

SURGICAL MEMORANDA.

EARLY EXCISION OF THE KNEE-JOINT.

WITH reference to Professor Stokes's paper in the BRITISH MEDICAL JOURNAL of December 10th, it appears to me that much of the existing difference of opinion regarding the propriety of excising the knee-joint arises from the varying conditions under which our patients exist. In London, surgeons at the present time have almost unlimited opportunities for sending hospital patients to convalescent homes in the country. Many of these establishments are admirably managed, and situated in healthy seaside localities. Under these circumstances, the children of our poor suffering from the early stage of joint-disease have all the advantages enjoyed by their more wealthy neighbours. Most of these patients may be cured without excision of the affected joint.

For instance, some days ago, when going round the Westminster Hospital, two patients (one a boy aged 11, and the other a girl aged 9) came to visit me on their return from a residence respectively of nine and seven months in convalescent homes. Both patients had been under my care for pulpy degeneration or granular synovitis of the right knee-joint; and, in addition, the boy's hip-joint was similarly affected. His was almost as unpromising a case as it was possible to meet with; but he has returned a healthy lad, full of vigour, and with joints that will stand many a long day's work.

I cannot comprehend the necessity that exists for excising a slice from the ends of the bones in the early stages of pulpy degeneration of the knee, because the disease (unlike tuberculous osteitis) commences, as a rule, in the synovial membrane, the bones being subsequently implicated. It is true, after removing the ends of the bones, we can get them to grow together, and so do away with the necessity for a synovial membrane. Nor can I understand the objection so strongly urged by Professor Stokes against the plan of treating these cases by incision and drainage, because of the danger incurred by suppuration. The treatment by incision and drainage can be carried out antiseptically, with results, so far as inflammation is concerned, as favourable as that attending his practice after resection of the joint. Before, however, disease of the knee has reached the stage when some surgeons advocate treatment by incision and drainage, Professor Stokes