inferior vena cava to the heart, and how almost immediately underneath the right auricle we have the fibrous opening in the diaphragm for the passage of this vessel, there seems good reason for the opinion, so generally entertained, of the great danger of a complete dislocation of the heart to the right side; the inferior vena cava must be very seriously interfered with, and hence the dread of sudden death in these cases seems only reasonable. In my opinion, however, the great danger of this condition rests on the rapidity of the dislocation; for, if the effusion only increase very gradually, the heart, like every other organ in the body under like circumstances, has time to adapt itself to the new state of things. D.'s case also proves that it is not impossible for an empyema to become absorbed after one or two tapplings; and that it is not necessary to open freely into the pleura, unless symptoms of hectic are present. It tells us, too, that the presence of an effusion for months does not always cause such a binding down of the lung by constricting bands of lymph, as to prevent all chance of its expansion. If this is possible in such a severe case as D.'s, surely it must be much more likely in cases where the effusion is comparatively recent; and hence answers forcibly the argument of those advocates of paracentesis who would in all cases make use of this expedient, without trying, in the first instance, any other form of medicinal treatment. Case No. III also speaks against those who pin their practice on this argument: for here we have a pleurisy existing for an indefinite period, and still the lung completely recovered. Indeed, in the town of Omagh, I have frequently the opportunity of examining the condition of the lungs of a lad who suffered, some years since, from a very considerable acute pleuritic effusion. His illness lasted close on three months, and yet I would defy the most skillful diagnostician or stethoscopist to detect the minutest trace of this boy's previous illness; and yet he was treated medicinally, and not by paracentesis. Finally: Case No. II teaches us the great usefulness of paracentesis, in suitable cases, and how quickly a state of comparative comfort can be accorded by its means to a patient even in a semi-moribund condition.

Case No. III has not so many points of importance, still it conveys lessons of its own, one of which I have just referred to. Does it not also tell us that a considerable pleural effusion may exist without the knowledge of the patient, and without apparently causing the slightest inconvenience?

Since my appointment to the Tyrone Infirmary, I have met with thirty-two cases of pleural effusion; and, in nine instances, I tapped the pleura—in one instance performing the operation five times, in another three times, in four twice, and in three once—with the following results: three were greatly relieved, four were completely cured, and two died apparently directly from the results of the operation. The remaining twenty-three patients were not operated on, and they all completely recovered under careful mercurial treatment (principally Bailey's pill) and repeated blistering. My experience, therefore, of the operation of paracentesis thoracis has not been very favourable; and I would never now attempt it, except in very pressing cases. But, what are the cases you consider pressing? is a question you may justly ask; and to this question, which is of a twofold nature and of the first importance, I shall now very briefly endeavour to supply an answer.

What cases do require paracentesis of the thorax? All cases where an empyema can be diagnosed; or where, owing to displacement or pressure, the action of the heart or lungs is greatly impeded; or when symptoms of death by apnoea or syncope are present; wherever, in fact, with the physical signs of pleuritic effusion, we have orthopnoea, quick feeble pulse, lividity of the countenance, a tendency to delirium, or great displacement of the heart, with discoloration of the nails, congestion of the face and lips, and the other usual signs of non-aeration of the blood. Again: where patients, suffering from a pleural effusion, are manifestly losing ground, and when there is no other ascertainable cause to account for the progressive sinking. The operation may also be necessary to relieve suffering and prolong life in specific lung affections—such as malignant or tubercular disease, and in some cases of pneumo-thorax, uncomplicated with phthisis or other specific disease.

Such are the cases that, in my opinion, require the operation. In empyema, if the fluid reaccumulate after the first tapping, with signs of wasting and much hectic, the pleural cavity should be opened into freely, a thorough drainage established, and the cavity syringed out daily with some weak antiseptic solution.

The cases that do not require paracentesis are, as a general rule, acute pleuritic effusions, or chronic effusions, unaccompanied by any of the urgent symptoms just referred to. In specific diseases of the lungs, or in cases of effusion resulting from disease of other organs—such as hydro-thorax in Bright's disease—it is generally useless, except of course as a palliative, to attempt paracentesis. Sometimes, I have found a partial tapping (the patient being under the influence of mercury) followed by rapid absorption of the fluid. I attribute the satisfactory

result thus obtained to the restoration of the absorbing powers of the pleura, which, if exposed to considerable pressure from a large effusion, must of course be greatly interfered with.

In performing the operation of paracentesis thoracis, it should never be forgotten to make, in the first instance, an exploratory puncture, either with the small needle of an aspirator, or of a hypodermic syringe. This proceeding does no harm, and it confirms the physician in the correctness of his diagnosis.

In conclusion, gentlemen, it is almost superfluous to remark that I am not a very strong advocate of the operation of paracentesis. I was enamoured at first with a few successful cases; and, having carefully studied Trousseau's writings on this subject, I proceeded to follow out his line of practice indiscriminately. Two deaths, however, in rapid succession and from unaccountable causes, quickly made me pause; and I have now learned that the operation is not only one of the most useful, but also one of the most dangerous in surgery, and that great judgment is necessary in selecting the cases suitable for it. Niemeyer, indeed, goes almost further than Trousseau in his praise of the operation; and even asserts that, without it, it is useless to attempt the cure of pleural effusions. He seems entirely to discard medicinal treatment, which, in my hands, has certainly been attended with the most successful results.

TUMOUR OF BOTH FRONTAL LOBES.

By JAMES RUSSELL, M.D., F.R.C.P.,
Physician to the General Hospital, Birmingham.

THE details of the following case are more scanty than its importance merits; they are, however, of a sufficiently distinct and positive nature to justify their being placed on record. They coincide very accurately with the description given by Dr. Ferrier of the result of injury to the antero-frontal lobes, and support the opinion he has expressed as to the probable function of this part of the brain. Indeed, I find that the expressions used by Dr. Harvey and myself in describing the patient's condition were precisely those which Dr. Ferrier had employed in referring to the mental phenomena exhibited by monkeys in which the antero-frontal lobes had been destroyed. The absence of the more ordinary symptoms of cerebral disease, those usually associated with lesion of other parts of the cerebrum, so far as my imperfect examination extended, is equally striking with the positive aspect of the case, and again agrees with general experience in relation to this class of cases. I may call attention to the fact of the disease existing in the corresponding part in each hemisphere.

I only received a single visit from the patient, and on that occasion my examination unfortunately did not extend beyond the prominent symptoms. He was fifty-three years of age, and was sent to me by my friend Dr. Harvey. His wife, who accompanied him, complained that through the winter her husband had been in a kind of lethargy, as she said; he sat and seemed to doze, and was indifferent to things going on about him. He also made mistakes in business, and sometimes passed his urine in bed. She spoke of a tendency to delusion; but this tendency, Dr. Harvey tells me, never exceeded "a wrong notion about the time of the day or the day of the week, and sometimes a wondering whether he were in his own house". When he visited me, there was no appearance of paralysis. Sensation was not tested. He walked into my room, and one would hardly have noticed that there was much amiss besides a considerable amount of indifference and a degree of dulness. He answered my questions rather slowly, but clearly, but did not volunteer any statement; all information had to be obtained by direct questioning. He had some frontal headache; but it was not prominent. One positive symptom there was—greatly obscured vision; and this symptom seemed to have appeared with rapidity. He could just read No. 16 Jäger, and his wife believed he could have read No. 1 six weeks previously. The pupils were large; both discs were swollen, congested, spotted with blood, and the vessels were concealed. The patient died about six weeks after this visit, the lethargy deepening, but no new symptom appearing, except a degree of paralysis of the left third nerve at the last.

On *post mortem* examination, a tumour, no doubt malignant, closely connected with the brain-tissue, and rather under the size of a Tangerine orange, was found in the anterior part of the right frontal lobe. A similar but much smaller tumour existed in the corresponding part of the left hemisphere; and one of the same size with the last mentioned at the extremity of the right occipital lobe. All other parts of the brain were perfectly healthy. The paralysis of the third nerve was not accounted for by any *post mortem* appearance.