It was thought that it would be safer and easier to remove the whole mass, which was accordingly done, rather than attempt removal of the tumour alone. The stump was secured by a wire clamp, and its serous outer surface was attached by silk to the abdominal wound. Two thick silk ligatures were also tied round the stump for security. Very little blood was lost, and much care was taken to prevent any from being left behind the bladder, or on the vaginal roof, which appeared to be considerably dragged upwards by the clamped stump. A glass drainage-tube was inserted just above the stump, and perchloride of iron was applied to the end of the stump. After the operation, which lasted about an hour and a half, the finger passed into the vagina detected the cervix high up, but otherwise normal, showing that the wire had encircled the uterus at about the level of the inner os, and had not included any of the vaginal roof with danger to the ureters. Only about four ounces of ether were administered. The mass removed weighed nearly 9lbs., and contained the foetus with membranes intact. The recovery after this formidable operation was uninterrupted, and is very largely to be attributed to the care and skilful treatment of Dr. Dukes who had the sole charge of the patient after the operation. She may now (August 25th) be said to be quite well. The breasts secreted milk on the fourth day. The clamp came away on the twenty-first day.

This is the second time that Porro's operation has been successfully performed in this country, so far as I have been able to learn. The first case was, very curiously, done by Mr. Knowsley Thornton at the Samaritan Hospital the day before this one (see BRITISH MEDICAL JOURNAL, July 22nd); and it seems to me that it ought to be more generally successful than the records from abroad show it to have been. Dr. Alexander Simpson's compiled table shows recoveries 41.6 per cent., and deaths 58.3 per cent.; and in Italy, where it has been done 38 times, the deaths were 24, or 63 per cent., and the recoveries 14, or 37 per cent. Experience and observation up to the present time incline me to think that the extraperitoneal method of treating the stump will be found to be the best, more frequently than the intraperitoneal.

PRELIMINARY IRIDECTOMY IN EXTRACTION OF CATARACT.*

By FRANK H. HODGES, F.R.C.S. Edin.,
Ophthalmic Surgeon to the Leicester Infirmary.

ON my appointment as Ophthalmic Surgeon to the Leicester Infirmary, five years ago, I sought the advice of Mr. Critchett and Dr. Snellen of Utrecht, as to the safest method of extracting cataract. Both were emphatically in favour of performing an iridectomy some weeks prior to the extraction; Mr. Critchett remarking, "If my eternal salvation depended upon the success of a single case of cataract, I would do a preliminary iridectomy;" and Dr. Snellen, "Were I to operate on my own father I should do a preliminary iridectomy." Dr. Snellen further informed me that his custom was to tell his patients that the usual operation at one sitting (Graëfe's method) was a very successful procedure, but that dividing the operation by doing an iridectomy first, and extracting the cataract later on, was somewhat safer, though necessarily the cure was a little delayed; then leaving the choice of operation to the patient.

With regard to the interval that should elapse between the iridectomy and the extraction, Dr. Snellen's rule was "not before six weeks, preferably eight." I have hitherto maintained the latter interval, though I think the rule that my friend, Mr. Priestley Smith, adopts, of doing the extraction as soon as any vascularity in the region of the iridectomy wound has subsided, is perhaps better than any arbitrary fixed interval.

Of the comparative advantages of this method of extraction, if any, over the more usual methods, I have no personal experience, having during the last five years performed preliminary iridectomy in all cases of extraction of senile cataract. Experience of it, however, leads me to claim several conspicuous advantages; and of these I would place first, "absence of hæmorrhage from the iris at the time of the extraction." In my first case in private practice, the subject of which was a gouty old lady, on division of the iris the anterior chamber filled with blood, which occupied six weeks in absorption. A sharp attack of gout came on four days after the iridectomy, but fortunately did not affect the eye. At the extraction eight weeks later, another attack of gout occurred, and this time the iris became involved, but yielded promptly to leeches and atropine, with colchicum internally. The old lady made an excellent recovery, and can now read No. 1 fluently. The cataract had a small hard nucleus with abundant soft cortical substance, which I

could not completely remove, and I believe that the preliminary iridectomy turned the scale from failure to success; for, had the complication of hæmorrhage in a chamber with cortical substance been added to the existing one of gouty iritis, it is highly probable the eye would have been lost.

Besides the greater safety to the eye from the absence of blood in the anterior chamber, it is a manifest advantage to the operator not to have the parts obscured in the important stages of laceration of the capsule and extraction of the cataract. Another advantage of dividing the operation is the experience gained at the first as to how the anæsthetic is borne. My practice is to always to give an anæsthetic for the iridectomy, ether or ethidene bichloride for patients under sixty, and chloroform for those over*; and then, if this be badly borne, or cause much after-sickness, for the extraction either to give no anæsthetic or substitute another. Of the minor advantages, I would instance the tranquility of mind with which a patient awaits the extraction after a painless iridectomy followed by the usual rapid recovery. I have had several patients, who prior to the iridectomy were dreadfully nervous about it, or more usually the anæsthetic; who, upon suffering no inconvenience from the anæsthetic nor pain from the iridectomy, submitted calmly to the subsequent extraction. Again, cataract not unfrequently happens to people in excellent health, who have never been laid up for a day, and to whom confinement is necessarily irksome. The little preparatory training which they get for the iridectomy stands them in good stead for the all-important extraction, and accustoms them to the new sensation of having the eyes bandaged, and of lying on only one side or the back, and to the maintenance of perfect quiet. These latter may seem trifles, but success in eye-surgery often depends upon scrupulous attention to many minor points of detail.

CLINICAL MEMORANDA.

NOTE ON THE AURICULAR PULSATION OF MITRAL STENOSIS.

AS I finish the perusal of Dr. Balfour's paper in the current number of the BRITISH MEDICAL JOURNAL for August 12th, and the discussion thereupon, which took place at the late annual meeting, I am moved to write, while the subject is in memory, to say that I am quite in accord with the views enunciated by Dr. Broadbent, that any pulsation on the chest-wall, and tracings derived therefrom, can hardly own an auricular source. I say this upon the strength of a good deal of observation. Dr. Balfour's writings on this as on all subjects, interesting and pregnant as they ever are, arrested my attention long ago, as also did Dr. Gibson's paper and tracings when they were first published; and, so far as opportunity has offered since that time, I have taken occasion to ascertain in the deadhouse the relations existing between the left auricular appendix and left auricle at the surface, in cases of mitral constriction; and, though I have seen many cases where the auricle and the auricular appendix have been dilated, neither has ever come so near to the surface as to make it at all probable, to my mind, that any pulsation could be appreciable enough to record upon the chest-wall. I have never seen even the most dilated left appendix which was not a considerable depth below the surface; and, discounting the dead for the living position, I still am more than in doubt. On the other hand, three or four cases have occurred in which, during life, a marked pulsation has been observed in the second left interspace, and in all, it has been adequately explained by the existence at the necropsy of a much dilated pulmonary artery. Dilatation of the pulmonary artery is a condition which occurs under diverse circumstances, and it is not uncommon. Its not unfrequent occurrence disposes, I think, of Dr. Balfour's objections to the basic murmur being of pulmonary origin, viz., that there is no possible cause of murmur in the pulmonary artery which is not equally operative at the aortic orifice, or indeed more so. Given high tension in both systemic and pulmonary circulation, I believe the pulmonary artery dilates readily, and the aorta much less so.

I should be sorry to think that the existence of curable dilatation of the left ventricle in chlorosis, so ably maintained by Dr. Balfour, was in any way endangered by a disbelief in the auricular impulse; but, whether it be so or not, I am of opinion that the pulsation observed in some of these cases, and the *bruit* heard in many are probably due to waves in a dilated pulmonary artery, the dilatation being consequent upon mitral regurgitation, and to the resulting obstruction to the pulmonary circulation.

JAMES F. GOODHART, M.D., F.R.C.P.,
Assistant Physician to Guy's Hospital.

* Read before the Annual Meeting of the Midland Branch.

* In deference to the opinion of Mr. Jonathan Hutchinson that, for patients over sixty, chloroform is the safest anæsthetic.