Managing uninvestigated dyspepsia in primary care

Test and treat for *Helicobacter pylori* is still as good as proton pump inhibitors first

RESEARCH, p 651

Naoki Chiba associate clinical professor of medicine, McMaster University, Hamilton, ON, Canada, L8S4L8

naoki.chiba@rogers.com

Competing interests: NC has received honoraria from Abbott, Astra-Zeneca, Janssen, and Nycomed for giving talks about proton pump inhibitor and dyspepsia. NC has also received research support from Astra-Zeneca.

Provenance and peer review: Commissioned; not externally peer reviewed.

BMJ 2008;336:623-4

doi: 10.1136/bmj.39497.622720.80

In their accompanying paper, Delaney and colleagues report a randomised controlled trial of 699 people with uninvestigated dyspepsia, which compares the strategy of “test and treat” for *Helicobacter pylori* infection against empirical acid suppression using a proton pump inhibitor.¹ The test and treat group was tested for *H pylori* infection and positive patients were given eradication treatment, whereas negative patients were given proton pump inhibitors. Previous studies have looked at people who were *H pylori* positive² or *H pylori* negative,³ but none has used test and treat as an overall strategy—instead, if patients were *H pylori* positive they were randomised to eradication treatment or placebo; if they were negative they were randomised to proton pump inhibitors, other drugs, or placebo.

Dyspepsia is a common symptom complex of epigastric pain or discomfort—which includes symptoms of heartburn, acid regurgitation, excessive belching, increased abdominal bloating, nausea, feeling of abnormal or slow digestion, or early satiety—for which patients seek medical care.⁴ Upper gastrointestinal endoscopy should be performed in patients with alarm symptoms such as weight loss, vomiting, or overt bleeding. However, most patients can be safely managed with empirical treatment.^{2 3 5 6} Patients who present to primary care with dyspepsia usually have several symptoms, but the symptoms do not predict the diagnosis.⁶ Recent algorithms for the management of dyspepsia have been published in the United Kingdom,⁷ Canada,⁴ and the United States.⁸

In the UK, the National Institute for Health and Clinical Excellence (NICE) recommends that patients with epigastric pain and heartburn should be managed in the same way, rather than arbitrarily considering that epigastric pain represents dyspepsia (which is not a diagnosis) and that heartburn diagnoses gastro-oesophageal reflux disease.⁷ In Canada, it is recommended that patients with uninvestigated dyspepsia who have mainly reflux symptoms should be treated as if they have gastro-oesophageal reflux disease and given empirical proton pump inhibitors.⁴ In patients with non-heartburn predominant symptoms of dyspepsia, *H pylori* can be tested for non-invasively and eradicated if the result is positive. It is not cost effective to perform endoscopy before treatment.⁹ In the Canadian CADET-Hp study, patients with uninvestigated dyspepsia who were *H pylori* positive were given *H pylori* eradication treatment or empirical treatment with a proton pump inhibitor.² Treatment was successful—a final score of 1-2 (none or minimal) on a seven point Likert scale—in 50% of

patients who had eradication treatment compared with 36% in those given a proton pump inhibitor ($P=0.02$). The number needed to treat over one year was seven for one treatment success. This strategy was also cost effective.¹⁰ In a post hoc subgroup analysis, patients with predominant reflux symptoms who had *H pylori* eradication had 43% treatment success compared with 32% treatment success in those who received placebo antibiotics ($P=0.08$). Dyspepsia did not seem to worsen in this group of patients.¹¹

Delaney and colleagues followed the NICE guidelines and studied patients with dyspepsia who had epigastric pain, heartburn, or both.¹ This was a pragmatic trial that allowed the practitioner to manage the patient according to their discretion during the next year. At 12 months, the patients in the two treatment groups did not differ significantly in terms of dyspeptic symptoms, quality adjusted life years, or costs. The test and treat arm had higher initial costs, but over the course of the year there was a trend towards reduced use of resources (for example, fewer ultrasound scans, endoscopies, primary care consultations), although the only significant reduction was in *H pylori* testing. Delaney and colleagues’ trial lasted one year, so if *H pylori* was responsible for dyspepsia in some patients, cost effectiveness might have improved over a longer time period. However, this might be offset by the large numbers of patients with ongoing symptoms who need continued drug treatment.

Eradicating *H pylori* can be beneficial; for example, it can cure ulcers. Eradication may reduce the potential risk of developing gastric adenocarcinoma or mucosa associated lymphoid tissue (MALT) lymphoma in some patients. Eradication may also provide a small benefit in people with investigated or functional dyspepsia.¹² Primary care patients at first presentation could have any of the above diagnoses, so the strategy of eradicating *H pylori* as a first line strategy continues to have merit.

The prevalence of *H pylori* is declining in the general population. Delaney and colleagues’ study indicates that the prevalence of *H pylori* in the UK is about 30%, which is comparable to that seen in Canada.² Where the prevalence of *H pylori* is lower, empirical treatment with proton pump inhibitors may be more cost effective, whereas test and treat may be better when the prevalence of *H pylori* is higher. Delaney and colleagues pointed out that it was difficult to recruit for their study as so many patients with dyspepsia had already been treated for *H pylori*.

So what strategy should we use? Although this study suggests that the test and treat strategy and the policy of

giving proton pump inhibitors first are equally effective, practitioners can choose which strategy to use according to the predominant symptom complex. Delaney and colleagues' study included patients with both predominant heartburn and epigastric pain. Simply having heartburn does not mean that the patient has gastro-oesophageal reflux disease. Indeed, some patients with heartburn and epigastric pain might respond well to *H pylori* eradication. The good news is, if the prevalence of *H pylori* is high enough, we cannot go wrong with either strategy. If *H pylori* test and treat is used, a reliable method such as urea breath test or stool antigen must be used because *H pylori* serology is unreliable. In uninvestigated patients without alarm symptoms, the initial choice of treatment still depends on individual considerations for each patient.

- 1 Delaney BC, Qume M, Moayyedi P, Logan RFA, Ford AC, Elliott C, et al. Helicobacter pylori test and treat versus proton pump inhibitor in initial management of dyspepsia in primary care: multicentre randomised controlled trial (MRC-CUBE trial). *BMJ* 2008 doi: 10.1136/bmj.39479.640486.AE.
- 2 Chiba N, Veldhuyzen van Zanten SJO, Sinclair P, Ferguson RA, Escobedo S, Grace E. Treatment of *H pylori* infection in primary care patients with uninvestigated dyspepsia improves symptoms: the CADET-Hp (Canadian adult dyspepsia empiric treatment-Helicobacter pylori positive) randomised controlled trial. *BMJ* 2001;324:1012-6.
- 3 Veldhuyzen van Zanten SJ, Chiba N, Armstrong D, Barkun A, Thomson A, Smyth S, et al. A randomized trial comparing omeprazole, ranitidine, cisapride, or placebo in Helicobacter pylori negative, primary care patients with dyspepsia: The CADET-HN study. *Am J Gastroenterol* 2005;100:1477-88.
- 4 Veldhuyzen van Zanten SJ, Flook N, Chiba N, Armstrong D, Barkun A, Bradette M, et al. An evidence-based approach to the management of uninvestigated dyspepsia in the era of Helicobacter pylori. Canadian Dyspepsia Working Group. *CMAJ* 2000;162(suppl 12):S3-23.
- 5 Armstrong D, Veldhuyzen van Zanten SJ, Barkun AN, Chiba N, Thomson AB, Smyth S, et al. Heartburn-dominant, uninvestigated dyspepsia: a comparison of "PPI-start" and "H2-RA-start" management strategies in primary care—the CADET-HR study. *Aliment Pharmacol Ther* 2005;21:1189-202.
- 6 Thomson AB, Barkun AN, Armstrong D, Chiba N, White RJ, Daniels S, et al. The prevalence of clinically significant endoscopic findings in primary care patients with uninvestigated dyspepsia: the Canadian adult dyspepsia empiric treatment-prompt endoscopy (CADET-PE) study. *Aliment Pharmacol Ther* 2003;17:1481-91.
- 7 National Institute for Health and Clinical Excellence. *The management of dyspepsia in adult patients in primary care*. London: NICE, 2004.
- 8 Talley NJ, Vakil NB, Moayyedi P. American gastroenterological association technical review on the evaluation of dyspepsia. *Gastroenterology* 2005;129:1756-80.
- 9 Ford AC, Qume M, Moayyedi P, Arents NL, Lassen AT, Logan RF, et al. Helicobacter pylori "test and treat" or endoscopy for managing dyspepsia: an individual patient data meta-analysis. *Gastroenterology* 2005;128:1838-44.
- 10 Chiba N, Veldhuyzen van Zanten SJ, Escobedo S, Grace E, Lee J, Sinclair P, et al. Economic evaluation of Helicobacter pylori eradication in the CADET-Hp randomized controlled trial of *H pylori*-positive primary care patients with uninvestigated dyspepsia. *Aliment Pharmacol Ther* 2004;19:349-58.
- 11 Chiba N, Veldhuyzen van Zanten SJO, Thomson ABR, Barkun AN, Armstrong D, Sinclair P, et al. Reflux symptoms should not be excluded from the definition of uninvestigated dyspepsia: results from the CADET-Hp study. *Am J Gastroenterol* 2001;96(suppl 1):A20.
- 12 Moayyedi P, Soo S, Deeks J, Delaney B, Harris A, Innes M, et al. Eradication of Helicobacter pylori for non-ulcer dyspepsia. *Cochrane Database Syst Rev* 2006;(4):CD002096.

Statins for people with kidney disease

Criteria for treatment should be the same as for people with normal kidney function

RESEARCH, p 645

M Clase associate professor of medicine, McMaster University, Hamilton, ON L8N 1Y2
clase@mcmaster.ca

Competing interests: CMC has organised educational rounds sponsored by drug companies that make statins.

Provenance and peer review: Commissioned; not externally peer reviewed.

BMJ 2008;336:624-5
doi: 10.1136/bmj.39483.665139.80

In the accompanying paper, Strippoli and colleagues report a meta-analysis of the benefits and harms of statins in patients with chronic kidney disease. They report outcomes for people with low glomerular filtration rate, those on dialysis, and those with kidney transplants.¹

Chronic kidney disease (glomerular filtration rate < 60 ml/min/1.73 m² or urine abnormalities) and cardiovascular disease often coexist; chronic kidney disease also predicts cardiovascular events. A recent meta-analysis reported that low glomerular filtration rate was associated with an unadjusted relative risk of 2.5 (95% confidence interval 1.4 to 4.3) for cardiovascular events.² In adjusted analyses, some of this effect is explained by the high prevalence of traditional risk factors for cardiovascular disease in people with low glomerular filtration rate or albuminuria.^{3 4} However, non-traditional risk factors such as anaemia (which in patients on dialysis predicts left ventricular hypertrophy, systolic dysfunction, and congestive heart failure⁵) and factors favouring vascular calcification (which in patients on dialysis predicts death⁶) may contribute to death from cardiovascular disease in people with a low glomerular filtration rate or those on dialysis. So statins may not be as effective for preventing cardiovascular events in these groups as they are in the general population.⁷ Perhaps because of this concern, or because of worries about increased harm or polypharmacy,

people with low glomerular filtration rate are less likely to be prescribed statins after a coronary event.⁸

In people at high risk of cardiovascular events because of established cardiovascular disease or conventional risk factors, does a low glomerular filtration rate affect the benefits or harms of statins? In Strippoli and colleagues' analysis of patients with pre-dialysis (stages I-IV) chronic kidney disease, which includes five studies, 18 176 people with low glomerular filtration rate, and 1591 events, the relative risk of all cause mortality was 0.81 (0.74 to 0.89) in patients randomised to statins. This is consistent with data from the general population^{9 10}: statins produced a clinically important reduction in the overall risk of death of 19%. However, this analysis is based on studies mainly involving patients with a glomerular filtration rate above 30 ml/min/1.73 m² (stages I-III). Only a few people who are not on dialysis and have a glomerular filtration rate below 30 ml/min/1.73 m² (stage IV) have been included in trials.

The meta-analyses of studies of people on dialysis and those with kidney transplants found no benefit in terms of all cause mortality. Strong trends to reductions in death from cardiovascular events were seen in both groups, however, with point estimates of a 17% reduction in death from cardiovascular disease in people on dialysis and 32% in people with kidney transplants. In terms of harm, they found no significantly increased

risk of abnormalities of liver function or creatine kinase (relative risk 1.5, 0.9 to 2.6).

These data suggest that the benefit of statins seen in the general population also applies to people with low glomerular filtration rate, those on dialysis, and those with kidney transplants, provided they have established cardiovascular disease or a cardiovascular risk profile that justifies treatment.

In people who are at high risk of cardiovascular disease mainly because of chronic kidney disease (low glomerular filtration rate or albuminuria), do statins prevent cardiovascular events? No large scale randomised controlled trials are available to answer this question. The SHARP (study of heart and renal protection) should be able to answer this question directly. The trial has enrolled 9000 people who are on dialysis or have a low glomerular filtration rate, but no previous cardiac event, and has randomised them to combined simvastatin and ezetimibe or to placebo.^{7 11} But until these results are available, low glomerular filtration rate or dialysis alone should not be considered indications for treatment with statins.

Finally, in people with low glomerular filtration rate, do statins prevent progression of renal insufficiency and progression to end stage renal disease? Strippoli and colleagues found that statins modestly reduced proteinuria but did not affect the rate of decline of glomerular filtration rate. A previous meta-analysis found similar effects on proteinuria and detected a small reduction in the rate of decline of glomerular filtration rate,¹² perhaps because a difference in inclusion criteria led to increased power. None the less, surrogate outcomes such as these are unconvincing, and no trials have reported the clinically important outcome of progression to end stage renal disease. The effect of treatment to lower cholesterol on this outcome will be looked at in a secondary analysis of the SHARP study.

Available evidence suggests that the threshold of cardiovascular risk for treating most patients with a

low glomerular filtration rate (30-60 ml/min/1.73 m²) should be the same as for those with a higher rate. This approach should probably be extended to people with a glomerular filtration rate below 30 ml/min/1.73 m², those on dialysis, and those with transplants. The SHARP trial will definitively answer the question of whether low glomerular filtration rate or dialysis are indications for cholesterol lowering treatment for the prevention of cardiovascular events and progression to end stage renal disease. Pending its results, data do not support the use of statins for these reasons.

- 1 Strippoli GFM, Navaneethan SD, Johnson D, Perkovic V, Pellegrini F, Nicolucci A, et al. Effects of statins in patients with chronic kidney disease: meta-analysis and meta-regression of randomised controlled trials. *BMJ* 2008 doi: 10.1136/bmj.39472.580984.AE.
- 2 Tonelli M, Wiebe N, Culleton B, House A, Rabbat C, Fok M, et al. Chronic kidney disease and mortality risk: a systematic review. *J Am Soc Nephrol* 2006;17:2034-47.
- 3 Hallan S, Astor B, Romundstad S, Aasard K, Kvenild K, Coresh J. Association of kidney function and albuminuria with cardiovascular mortality in older vs younger individuals: the HUNT II study. *Arch Intern Med* 2007;167:2490-6.
- 4 Tonelli M, Jose P, Curhan G, Sacks F, Braunwald E, Pfeffer M, et al. Proteinuria, impaired kidney function, and adverse outcomes in people with coronary disease: analysis of a previously conducted randomised trial. *BMJ* 2006;332:1426.
- 5 Parfrey PS, Foley RN, Harnett JD, Kent GM, Murray DC, Barre PE. Outcome and risk factors for left ventricular disorders in chronic uraemia. *Nephrol Dial Transplant* 1996;11:1277-85.
- 6 Matsuoka M, Iseki K, Tamashiro M, Fujimoto N, Higa N, Touma T, et al. Impact of high coronary artery calcification score (CACS) on survival in patients on chronic hemodialysis. *Clin Exp Nephrol* 2004;8:54-8.
- 7 Baigent C, Landry M. Study of heart and renal protection (SHARP). *Kidney Int Suppl* 2003;63:207-10.
- 8 Keough-Ryan TM, Kiberd BA, Dipchand CS, Cox JL, Rose CL, Thompson KJ, et al. Outcomes of acute coronary syndrome in a large Canadian cohort: impact of chronic renal insufficiency, cardiac interventions, and anemia. *Am J Kidney Dis* 2005;46:845-55.
- 9 Baigent C, Keech A, Kearney PM, Blackwell L, Buck G, Pollicino C, et al. Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90 056 participants in 14 randomised trials of statins. *Lancet* 2005;366:1267-78.
- 10 Ross SD, Allen IE, Connelly JE, Korenblat BM, Smith ME, Bishop D, et al. Clinical outcomes in statin treatment trials: a meta-analysis. *Arch Intern Med* 1999;159:1793-802.
- 11 ClinicalTrials.gov. SHARP: study of heart and renal protection. 2008. <http://clinicaltrials.gov/ct2/show/NCT00125593?term=sharp&rank=9>.
- 12 Sandhu S, Wiebe N, Fried LF, Tonelli M. Statins for improving renal outcomes: a meta-analysis. *J Am Soc Nephrol* 2006;17:2006-16.

Assessing the options available to Lord Darzi

Time to look at both the system and the patient as a whole

Martin Roland director, National Primary Care Research and Development Centre, University of Manchester, Manchester M13 9PL
m.roland@manchester.ac.uk

Competing interests: MR receives research funding from the Department of Health. The opinions are those of the author and not of the Department of Health.

Provenance and peer review: Commissioned; not externally peer reviewed.

BMJ 2008;336:625-6
doi: 10.1136/bmj.39510.702234.80

The Department of Health in the United Kingdom is undertaking a major review of National Health Service policy ("the Darzi review"). Which options are most likely to produce an effective, efficient, and patient centred health service? The UK has made major progress in reducing avoidable deaths and improving care for chronic diseases.^{1 2} However, many areas still have considerable room for improvement.

Strong primary care is associated with reduced costs and improved outcomes.³ The review therefore needs to strengthen primary care. The UK system of universal registration with a single general practice promotes equity, provides a "medical home" for coordinating care, and is an effective mechanism for holding pro-

viders to account for the quality of care provided. This must be retained, even though patients may occasionally consult other practitioners—for example, a doctor near their workplace.

Much has been written about "polyclinics"—facilities that group together primary care practitioners with diagnostic and specialist services. Despite lack of clarity about their purpose, several are already planned. Part of the debate about polyclinics relates to a perceived need to merge small practices in highly deprived areas where the quality of medical services is low. Providing good premises and facilities for such practices could make a big difference to care and would enable the practices to pool resources to

EDDIE MULLHOLLAND/REX

You can listen to Ara Darzi in conversation with BMJ editor Fiona Godlee on our website. Visit www.bmj.com/audio/archive/ara-darzi.dtl to access the interview

provide a wider range of services. However, whether practices should merge within such facilities is a separate question.

The NHS goal of providing patient choice in primary care is not realised in many parts of the country where patients have little real choice of practice. Increased patient choice requires more high quality practices, not the small number of large practices that some polyclinic models suggest. We know that patients in small practices rate their care more highly in terms of both access and continuity.⁴ Indeed, although small practices show more variation in quality, on average, they achieved slightly higher levels of clinical quality than larger practices in the quality and outcomes framework.⁵ One approach would be to use polyclinics to provide extra community based resources but to use them to house general practitioner practices only when local buildings or access are poor. Additional practices are still needed to increase patient choice.

Some models for polyclinics include a greater role for specialists working in the community (bringing services "closer to home"), and government policies are already moving specialists out of hospitals and training primary care staff to take on new specialist roles. However, specialists may be less efficient when deployed outside hospitals.⁶ A recent evaluation of NHS closer to home demonstration sites suggests that these new services improve access when provided as an additional resource. However, their economic justification is much less clear when they are designed to substitute for existing hospital based services. Specialists and practitioners have raised concerns about the quality and safety of some of these new services, although there is little hard evidence of problems, and patients rate them highly.⁷ The justification for moving specialist services into the community is greatest where patients have long travelling times, and the benefits are small in many urban areas. Training new types of practitioner to provide specialist services can provide effective additional capacity, but only if quality control is good.

Better support for self management is needed for people with long term conditions. Practitioners need to provide patients with better information about their care and involve them more in decisions about care. These are areas where NHS performance is poor.⁸ Existing NHS models of self management need to change so that they are better integrated with existing services, include support and training for practitioners, and organise services around the needs of patients.⁹

The greatest demand on the future NHS will be to provide high quality coordinated care for patients with multiple chronic diseases. Recent NHS initiatives have increased the range of providers in both primary care (for example, walk-in centres) and secondary care (for example, independent sector treatment centres). This has the potential to worsen coordination of care—an area in which UK performance is already poor compared with other countries.¹⁰

Several changes are necessary to improve continuity and coordination of care. Firstly, the general practi-

tioner contract should be modified to incentivise personal continuity of care as well as access, especially as three recent studies suggest that many patients in England value personal continuity more highly than rapid access to care.¹¹⁻¹³ Secondly, financial incentives for hospitals to admit patients as emergencies need to be reduced. Instead, hospitals should have incentives to work with providers within the community to provide high quality alternatives to admission. Thirdly, new approaches to commissioning health care are needed. Patients with multiple conditions need better coordination of care between the various sectors of the NHS, and involving hospital specialists in joint commissioning with general practitioners would provide better integration than a policy that focuses exclusively on commissioning led by primary care.

Vigorously pursued policies may deliver on their stated goals but have other unintended effects. For example, strategies designed to reduce waiting times to see general practitioners have made it more difficult for patients to book in advance,¹⁴ and an emphasis on seeing non-urgent new patients within target times can result in other patients coming to harm because their follow-up appointment is delayed.¹⁵ Now is the time to look at both the system and the patient as a whole. That is the challenge for the Darzi review.

- 1 Nolte E, McKee M. Measuring the health of nations: updating an earlier analysis. *Health Affairs* 2008;27:58-71.
- 2 Campbell S, Reeves D, Kontopantelis E, Sibbald B, Roland M. Quality of primary care in England with the introduction of pay for performance. *N Engl J Med* 2007;357:181-90.
- 3 Starfield B, Shi L, Macinko J. Contribution of primary care to health systems and health. *Milbank Q* 2005;83:457-502.
- 4 Campbell JL, Ramsay J, Green J. Practice size: impact on consultation length, workload, and patient assessment of care. *Br J Gen Pract* 2001;51:644-50.
- 5 Doran T, Fullwood C, Gravelle H, Reeves D, Kontopantelis E, Hiron U, et al. Pay for performance programs in family practices in the United Kingdom. *N Engl J Med* 2006;355:375-84.
- 6 Sibbald B, McDonald R, Roland M. Shifting care from hospitals to the community: a review of the evidence on quality and efficiency. *J Health Serv Res Policy* 2007;12:110-7.
- 7 Leese B, Bohan M, Gemmell I, Hinder S, Mead N, Pickard S, et al. Evaluation of "closer to home" demonstration sites. Final report. Manchester: National Primary Care Research and Development Centre, 2007. www.npcrdc.ac.uk/r5.77.
- 8 Coulter A. Engaging patients in their healthcare. *How is the UK doing relative to other countries*. Oxford: Picker Institute Europe, 2006. www.pickereurope.org/Filestore/Publications/Six_country_study_with_ISBN_web.pdf.
- 9 Kennedy A, Rogers A, Bower P. Support for self care for patients with chronic disease *BMJ* 2007;335:968-70.
- 10 Schoen C, Osborn R, Doty MM, Bishop M, Peugh J, Murukutla N. Toward higher-performance health systems: adults' health care experiences in seven countries, 2007. *Health Affairs* 2007;26:w717-34.
- 11 Turner D, Tarrant C, Windridge K, Bryan S, Boulton M, Freeman G, et al. Do patients value continuity of care in general practice? An investigation using stated preference discrete choice experiments. *J Health Serv Res Policy* 2007;12:132-7.
- 12 Rubin G, Bate A, George A, Shackley P, Hall N. Preferences for access to the GP: a discrete choice experiment. *Br J Gen Pract* 2006;56:743-8.
- 13 Cheraghi-Sohi S, Hole A, Mead N, McDonald R, Whalley D, Bower P, et al. What do patients want from a primary care consultation? A discrete choice experiment to identify patient priorities. *Ann Fam Med* 2008;6:107-15.
- 14 Salisbury C, Goodall S, Montgomery AA, Pickin DM, Edwards S, Sampson F, et al. Does advanced access improve access to primary health care? Questionnaire survey of patients. *Br J Gen Pract* 2007;57:615-21.
- 15 House of Commons. Public Administration Select Committee. Fifth report of session 2002-2003. *On target: government by measurement*. 2003. www.publications.parliament.uk/pa/cm200203/cmselect/cmpubadm/62/6202.htm.

Observational evidence for determining drug safety

Is no substitute for evidence from randomised controlled trials

Nick Freemantle professor of clinical epidemiology and biostatistics, Health Care Evaluation Group, University of Birmingham, Edgbaston, Birmingham B15 2TT
N.Freemantle@bham.ac.uk

Atar Irs lecturer in clinical pharmacology, Institute of Pharmacology, University of Tartu, 50411 Tartu, Estonia

Competing interests: NF has received funding from drug companies (but not the manufacturer of aprotinin) for research, consultation, and educational services in the area of drug efficacy and safety. AI is employed by the Estonian State Agency of Medicines (SAM) and is a member of the European Medicines Agency's (EMA) committee for medicinal products for human use (CHMP) and paediatric committee (PDCO). The views expressed in the article are his own and not those of SAM or EMA.

Provenance and peer review: Commissioned, not externally peer reviewed.

BMJ 2008;336:627-8
doi: 10.1136/bmj.39491.493252.80

Drug regulators are constantly criticised for their handling of drug safety. Comment on the activities of the US Food and Drug Administration with regard to drug safety has become particularly loud and damning. A recent article that proposed sweeping changes to the system criticised the FDA for lacking relevant expertise, being too close to the drug industry, and failing to ensure that companies deliver on their safety commitments with new drugs.¹ The main practical recommendation was that systems of monitoring and reporting adverse events should be enhanced, highlighting a perceived role for observational studies. Drug safety is also an important concern in Europe, and the European Commission Enterprise and Industry Directorate General is consulting with stakeholders on changes of legislation to strengthen and rationalise pharmacovigilance in Europe.² It describes a strategy that also focuses on observational studies of drug safety. But can observational studies alone improve judgments on drug safety?

Using adverse event reporting and observational studies to examine safety might seem attractive, but for several reasons observational studies alone cannot provide reliable estimates of treatment effects. For example, safety outcomes can be confounded with patients' prognosis, drugs may be allocated according to clinician's choices rather than by a quasi-random process, and clinicians may be influenced by trying to improve outcomes in people they think are at high risk.

In a properly randomised controlled trial, experimental conditions differ according to chance and the intended treatment, which enables robust inference. In observational studies on drug safety, the way that clinicians select patients can have important effects. Patients treated with an agent of interest may be at increased risk as described by known prognostic factors. This was the case in a recent highly publicised observational study of aprotinin for the reduction of bleeding and risk of transfusion in cardiac surgery,^{3,4} which suggested a 48% increase in the risk of death in the long term associated with the use of aprotinin.⁴ Multivariable methods may help but inclusion of such prognostic factors in a multivariable model does not fully account for their effects. Identified risk factors and their inter-relations are often complex and statistical models deal with these only crudely and partially. Thus a statistical model may include hypertension as just a "yes or no" dichotomy, rather than consider the level of raised blood pressure. In addition, the effects of raised blood pressure may differ in people who do, or do not, have other risk factors in ways that cannot be readily dealt with in analysis. In any case, the multivariable analysis simply estimates the effects of known risk factors from the available data, and the best estimate available from the data may not be a very good one.

Observational studies are further compromised if clinicians make a judgment on whether or not to treat a patient with the drug of interest. In part, this may be haphazard or essentially quasi-random, or it may be based at least on the known prognostic factors and a range of other factors that we cannot fully understand from the summary data collected. Where this clinical judgment accurately predicts outcome for a patient beyond the effects of known risk factors, it represents a latency that is confounded with drug exposure. This will lead to a negative bias in the results of multivariable analyses.

A further problem associated with observational studies is that they can affect our judgment on more reliable measurements. Would the BART trial comparing aprotinin with the antifibrinolytic agents tranexamic acid or epsilon-aminocaproic acid in patients undergoing cardiac surgery have been stopped and aprotinin withdrawn had it not been for the previous observational study indicating risk associated with aprotinin?⁵ This happened even though a recent systematic review of randomised trials found no increased risks associated with treatment.⁶

Information on the size of the increased risk seen in the BART trial is currently limited, but an FDA statement indicates that the adverse signal had not reached conventional levels of statistical significance at the interim analysis of the trial.⁵ It is also not clear whether this conventional level of statistical significance had accounted for multiple testing as is appropriate in interim safety monitoring. Given the lack of a placebo group, the observed difference between the groups may have been caused by superiority of the comparator agents rather than increased risk with aprotinin, although chance also seems a plausible explanation, and stopping the trial early rules out more conclusive evidence on this important question.

So what is the answer to concerns about drug safety? Only properly randomised trials can provide truly reliable evidence on adverse events, just as these are the only source of convincing data on drug efficacy. Observational studies may provide some limited reassurance that a drug is safe, or they may provide an early indication of a problem, but by design they cannot provide reliable evidence on questions of drug safety. Regulators can, and do, require large scale randomised trials to be conducted after launch that look at important emerging safety concerns, and their power and determination to ensure this happens should be enhanced. For example, they could make continued marketing authorisation conditional on the provision of new data from trials which meet pre-agreed standards. If these trials are properly designed they can provide important answers not just about safety concerns, but crucially about the balance of risks and benefits associated with the use of a particular agent.

- 1 Fruberg CD, Levin AA, Gross PA, Shapiro RS, Strom BL. The FDA and drug safety: a proposal for sweeping change. *Arch Intern Med* 2006;166:1938-42.
- 2 European Commission. Strategy to better protect public health by strengthening and rationalising EU pharmacovigilance: public consultation on legislative proposals. 2007. http://ec.europa.eu/enterprise/pharmaceuticals/pharmacovigilance_acs/docs/public-consultation_12-2007.pdf.
- 3 Mangano DT, Tudor IC, Dietzel C. The risk associated with aprotinin in cardiac surgery. *N Engl J Med* 2006;354:353-65.
- 4 Mangano DT, Miao Y, Vuylsteke A, Tudor IC, Juneja R, Filipescu D, et al. Mortality associated with aprotinin during 5 years following coronary artery bypass graft surgery. *JAMA* 2007;297:471-9.
- 5 Food and Drug Administration. *Early communication about an ongoing safety review aprotinin injection (marketed as trasylol)*. 2007. www.fda.gov/cder/drug/early_comm/aprotinin.htm.
- 6 Henry DA, Carless PA, Moxey AJ, O'Connell D, Stokes BJ, McClelland B, et al. Anti-fibrinolytic use for minimising perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* 2007;(4):CD001886.

Better radiology in the *BMJ*

Despite improvements already made, we still have more to do

Amy Davis editorial registrar,
BMA House, London WC1H 9JR
adavis@bmj.com

Competing interests: None
declared.

Provenance and peer review:
Commissioned; not externally
peer reviewed.

BMJ 2008;336:628
doi: 10.1136/bmj.39504.640104.80

In 1976 we published a review of 62 *BMJ* research papers, which showed that 32 of them had statistical faults and five had made claims in their abstracts that were not supported by their results.¹ These mistakes occurred because at that time statisticians were not commonly involved in research papers, either as authors or peer reviewers.² To improve the reliability of published research, the *BMJ* was one of the first medical journals to introduce statistical review of all published research papers, and the role of statisticians in peer review continues to be of interest today.³ We now plan to do the same for radiology, using external advisors and radiologists to guide us and improve our content.

Twenty years ago radiology was mainly plain radiographs viewed on light boxes. Now digital technology means that multislice computed tomography, magnetic resonance imaging, Doppler ultrasound, and even positron emission tomography can be viewed on computers in hospitals and across the world via the internet. We are concerned that the radiology content and images in the *BMJ* may not have kept up with these changes. In June 2007, an external review of the *BMJ* from one of our radiology advisors showed that we had used radiological images in only six articles in the previous six months—four of them in the Practice section and two in the Minerva section. Of these six images, most were not adequately explained, and none had arrows to indicate the relevant abnormalities or findings.

So what have we done to remedy this? Firstly, we have set up a panel of radiologists who will review all images for clinical articles before they are published in the *BMJ* (<http://resources.bmj.com/bmj/about-bmj/radiology-reviewers>). (We do not currently peer review images in the News section as these are journalistic rather than educational.) We aim to use high quality images, with accurate figure legends and arrows to identify the abnormalities. Secondly, we have increased the amount of educational content on radiology. The *BMJ* series Rational Imaging⁴⁻⁶ provides updates for doctors on the best use of different imaging methods for common or important clinical presentations. We welcome submissions for this series (see instructions for authors on bmj.com). Also, from April we will feature a Picture Quiz using radiological and clinical

Guide for *BMJ* image bank

Images must be high resolution at 300 dpi
For the Picture Quiz, images need to be at least 8 cm wide; for all other articles they should be at least 12 cm wide
All images should be sent to us as jpeg files
We need two versions of each image, one that shows just the image and one that includes arrows to indicate the abnormality
All images need to have a figure legend
In the initial stages we would prefer images that show "classic" signs of common illnesses
All images must have a signed *BMJ* patient consent form that specifically allows publication in the *BMJ*
Consent forms can be found at: <http://resources.bmj.com/bmj/authors/checklists-forms/patient-consent-form>

images (<http://resources.bmj.com/bmj/authors/types-of-article>). Thirdly, we are creating a bank of radiology images to use with different articles and would like to encourage readers to send us interesting images, provided they have adequate consent from the patient for publication (box). Finally, we intend to keep readers abreast of "hot topics" and developments in radiology. For example, after discussions with radiologists at the Royal College of Radiology's annual meeting in September 2007, we commissioned a feature about the ethics of private companies offering whole body computed tomography screening.⁷ Other articles are planned.

Like statistics, radiology is an increasingly complex field in which it is possible to get things embarrassingly and dangerously wrong. We hope our panel of radiologists will save our blushes and help you to raise your radiological game. We welcome your feedback and suggestions for Picture Quiz and Rational Imaging articles, which can be emailed to Amy Davis (adavis@bmj.com).

- 1 Gore SM, Jones IG, Rytter EC. Misuse of statistical methods: critical assessment of articles in the *BMJ* from January to March 1976. *BMJ* 1977;i:85-7.
- 2 Lock S. *A difficult balance: editorial peer review in medicine*. London: BMJ Publishing Group, 1985.
- 3 Hughes S. European regulatory agencies should employ full time statisticians. *BMJ* 2007;335:250.
- 4 Rahman NM, Davies RJO, Gleeson FV. Investigating suspected malignant pleural effusion. *BMJ* 2007;334:206-7.
- 5 Burling D, East J, Taylor SA. Investigating rectal bleeding. *BMJ* 2007;335:1260-2.
- 6 Sinha R, Nwokolo C, Murphy P. Magnetic resonance imaging in Crohn's disease. *BMJ* 2008;336:273-6.
- 7 Kmietowicz Z. Computed tomography screening: better safe than sorry? *BMJ* 2007;335:1182-4.