Polymerase chain reaction for salmonella virulence-associated plasmid genes detection: a new tool in salmonella epidemiology

L. REXACH, F. DILASSER AND P. FACH*

Centre National d'Etudes Vétérinaires et Alimentaires/Laboratoire Central d'Hygiène Alimentaire, Unité d'Epidémiologie et d'Innovation Technique en Microbiologie, 43 rue de Dantzig, 75015 Paris, France

(Accepted 22 June 1993)

SUMMARY

The important role of plasmid genes in assessing virulence for BALB/c mice in salmonella, and the difficulty of using standard techniques to detect them, led us to develop a detection method by gene amplification.

One hundred and forty-three strains (71 serovars) of salmonella and 35 strains of other species were tested using specific oligonucleotide primers. The amplification products were identified by a specific oligonucleotide probe. Forty-nine salmonella strains from ten serovars (*S. abortus ovis*, *S. choleraesuis*, *S. dublin*, *S. enteritidis*, *S. gallinarum/pullorum*, *S. hessarek*, *S. typhimurium*, *S. IIIa* 48:z₄, z₂₃, *S. IV* 43:z₄, z₂₃:-, *S. V* 28:a:-) produced a positive and specific response.

Because of various origins of the strains possessing the gene sought and the diversity of the responses, both from one serovar to another and in the same serovar, this search has its place among the epidemiological markers in general use. This method appears well suited to the research and detection of plasmid genes associated with mouse virulence in salmonella.

INTRODUCTION

Animals are known to constitute a vast reservoir of salmonella. The general problem of environmental contamination by organic waste in regard of human and animal salmonellosis [1] has acquired a new importance for the developed countries, due to extended livestock farming. Thus any information which can be obtained on the potential virulence of salmonella is of great interest.

For some years, various publications have illustrated the role of plasmids as virulence-related factors in various serovars of salmonella [2–9]. The genetic determinants of invasiveness are chromosomal [10]. Chromosomal DNA is an important factor in the expression of virulence, especially in the capacity of strains to survive and multiply in reticulo-endothelial system cells [11]. Nevertheless, the virulence of salmonella strains is linked to a combination of chromosomal and plasmid factors [6, 10–13]. So far, no role other than that of their connection with virulence [7, 14] has been attributed to these plasmids, yet for *S. enteritidis*, increased production of long chains of lipopolysaccharides has been noted [15]. In

* Author for correspondence.

1982 Jones [6] showed that these plasmids were linked to adhesion and invasion of HeLa cells in the mouse. In 1985 Chikami [12] showed that, without plasmids, salmonella could invade intestinal mucous membranes, but were unable to propagate infections, at given rates of contamination. Since then, in 1986, Hackett [10] and Pardon [13] have shown that the plasmids were necessary for colonization of the spleen and liver in the mouse. Also, Hackett [10] has demonstrated that strains possessing the plasmids were not phagocytosed by macrophages and thus remained in the organism. Other studies have shown that among various serovars of salmonella there exists a high degree of structural homology in all or some of the plasmids linked with virulence [11, 16–18], to such an extent that cross-immunization has been achieved [19–20].

In the light of the above, it appeared interesting, for salmonella strains, to propose a method for demonstrating the genetic virulence potential for mice linked to the presence of this type of plasmid. Use of a polymerase chain reaction (PCR) seemed a quick, simple and reliable method capable of achieving the objectives set.

MATERIALS AND METHODS

Bacterial strains

One hundred and forty-three strains of salmonella, i.e. 71 serovars and 35 strains of other species of bacteria, were used in this study (Table 1). They were all isolated either from animals (51 strains) or from food products (73 strains) or from the environment (53 strains); one strain was isolated from a human faecal culture following salmonellosis. They were all subjected to biochemical and serological analysis. Bacteria cells were grown in Trypticase Soy Broth (Difco) overnight at 37 °C.

Boiling DNA extraction

A simple and fast technique was adopted to obtain target DNA for PCR analysis. Overnight bacterial culture (1 ml) was heated to 100 °C for 10 min to inactivate any protease and DNase which may have been present. Then the crude broth culture was cooled to 0 °C for 10 min and 5 µl portions were submitted to PCR, as described below.

Primers and probe

The computer analysis of the *S. typhimurium mkfA* gene [21–24] and the *S. choleraesuis* virulence plasmid *mba* region [25] allowed us to find a primers set and an internal probe, after sequence alignment. Primers (VIR 113, VIR 561) and the internal probe (VIR 334) were chosen according to the following criteria: no cross-hybridization with other known sequences, high G+C content at 3' extremity, minimum of dimer formation and self-complementarity. The primers set generates a 472 bp amplified DNA fragment. Positions and sequences of the oligonucleotides are shown in Table 2.

PCR conditions and electrophoresis

Bacterial DNA was amplified with a PTC-100 Programmable Thermal Controller (MJ Research Inc., Watertown, MA, USA). *Taq* polymerase (2 units, Boehringer Mannheim) was added to 50 µl of a solution of Tris-HCl pH 8.3 (10 mM), KCl (50 mM), MgCl₂ (1.5 mM), gelatin (0.1 mg/ml), deoxynucleotide

Table 1. Ability of PCR assay to distinguish the virulence-associated gene among salmonella serovars and other bacterial strains

Strains	Origins	Results
<i>abortus ovis</i>	Ovine abortion product	+
<i>abortus ovis</i>	Ovine abortion product	—
<i>agona</i>	Breeding environment	—
<i>agona</i>	Guinea fowl viscera	—
<i>alachua</i>	fish meal	—
<i>anatum</i>	Pork butchery	—
<i>banana</i>	Foodstuffs	—
<i>blockley</i>	Hen viscera	—
<i>bovis morbificans</i>	Bovine faecal sample	—
<i>bovis morbificans</i>	Pork butchery	—
<i>bovis morbificans</i>	Sea sediment	—
<i>braenderup</i> (× 2)	Bovine faecal samples	—
<i>braenderup</i> (× 2)	Breeding environment	—
<i>braenderup</i>	Chicken meat	—
<i>braenderup</i>	Dog faecal sample	—
<i>braenderup</i>	Surface water	—
<i>braenderup</i>	Raw milk	—
<i>bredeney</i>	Pork butchery	—
<i>bredeney</i>	Sea water	—
<i>bredeney</i>	Sea sediment	—
<i>broughton</i>	Beef meat	—
<i>cerro</i>	Dog faecal sample	—
<i>choleraesuis</i>	Young wild boar	+
<i>cubana</i>	Foodstuffs	—
<i>derby</i>	Calf offals	—
<i>derby</i>	Pork butchery	—
<i>derby</i>	Sea sediment	—
<i>dublin</i>	Beef meat	—
<i>dublin</i> (× 2)	Bovine abortion product	+
<i>dublin</i> (× 2)	Bovine faecal sample	+
<i>eboko</i>	Bovine abortion product	—
<i>enteritidis</i>	Bovine faecal sample	+
<i>enteritidis</i> (× 3)	Breeding environment	+
<i>enteritidis</i>	Cacao	+
<i>enteritidis</i>	Cooked food	+
<i>enteritidis</i> (× 2)	Duck viscera	+
<i>enteritidis</i> (× 2)	Egg product	+
<i>enteritidis</i>	Hen viscera	—
<i>enteritidis</i> (× 2)	Hen viscera	+
<i>enteritidis</i>	Pastry	+
<i>enteritidis</i>	Pastry	—
<i>enteritidis</i>	Raw milk	+
<i>enteritidis</i>	River water	+
<i>enteritidis</i>	Turkey meat	+
<i>gallinarum/pullorum</i> (× 4)	Hen viscera	+
<i>gloucester</i>	Turnsol meal	—
<i>goldcoast</i>	Beef meat	—
<i>hadar</i>	Chicken meat	—
<i>heidelberg</i>	Hen viscera	—
<i>hessarek</i>	Surface water	+
<i>idikan</i>	Fish meal	—
<i>indiana</i>	Beef meat	—
<i>infantis</i>	Beef meat	—

Table 1 (*cont.*)

Strains	Origins	Results
<i>infantis</i>	River water	—
<i>isangi</i>	Shellfish	—
<i>kedougou</i>	Hen viscera	—
<i>kottbus</i>	Breeding environment	—
<i>llandoff</i>	Breeding environment	—
<i>lille</i>	Foodstuffs	—
<i>london</i>	Pork butchery	—
<i>mbandaka</i>	Soya meal	—
<i>meleagridis</i>	Beef meat	—
<i>montevideo</i>	Fish meal	—
<i>montevideo</i>	Hen viscera	—
<i>münchen</i>	Processing plant	—
<i>münster</i>	Horse meat	—
<i>newport</i>	Pastry	—
<i>newport</i>	Turkey viscera	—
<i>nima</i>	Poultry faecal sample	—
<i>panama</i>	Beef meat	—
<i>paratyphi B</i>	Beef meat	—
<i>paratyphi B</i>	Bovine viscera	—
<i>paratyphi B</i>	Fish	—
<i>paratyphi B</i>	Ice cream	—
<i>paratyphi B</i>	Treatment plant	—
<i>reading</i>	Turkey faecal sample	—
<i>regent</i>	Breeding environment	—
<i>rissen</i>	Turnsol meal	—
<i>rubislaw</i>	Fish viscera	—
<i>saintpaul</i> (× 2)	Poultry meat	—
<i>sandiego</i>	Turkey viscera	—
<i>schwarzengrund</i>	Hen viscera	—
<i>senftenberg</i>	Hen viscera	—
<i>senftenberg</i>	Manioc	—
<i>stanley</i>	Bovine faecal sample	—
<i>tennessee</i>	Beef meat	—
<i>thompson</i>	Poultry meat	—
<i>typhimurium</i>	Bovine faecal sample	+
<i>typhimurium</i> (× 4)	Breeding environment	+
<i>typhimurium</i>	Chicken meat	+
<i>typhimurium</i>	Dog faecal sample	+
<i>typhimurium</i> (× 2)	Duck viscera	+
<i>typhimurium</i>	Egg product	+
<i>typhimurium</i>	Foodstuffs	+
<i>typhimurium</i>	Foodstuffs	—
<i>typhimurium</i>	Human faecal sample	+
<i>typhimurium</i> (× 2)	Pork butchery	+
<i>typhimurium</i>	Pork butchery	—
<i>typhimurium</i>	Pork meat	+
<i>typhimurium</i>	Poultry meat	+
<i>typhimurium</i>	River water	+
<i>typhimurium</i>	Treatment plant	+
<i>virchow</i>	Hen viscera	—
<i>virchow</i>	Poultry meat	—
<i>wien</i>	Pork butchery	—
<i>worthington</i>	Pork butchery	—
<i>zanzibar</i>	Turkey viscera	—
<i>S. I 4</i> :-:-	Quail viscera	—
<i>S. I 6,7</i> :-:-	Processing Plant	—

Table 1 (cont.)

Strains	Origins	Results
<i>S. I</i> 9,12:-:-	Bovine viscera	—
<i>S. I</i> 1,3,19:-:-	Hen viscera	—
<i>S. I</i> 1,3,19:i:-	Fish meal	—
<i>S. I</i> 1,3,19:z ₂₇	Colza meal	—
<i>S. I</i> 1,3,19:z ₂₇	Foodstuffs	—
<i>S. II</i> 42:b:enz ₁₅	Frog leg	—
<i>S. IIIa</i> 48:z ₄ , z ₂₃	Foodstuffs	+
<i>S. IIIb</i> 38:r:z	Treatment plant	—
<i>S. IIIb</i> 61:i:z ₅₃	Equine viscera	—
<i>S. IIIb</i> 61:i:z ₅₃	Pork butchery	—
<i>S. IIIb</i> 61:k:1,5,7	Ovine viscera	—
<i>S. IV</i> 43:z ₄ , z ₂₃ :-	Wild boar meat	+
<i>S. V</i> 28:a:-	Gelatine	+
<i>S. rough</i>	Pork butchery	—
<i>S. rough</i> (× 2)	Sea water	—
<i>Aeromonas</i> sp. (× 2)	Estuary water	—
<i>Citrobacter freundii</i>	Poultry breeding	—
<i>Citrobacter freundii</i>	Poultry meat	—
<i>Citrobacter freundii</i>	Surface water	—
<i>Enterobacter aerogenes</i>	Cooked food	—
<i>Enterobacter cloacae</i>	Estuary water	—
<i>Enterobacter cloacae</i>	Poultry breeding	—
<i>Enterobacter hafniae</i>	Bovine faecal sample	—
<i>Enterobacter hafniae</i>	Estuary water	—
<i>Escherichia coli</i>	Bovine breeding	—
<i>Escherichia coli</i> (× 2)	Estuary water	—
<i>Escherichia coli</i> (× 2)	Poultry breeding	—
<i>Klebsiella pneumoniae</i>	Poultry breeding	—
<i>Moraxella</i> sp.	Estuary water	—
<i>Listeria innocua</i>	Cheese	—
<i>Listeria ivanovii</i>	Cheese	—
<i>Listeria monocytogenes</i>	Cheese	—
<i>Proteus mirabilis</i>	Horse meat	—
<i>Proteus morgani</i>	Pig offals	—
<i>Proteus rettgeri</i>	Bovine viscera	—
<i>Providencia</i> sp.	Estuary water	—
<i>Providencia</i> sp.	Poultry breeding	—
<i>Pseudomonas</i> sp. (× 4)	Estuary water	—
<i>Sarcina lutea</i>	Bovine breeding	—
<i>Shigella sonnei</i>	Slaughterhouse	—
<i>Staphylococcus aureus</i>	Cheese	—
<i>Staphylococcus epidermidis</i>	Milk	—
<i>Vibrio vulnificus</i>	Estuary water	—
<i>Yersinia enterocolitica</i>	Poultry breeding	—

(200 µM each, Boehringer), primers (0.5 µM each) and target DNA. The reaction mixture was heated for 5 min at 95 °C and overlaid with mineral oil to prevent evaporation. PCR reaction was performed with 30 cycles as follows: 5 sec primer annealing at 60 °C, +1 °C/3 sec up to 95 °C, 5 sec denaturation at 95 °C, -1 °C/sec down to 60 °C. After the 30th cycle, the extension reaction was continued for another 10 min at 72 °C. Samples of the reaction mixture (10 µl) were loaded on to a 2% agarose gel containing ethidium bromide (0.5 µl/ml) for electrophoresis. After 30 min under 100 V, the gel was examined under u.v. light

Table 2. *Positions and sequences of oligonucleotides*

Oligonucleotides	Sequences (5'-3')	Corresponding position in virulence plasmid gene [22]
VIR 113	TTGTAGCTGCTTATGATGGGGCGG	113-136
VIR 561	TGGAGAAACGACGCACTGTACTGC	561-583*
VIR 334	CGAGAATCACCTCAGTCTCAGGGC	334-357

* Complementary DNA strand.

(312 nm) and the amplification product size was compared to DNA fragments size of a molecular weight marker (Marker VI, Boehringer). Negative controls containing all reagents except template DNA were performed in each amplification set. To avoid contamination, sample preparation, PCR amplification and electrophoresis were performed in three different rooms.

Probe labelling

The probe labelling was performed with the 'DNA Tailing Kit' (Boehringer). According to the manufacturer's instructions, tailing buffer, CoCl_2 , dTTP, digoxigenin-11-dUTP, sterile distilled water, terminal desoxynucleotidyl transferase and the oligonucleotide to be labelled (100 p.m.) were mixed and incubated for 15 min at 37 °C. To end the reaction, a stop solution (Glycogen, EDTA), was added. The labelled probe was purified on a G 50 Sephadex column (Pharmacia, Uppsala, Sweden) with a Tris-HCl pH 8 (10 mM), EDTA (1 mM) elution buffer.

Southern blot and hybridization

After agarose gel electrophoresis, DNA denaturation was performed for 15 min with NaOH (0.4 N), followed by neutralization with Tris-HCl pH 7.2 (0.5 M), NaCl (1.5 M) for 45 min to 1 h 30 min. Then amplified DNA fragments were transferred to a positively charged nylon membrane (Boehringer) using saline sodium citrate buffer (SSC, 20 ×) in a vacuum blotter (10 mm mercury pressure) for 45 min at room temperature, and fixed on to the membrane by heating at 120 °C for 30 min. The filter was pre-hybridized in SSC (5 ×) buffer, *N*-laurylsarcosine (0.1%), sodium dodecyl sulphate (SDS, 0.02%) and blocking reagent (1%, Boehringer) at 65 °C for 1 h. For hybridization, labelled probe VIR 334 (25 ng/ml) was added and the filter was incubated at 55 °C for 2 h 30 min. After SSC (2 ×), SDS (0.1%) washing, the filter was then washed to a final stringency of 55 °C in SSC (0.1 ×), SDS (0.1%). Detection was performed with alkaline phosphate-labelled anti-digoxigenin antibody (Boehringer), nitro blue tetrazolium (NBT) and bromo-4-chloro-3-indolyl phosphate (BCIP) according to the manufacturer's instructions (Boehringer).

RESULTS

Optimization of the amplification conditions

In this study it was not necessary to perform three thermal steps. A very good amplification yield was obtained by using only two temperatures with a ramping time of +1 °C/3 sec between annealing and denaturation temperatures. This allowed reduction of the reaction time in comparison with standard PCR. The global reaction time was thus reduced to 1 h 30 min, with results equivalent to

Fig. 1. Salmonella PCR results. Lanes: 1, *S. typhimurium*; 2, *S. enteritidis*; 3, *S. hessarek*; 4, *S. choleraesuis*; 5, *S. dublin*; 6, molecular weight marker VI (Boehringer); 7, *S. gallinarum/pullorum*; 8, *S. abortus ovis*; 9, *S. IIIa* 48:z₄, z₂₃; 10, *S. IV* 43:z₄, z₂₃:-; 11, *S. V* 28:a:-. (Numbers on figure are molecular weights in base pair.)

those obtained using conventional amplifications, which incorporate a temperature hold time during each step (data not shown).

PCR/DNA hybridization assay specificity

After PCR assay, no cross-reaction was observed with the 35 non-salmonella strains derived from 15 species (Table 1).

No amplification products, specific or not, could be observed for 94 salmonella strains from 66 serovars (Table 1), irrespective of the source under consideration.

An amplification product of expected molecular size (Fig. 1) has been obtained for 49 salmonella strains, from 10 serovars (Table 1). The 10 serovars' amplification products have all hybridized with the digoxigenin-labelled probe VIR 334 (Fig. 2).

DISCUSSION

We have developed a PCR assay for the detection of salmonella virulence-associated plasmid gene. A primers set, derived from the virulence gene, allowed us to detect 10 serovars among 71 tested serovars. No amplification product was observed with 35 strains from 15 other bacterial species, and an internal probe ensured the specificity of the salmonella amplification products using a DNA/DNA hybridization test. Thus the 472 bp amplification product revealed the presence of a plasmid gene linked to virulence in salmonella.

As shown in Table 3, the strains responding to amplification were of different origins: human, 1; animal pathology, 19; food hygiene, 18; environment, 11; illustrating that strains with increased virulence are present in all biotopes.

Fig. 2. Vir 334 probe hybridization results, after Southern blot of the electrophoresis gel shown in Fig. 1. Lanes: 1, *S. typhimurium*; 2, *S. enteritidis*; 3, *S. hessarek*; 4, *S. choleraesuis*; 5, *S. dublin*; 6, molecular weight marker VI (Boehringer); 7, *S. gallinarum pullorum*; 8, *S. abortus ovis*; 9, *S. IIIa* 48:z₄, z₂₃; 10, *S. IV* 43:z₄, z₂₃:-; 11, *S. V* 28:a:-. (Numbers on figure are molecular weights in base pair.)

Seven of the ten serovars producing an amplified DNA fragment were mentioned in the literature as possessing virulence-related plasmids: *S. abortus ovis* [7, 17], *S. choleraesuis* [3, 16, 19, 25], *S. dublin* [12, 16–18], *S. enteritidis* [4, 8, 15, 26], *S. gallinarum/pullorum* [2, 3, 11], *S. hessarek* [18], *S. typhimurium* [5, 14, 19, 21]. The three other serovars were *Salmonella* subspecies IIIa, IV and V, which were encountered less frequently (Table 1).

Two of the 18 *S. enteritidis* strains studied showed no reaction (Table 1), proving that for a given serovar known to contain virulence-related plasmids [4, 8, 26–27], there is diversity. This was also confirmed for *S. abortus ovis*, since only one of the two tested strains possessed the plasmid gene associated with virulence (Table 1) and for *S. typhimurium*, since only 2 of the 20 tested strains did not respond to amplification (Table 1). Moreover, one serovar (two strains), described as possessing virulence-associated plasmid, i.e. *S. newport* [3, 17], did not produce a positive response when analysed by PCR during this study.

Several authors [2, 3, 17, 18, 28–29] have described serovars containing plasmids of various molecular size which did not contain the genes associated with virulence. The search for the plasmid content of strains was often used in the study of epidemiological markers [8, 29, 30–32]. While remaining of some interest, this approach does not provide any information as to the potential virulence of strains, since on the one hand the presence of plasmids does not necessarily imply the presence of the plasmid gene associated with mouse virulence [15, 17, 28], and on the other hand the plasmid/virulence combination is not inevitably linked to the molecular weights of the plasmids observed [4, 28, 32].

These facts illustrate the advantages of searching for genetic factors associated with virulence. The PCR technique used is perfectly suited to this search and has

Table 3. PCR salmonella tested strains distribution according to the origin

Origins	Tested strains number (serovars number)	PCR positive reaction strains number and involved serovars (number)
Human pathology	1	1 <i>typhimurium</i>
Animal pathology		
Bovine	13 (9)	6 <i>dublin</i> (4), <i>enteritidis</i> , <i>typhimurium</i>
Dog	3 (3)	1 <i>typhimurium</i>
Equine	1	0
Fish	2 (2)	0
Ovine	3 (2)	1 <i>abortus ovis</i>
Poultry	26 (18)	10 <i>enteritidis</i> (4), <i>gallinarum/pullorum</i> (4), <i>typhimurium</i> (2)
Young wild boar	1	1 <i>choleraesuis</i>
Food hygiene		
Beef meat	10 (10)	1 <i>dublin</i>
Cacao	1	1 <i>enteritidis</i>
Colza meal	1	0
Cooked food	1	1 <i>enteritidis</i>
Egg product	3 (2)	3 <i>enteritidis</i> (2), <i>typhimurium</i>
Feedstuffs	7 (6)	2 <i>typhimurium</i> , <i>S. IIIa</i> 48:z ₄ , z ₂₃
Fish meal	4 (4)	0
Frog leg	1	0
Gelatine	1	1 <i>S. V</i> 28:a:-
Horse meat	1	0
Ice cream	1	0
Manioc	1	0
Milk	2 (2)	1 <i>enteritidis</i>
Pastry	3 (2)	1 <i>enteritidis</i>
Pork butchery	12 (10)	2 <i>typhimurium</i>
Pork meat	1	1 <i>typhimurium</i>
Poultry meat	9 (7)	3 <i>enteritidis</i> , <i>typhimurium</i> (2)
Shellfish	1	0
Soya meal	1	0
Turnsol meal	2 (2)	0
Wild boar meat	1	1 <i>S. IV</i> 43:z ₄ , z ₂₃ :-
Breeding environment	29 (17)	11 <i>enteritidis</i> (4), <i>hessarek</i> , <i>typhimurium</i> (6)

the advantage of being quick, reliable and easy to use, in comparison with conventional techniques. The standard methods used to investigate the relationship between the presence of a plasmid and increased virulence call for preliminary extraction of the plasmids followed by the study, in mice, of the effect of strains with or without their plasmids, or the use of DNA/DNA hybridization techniques [7, 9, 10, 11, 13, 16, 21, 26]. In comparison with the PCR assay, these techniques are time-consuming and difficult to apply in a systematic way when investigating the presence of these genes.

The ubiquitous presence of the strains (Table 3), the diversity of the serovars involved and the preservation of plasmid genes associated with virulence during evolution [33] merit a proper place for this research among the ranks of epidemiological markers currently in use. The PCR method proposed gives the information sought directly and quickly, either by comparing strains from the same biotope or by the knowledge of the strains' capacity to develop in a given organism [6, 10, 12-13].

All these findings show that the search for plasmid genes associated with mouse virulence is of proven epidemiological interest and the method described above is perfectly suited to this task, in terms of its ease of use in comparison to tests on mice or DNA/DNA hybridization techniques, its reliability, reproducibility, rapidity (amplification; 1 h 30 min; detection, 30 min) and the possibility of investigating a large number of strains.

ACKNOWLEDGEMENTS

The authors thank J. P. Guillou (CNEVA/LCRV) for his invaluable help in the computer analysis for choosing the primers set and the probe. They are grateful also to S. Fremy, M. Goncalves, C. Oudart, C. Piquet and C. Pires Gomes (CNEVA/LCHA) for biochemical and serological analysis of salmonella isolates.

REFERENCES

1. Johnston WS, Hopkins GF, Lachlan GK, Sharp JCM. *Salmonella* in sewage effluent and the relationship to animal and human disease in the north of Scotland. *Vet Rec* 1986; **119**: 201-3.
2. Barrow PA, Simpson, JM, Lovell MA, Binns NM. Contribution of *Salmonella gallinarum* large plasmid towards virulence in fowl typhoid. *Infect Immun* 1986; **55**: 388-92.
3. Barrow PA, Lovell MA. The association between a large molecular mass plasmid and virulence in a strain of *Salmonella pullorum*. *J Gen Microbiol* 1988; **134**: 2307-16.
4. Chart H, Threlfall EJ, Rowe B. Virulence of *Salmonella enteritidis* phage type 4 is related to possession of a 38 MDa plasmid. *FEMS Microbiol Lett* 1989; **58**: 299-304.
5. Gulig PA, Curtis III R. Plasmid-associated virulence of *Salmonella typhimurium*. *Infect Immun* 1987; **55**: 2891-901.
6. Jones GW, Rabert DK, Svinaritch DM, Whitfield HJ. Association of adhesive, invasive and virulent phenotypes of *Salmonella typhimurium* with autonomous 60 MDa plasmids. *Infect Immun* 1982; **38**: 476-86.
7. Nakamura M, Sato S, Ohya T, Suzuki S, Ikeda T. Possible relationship of a 36 MDa *Salmonella enteritidis* plasmid to virulence in mice. *Infect Immun* 1985; **47**: 831-3.
8. Shah DB, Bradshaw JG, Peeler JT. Thermal resistance of egg-associated epidemic strains of *Salmonella enteritidis*. *J Food Sci* 1991; **56**: 391-2.
9. Terakado N, Sekizaki T, Hashimoto K, Naitoh S. Correlation between the presence of a 50 MDa plasmid in *Salmonella dublin* and virulence for mice. *Infect Immun* 1983; **41**: 443-4.
10. Hackett J, Kotlarski I, Mathan V, Francki K, Rowley D. The colonization of Peyer's patches by a strain of *Salmonella typhimurium* cured of the cryptic plasmid. *J Infect Dis* 1986; **53**: 1119-25.
11. Barrow PA, Lovell MA. Functional homology of virulence plasmid in *Salmonella pullorum*, *Salmonella gallinarum* and *Salmonella typhimurium*. *Am Soc Microbiol* 1989; **57**: 3136-41.
12. Chikami GK, Fierer J, Guiney D. Plasmid-mediated virulence in *Salmonella dublin*, demonstrated by use of Tn5-oriT construct. *Infect Immun* 1985; **85**: 420-1.
13. Pardon P, Popoff MY, Coynault C, Marly J, Miras I. Virulence associated plasmids of *Salmonella ser. typhimurium* in experimental murine infection. *Ann Inst Pasteur Microbiol* 1986; **137B**: 47-60.
14. Pohl P., Lintermans P, Bex F, Desmyter A, Dreze P, Fonteyne PA, Couturier M. Propriétés phénotypiques et génotypiques de 4 plasmides de virulence de *Salmonella typhimurium*. *Ann Inst Pasteur Microbiol* 1987; **138**: 523-8.
15. Chart H, Threlfall EJ, Rowe B. Virulence studies of *Salmonella enteritidis* phage types. *Lett Applied Microbiol* 1991; **12**: 188-91.
16. Beninger PR, Chikami G, Tanabe K, Roudier C, Fierer J, Guiney D. Physical and genetic mapping of the *Salmonella dublin* virulence plasmid psdL2. Relationship to plasmids from other *Salmonella* strains. *J Clin Invest* 1988; **81**: 1341-7.

17. Popoff MY, Miras I, Coynault C, Lasselin C, Pardon P. Molecular relationship between virulence plasmids of *Salmonella* ser. *typhimurium* and *dublin* and large plasmids of other *Salmonella* serotypes. *Ann Inst Pasteur Microbiol* 1984; **135A**: 385–98.
18. Woodward MJ, McLaren I, Wray C. Distribution of virulence plasmids within *Salmonella*. *J Gen Microbiol* 1989; **135**: 503–11.
19. Fierer J, Chikami G, Hatlen L, Heffernan EJ, Guiney D. Active immunization with LD342, a plasmid-cured strain of *Salmonella dublin*, protects mice against group D and B *Salmonella* infection. *J Infect Dis* 1988; **158**: 460–3.
20. Nakamura M, Sato S, Ohya T, Suzuki S, Ikeda T. Plasmid-cured *Salmonella enteritidis* AL1192 as a candidate for a live vaccine. *Infect Immun* 1985; **50**: 586–7.
21. Norel F, Coynault C, Miras I, Hermant D, Popoff MY. Cloning and expression of plasmid DNA sequences involved in *Salmonella* ser. *typhimurium* virulence. *Mol Microbiol* 1989; **3**: 733–43.
22. Norel F, Pisano MR, Nicoli J, Popoff MY. Nucleotide sequences of the plasmid-borne virulence gene *mftA*, encoding a 28 kDa polypeptide of *Salmonella typhimurium*. *Res Microbiol* 1989; **140**: 263–5.
23. Norel F, Pisano MR, Nicoli J, Popoff MY. Nucleotide sequence of the plasmid-borne virulence gene *mkfB*, from *Salmonella typhimurium*. *Res Microbiol* 1989; **140**: 455–7.
24. Norel F, Pisano MR, Nicoli J, Popoff MY. A plasmid-borne virulence region (2.8 Kb) from *Salmonella typhimurium* contains two open reading frames. *Res Microbiol* 1989; **140**: 627–30.
25. Matsui H, Kawahara K, Terakado N, Danbara H. Nucleotide sequence of a gene encoding a 29 kDa polypeptide in *mba* region of the virulence plasmid pKDSC50 of *Salmonella choleraesuis*. *Nucl Acids Res* 1990; **18**: 1055.
26. Helmuth R, Stephan R, Bunge C, Hoog B, Steinbeck A, Bulling E. Epidemiology of virulence associated plasmids and other membrane protein patterns within 7 common *Salmonella* serotypes. *Infect Immun* 1985; **48**: 175–82.
27. Hovi M, Sukupolvi S, Edwards MF, Rhen M. Plasmid-associated virulence of *Salmonella enteritidis*. *Microbiol Pathog* 1988; **4**: 385–91.
28. Clarke RC, Jorgensen ST, Harnett NM, Gyles CL. Plasmids of *Salmonella muenster* and their relation to virulence. *Can J Vet Res* 1987; **51**: 436–9.
29. Nastasi A, Mammina C, Villafrate MR, Massenti MF, Scarlata G, Diquattro M. Multiple typing of strains of *Salmonella enterica* subsp. *bongori* 48:z35:- isolated in southern Italy. *Ann Inst Pasteur Microbiol* 1988; **139**: 605–12.
30. Farrar WE. Molecular analysis of plasmids in epidemiologic investigation. *J Infect Dis* 1983; **148**: 1–6.
31. Martel JL, Corbion B, Carbonnelle B, Voisin E. A propos d'un foyer de salmonellose humaine et bovine à *S. typhimurium*. Une anthroozoonose par contact direct. *Bull Acad Vét France* 1990; **63**: 415–23.
32. Wray C, McLaren I, Parkinson NM, Beedell Y. Differentiation of *Salmonella typhimurium* DT 204c by plasmid profile and biotyping. *Vet Rec* 1987; **121**: 514–16.
33. Jones CS, Osborne DJ. Identification of contemporary plasmid virulence gene in ancestral isolates of *Salmonella enteritidis* and *typhimurium*. *FEMS Microbiol Lett* 1991; **80**: 7–12.