in him we have a careful observer and a conscientious worker, and one whose methods we will do well to imitate.

Before I conclude I would wish it to be understood that there is another side to this picture, or perhaps it may be the same picture looked at from a different standpoint. It is that much harm may be done by too excessive attention to rest. Evil may result from too prolonged rest. Mechanical rest may, in one sense, be antagonistic to physiological rest. Mechanical rest, in many cases, must be interfered with in order to attain physiological rest. An example will best show my meaning. Immediately after an injury the effusions into the tissues may, by their presence, interfere with the normal blood current through the part. At a later date these effusions are replaced by organised material which will also act in the same way. The nerve equilibrium will also be altered. The part will then be, from the vascular and nervous side, in a state of physiological unrest, and this unrest will be intensified by prolonged mechanical rest, because, unless there is a normal blood current, the effusions and fibrous material will not be removed. It is therefore necessary that, while we maintain mechanical rest after a part is injured, we should at the same time adopt some means to remove these products. It is here that massage is so valuable; lightly applied it has a marked soothing influence on the nerve disturbance; more strongly, though still gently applied, it will get rid of the effusions by causing a temporary congestion and free flow of blood through the part; still more strongly applied, it breaks down fibrous adhesions and gets rid of the pain felt in certain movements of the limb. While the massage interferes with the mechanical rest, it acts directly in relieving the physiological unrest. Experience alone will tell how far we can go with massage in order to attain the one object—the physiological rest—while at the same time we avoid doing harm by its over-use by interfering with the mechanical rest. In acute sprains and strains it may be begun at once, gently night and morning, using elastic pressure with wadding and a flannel bandage in the intervals of the massage. In sub-acute cases it may be used more freely, wearing an elastic bandage in the intervals, along with limited use of the injured limb. In chronic cases, which are non-tuberculous, adhesions may be freely broken down, often giving immediate relief after months of partial impairment of usefulness.

I am also strongly of opinion that in fractures near joints, as in Colles's and Pott's fracture, massage may with advantage be begun within a week, with the result that while the repair of the broken bone is in no way interfered with (I rather think it is aided), the limb is a useful one at a much earlier period than is the case if, as in the orthodox treatment, the limb is kept absolutely quiet for three or four weeks. If we think only of the broken bone and forget the injury to the surrounding soft parts, the result is a stiff and useless limb, which will for a long time be a source of discomfort and helplessness to the individual. It is a question exercising my mind whether we should not apply gentle massage in all fractures, as a matter of routine practice, so long as we can do so without displacing or causing movement between the broken fragments of the bone. The use of extension during the massage applied to the limb beyond renders this method much more feasible than it formerly was when we depended entirely on splints applied at the seat of fracture commanding the joints above and below. It is interesting to note in this connection that no fractures heal more kindly and quickly than broken ribs, in which it may truly be said that during the whole process of cure the act of breathing is keeping up a constant gentle movement, a nature's massage, which in no way interferes with the union of the broken bone.

In breaking down adhesions in old standing cases of fracture, sprain, or strain, one must act in a decided manner. Their presence is associated with limited movement, pain on movement, or pain on pressure, and the use of firmly applied rotatory massage, or the sudden stretching of the tissues which are matted together, often gives immediate and lasting relief.

In the case of nerve stretching in sciatica, the cases which are benefited are, in my opinion, those which may be called trade sciaticas, due to some special position adopted in the special trade pressing on and irritating the sciatic nerve.

You freely stretch the nerve, but do not interfere in any way with the sensory and motor functions, and the pain is relieved by breaking down the fibrous adhesions in the nerve sheath and among the nerve fibrils. May I say in passing that the operation is sometimes a source of psychical unrest to the operating surgeon if he does not easily find the nerve? This unrest is avoided if, in operating, the patient lying on his face, the surgeon will stand on the opposite side to the limb to be operated upon. If he then makes an incision over the nerve at the lower border of the gluteus maximus large enough to enable him to introduce his forefinger, which, using as a hook, he draws towards the middle line of the patient, he will at once find the nerve lying external to the muscles arising from the tuber ischii.

New lamps may have been expected of me to-day; if so, my hearers have been disappointed. "Let us make a stand on the ancient ways, and then look about us and discover what is the right and straight way, and so walk in it." Bacon was fond of quoting this passage, and it has been my motto. I have taken my stand on an ancient way; I have tried to polish and refill an old lamp.

James Hinton, another of my heroes, in one of his letters, writes: "Let me advise just once. I don't like an adviser much; but just this one thing—be reverent where you are ignorant, and attach no weight at all to your naturally feeling sure. We almost always feel sure wrongly—it is our own fate, it is our very being." The speaker to-day may be too sure, and may place too much reliance on rest as the most powerful therapeutic agent in surgical practice, but he can assure you that he has had Hinton's words constantly in his mind as he spoke, and what he has said is offered to this audience in the same spirit in which they were written by that philosophic surgeon. I began with a sentence from Hinton; I end with one from Hinton. These men had something in common. Hinton taught rest; Hinton sought it. In one of his last letters he writes these sad words: "I have tried for too much, and failed; but yet, perhaps, in that my failure God is giving me more than even I tried for. He has opened my eyes, at least a little, though I am blind and foolish still, no doubt. I will try and be wiser and look more, and care more what others feel." Strange words from one who spent his whole life for the good of others.

At a time fertile in unrest in religion, politics, and surgery, in the county in which Gilbert White spent his days, in the county in which he wrote one of the most restful—I had almost said *the* most restful—book I know, rest as a thesis is perhaps not altogether out of place, especially when I remember that to many of us this meeting is our annual holiday, our resting stage, and still more especially when I remember that we are enjoying the generous hospitality of the inhabitants of one of the main resting places in this country, where so many get that rest which enables them to go back to work with energies renewed and restored by the fresh air and restfulness of one of the most attractive rest resorts in Great Britain.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF OBSTETRICS.

At the Annual Meeting of the British Medical Association, held in Bournemouth, July, 1891.

By WILLIAM J. SMYLY, M.D.,

Master of the Rotunda Hospital, Dublin.

GENTLEMEN,—My first pleasing duty is to thank the Council for the honour they have conferred upon me in appointing me President of this Section. At the same time, I do not flatter myself that this selection has been made because of any peculiar fitness on my part to fill such an important post, but rather as a token of esteem for that great institution over which I have the good fortune to preside. The reputation of the Rotunda Hospital is due, I believe, not so much to its being the oldest and largest institution of its kind in these countries, but to the number of pupils which it attracts and sends forth annually to all parts of the world. The institution consists of four departments, and

the numbers annually treated in each are about as follows :—

Delivered in the Lying-in Hospital ...	...	...	1,200
Attended in their own homes ...	...	...	1,800
Admitted to Gynecological Hospital ...	...	...	450
Out-patient Department ...	...	...	8,000

A good teaching institution must, in my opinion, comprise these four departments. Midwifery and gynecology must go together; they are sciences which God has joined together and should never be put asunder. In this I am quite aware that I differ from some who have occupied this chair before me, who have expressed a hope that this Section may soon split into two—a wish which I most earnestly hope may never be realised. I would especially refer to the opening address of my immediate and much respected predecessor, Dr. Savage, of Birmingham, and to the arguments and illustrations which he used in support of his views—which to my mind, so far from proving the necessity for such a separation, most conclusively show that gynecology and obstetrics are inseparable, and that the obstetrician must also be a skilled gynecologist.

In the first place, he says: "An obstetrician, having failed to extract the foetus *per vaginam*, might hesitate to send for his friend, expert in abdominal operations, to perform Cæsarean section." But Cæsarean section should be performed in the first instance, after carefully weighing the indications for it, and not after unsuccessful attempts at extraction, since in the latter case the risk to mother and child would be enormously increased.

Some months ago a woman came to the Rotunda Hospital with a kyphotic pelvis, the tuberosities of the ischia being separated by an interval of only 2 inches; her former labour had been terminated by craniotomy, and she was very anxious to have a living child. I did not think that induction of premature labour would give any prospect of a living child, so I advised waiting until term and then Cæsarean section. To this she readily assented, so six weeks ago I performed the operation with a favourable result to mother and child. An exact diagnosis and careful measurement of the pelvis are essential preliminaries to scientific treatment, and these are the duty of the obstetrician and not of the pure gynecologist. For accurate pelvimetry I most strongly commend to you Skutsch's pelvimeter, which I now show, and which is the only instrument by means of which accurate measurements can be obtained.

He (Dr. Savage) next cited a case of puerperal fever, and asked who should be called in as a consultant—a general physician, a gynecologist, or an obstetrician? Well, I should certainly say the man who combines the greatest amount of skill with the widest familiarity with the disease in question, that is the man who practises both midwifery and gynecology. "Difficuly," he admits, "might arise in drawing the line" that is between midwifery and gynecology. I would go further and declare it an impossibility.

Let us now endeavour to define this line. A woman presents herself for examination. The first duty of the examiner is to determine the presence or absence of pregnancy—that is obstetrical; but if she happen to be sterile in consequence of endometritis she should, of course, be treated by a gynecologist. Under his judicious treatment she so far improves as to become pregnant; provided the pregnancy be uterine her case is obstetrical, otherwise gynecological. Even if the ovum be situated in the uterus her position is still uncertain; for if the conjugate diameter of her pelvic brim be 2½ inches or less she should go to the gynecologist, otherwise to the obstetrician. Her pregnancy probably ends in abortion, which does not improve the endometritis, and she acquires the habit of aborting, and again requires the gynecologist: becoming pregnant again she goes to term, but has placenta prævia. If this be partial the obstetrician is in place, but if complete abdominal section is, according to Mr. Tait, advisable, and so a specialist in this department is called in; but after delivery by the obstetrician she may be attacked with septic peritonitis, when, according to Dr. Savage, the gynecologist is again required. Should her perineum be ruptured the obstetrician may at once sew it up, but if primary union be not obtained she is once more handed over to the gynecologist. But, gentlemen, such a course would not only be absurd and contrary to the dictates of common sense: it would frequently be fraught with danger.

A few weeks ago my assistant, Dr. Bagot, was called to a patient in the extern maternity. In a tenement house not far from the hospital he found a poor woman, almost collapsed from severe accidental hæmorrhage, which had been at first concealed. The pulse was 148, very small and compressible; her lips were quite blanched, and her pupils dilated. There was jactitation and sighing respiration. The uterus was larger than the term of pregnancy, though the membranes had been ruptured before his arrival, in order to check the bleeding. The child presented in the first position, vertex; no foetal heart could be heard. The os, which was rigid and undilatable, admitted one finger only. On pushing up the head the blood flowed freely out of the uterus. Labour had not set in, though she had all night suffered from severe distension and pains in the uterus. Believing that owing to her collapsed state it would be impossible to deliver her alive by perforation followed either by version or extraction with the cranioclast, he at once performed Porro's operation, treating the pedicle extraperitoneally by means of a *serre-nœud* and pedicle pins made out of two Peaslee's perineum needles, as he had no time to procure proper pins. The placenta was found at the operation to be completely detached, and the uterus was full of clots; the child of course was dead. She was removed to the Rotunda Hospital on the fourth day after the operation, and has made an excellent recovery, though still very anæmic. Great credit is due to Dr. Bagot for his pluck and promptness in performing an operation under such disadvantageous circumstances, but credit is also due to Mr. Lawson Tait, who had pointed out to us this method of saving life in *ante-partum* hæmorrhages, which would otherwise prove fatal.

The arbitrary division between midwifery and gynecology is peculiar to this country, and is due to their separation in special hospitals, so that the gynecologist has come to despise midwifery, and the obstetric physician is often a poor operator. In his introductory address at Cardiff, the late Dr. Meadows, in recommending this separation, stated that the recent advances, especially in abdominal surgery, were due to the pure gynecologists, a statement, in my opinion, altogether contrary to fact. I do not wish for one moment to ignore the valuable and original work done by Mr. Lawson Tait and others in this country, but if we take gynecology as a whole and include our Continental brethren, the obstetricians can show quite as good gynecological work as the pure gynecologists.

I would not deny to any man the right to limit his practice as he may see fit; he may treat piles only, or fistula, or cancer, or may confine his attention to abdominal surgery, but what I do maintain is that, in teaching, gynecology must not be separated from midwifery, and their isolation places such difficulties in the way of students, that practical midwifery is not efficiently taught, and the study of gynecology is more or less optional. Clinical teaching can be efficiently carried out only in the wards of a hospital, where too a limited material can be utilised in teaching a number of pupils. In extern maternities each student requires a number of cases, and it is therefore not to be wondered at that we hear so much of the lack of cases for the instruction of medical students as well as midwives. This appears to me to be the true difficulty in the midwives question; they must first be taught before they can be registered. I do not deny that there are excellent lying-in hospitals in this country, but they are altogether insufficient in number, and their value is diminished by the want of a gynecological department. In order to appreciate the value of good midwifery, the student must also be shown the evil results of bad midwifery, and that he can only learn in a gynecological hospital.

Another important matter is accurate clinical observation, and this can only be carried out in hospitals. It was from observing the comparative mortality in two lying-in hospitals that Semmelweis first got on the track which led to his discovering the cause of puerperal fever, and pointed out its proper prophylaxis. I am a firm believer in the doctrines of that great observer. I believe that puerperal fever is a preventable disease, and therefore a greater responsibility rests upon me in conducting a great lying-in hospital.

Let me now in conclusion illustrate this by giving you an account of a septic outbreak which we experienced in the Rotunda Hospital during the spring and summer of last year,

and of the measures taken to eradicate the disease. When I was appointed to the hospital on November 1st, 1889, there had not been one death from any septic cause for 18 months, that is, out of 2,000 deliveries. These results were so brilliant that I thought it inadvisable to make any changes in the methods employed for the prevention of septic infection. I will briefly state what the routine practice then was. A patient was permitted to be examined abdominally by any number of students, but vaginally by three only and one pupil midwife. Previous to examination the external genitals were carefully washed with soap and water, irrigated with plain water, and finally bathed with corrosive sublimate solution, 1 in 500. The examiners' hands were thoroughly cleansed with soap and water and a good nailbrush, the soap removed by irrigation with carbolic lotion, and the hands then washed in the sublimate solution. Prophylactic douching was only employed in cases of purulent or septic discharges and where operation was required. Douching after delivery was employed only when specially indicated, for example, in hæmorrhage, after the introduction of the hand or instruments, after the birth of a putrid fœtus, for putrid or purulent discharge, and in fever. For some months prior to my appointment plain water alone was used for uterine douching, and though personally I was in favour of the use of antiseptics in such cases yet with such good results I felt that the experiment might be continued. This favourable state of affairs continued up to January, 1890, when a patient, sent up from the country for induction of premature labour, died of acute septicæmia. It was exceedingly difficult in this case to excite uterine action, the catheter had to be inserted several times, and on one occasion caused some hæmorrhage, having evidently caused a partial separation of the placenta. Hot douches, Barnes's bags, and rupture of the membranes had to be resorted to in order to procure delivery. Acute septicæmia set in the next day with a measly rash, and she died within a week. The health of the other patients during this month was excellent; in one other case only did the temperature reach 101°F.

In February there were 73 deliveries, and in 7 the temperature reached 101°. In March there were 108 deliveries, with 13 morbid cases in which the temperature reached 101°. This great increase in the morbidity alarmed me, and I exhorted the pupils to increased care in disinfection, but still the morbidity increased, twelve cases of high temperature—101° and over—occurring in the first fortnight. I therefore stopped all vaginal examinations, and all trouble ceased. It may seem strange to those who place reliance for information upon vaginal exploration only to hear that in a great institution where deliveries average 100 a month, we would give up this method of examination; but by abdominal palpation all necessary information can be obtained, excepting only as regards prolapse of the funis and the condition of the os uteri; but the frequent examination of the fetal heart minimises the danger of the former case, and the latter is of quite secondary importance.

In April vaginal examinations were resumed, and there were 12 morbid cases out of 96 deliveries.

In May 119 women were delivered, with 24 morbid cases and 1 death. In that case fever set in on the fifth day, and she died on the fifteenth. On looking over the bed cards I found that one gentleman who had examined her had also examined two other patients the same day, both of whom became dangerously ill, but finally recovered. I accordingly prohibited his attending the hospital for one month, and again forbade vaginal examinations, excepting only in particular cases when they were made by myself or one of my assistants.

In June vaginal examinations were again permitted, but only by one student, my object being to increase the sense of personal responsibility. During this month 106 women were delivered. There were 21 morbid cases, but all excepting one were unimportant; that one died on the fourteenth day. We now returned to antiseptic douching, but in July, out of 110 cases, there were 14 morbid, most of them severe, and 1 death. This patient had a normal delivery, and was not examined vaginally. Her convalescence was normal up to the fifth day, when her temperature rose to 102°. Her uterus was washed out with carbolic acid lotion, 1 in 40, and some putrid clots came away. She was again douched out on two successive days, but pyæmic symptoms set in, and she died at the end of

a week. Towards the end of August there were seven cases of severe illness, and I again resorted to the plan which I had previously found so successful, namely, stopping vaginal examinations altogether, but on this occasion without the former good results. Two cases proved fatal. On September 2nd another woman was confined, and ultimately died. The first of these three cases had been examined by one student only. Her temperature continued normal up to the fifth day, when it rose to 104.4°, with headache and shivering. The uterus was washed out, and the temperature sank to normal, and continued so until the eighth day, when she left the hospital, having signed a declaration that she did so contrary to our wishes. The same evening she had a rigor, and her temperature rose to 106°. A week later she was readmitted to hospital with pyæmia, of which she died six weeks from the date of delivery.

The second case was not examined vaginally. Her temperature continued normal up to the fifth day, when it rose to 100.4° F. Septic symptoms set in with diarrhœa, and she died on the fifteenth day.

The last case was delivered on September 2nd. Her labour was normal, and she was not examined vaginally. Her temperature continued normal until the fourth day, when she also became septic and died on the twenty-sixth day.

In considering these cases the following points were important: One only had been examined vaginally, and in all infection was late, and was due apparently to some error in the management of childbed rather than of the labours. The method adopted up to this time during convalescence was introduced by my predecessor, Dr. Macan. Each patient was given a basin night and morning containing water, and a large piece of tenax, and was directed to wash herself, the object being to prevent the carrying of infection from one patient to another by the nurse. It struck me that this method was faulty because it would be impossible for an ignorant woman lying in bed thoroughly to disinfect her fingers, and I therefore directed that the probationer who attended a patient during her delivery should continue to do so afterwards, washing her carefully twice a day, and using the same antiseptic precautions as if she were making a vaginal examination. Each patient was also to have her basin placed over her bed, and to have it carefully disinfected with corrosive sublimate, both before and after using it; they had previously been kept on a dresser, and washed with warm water. Since that date we have had upwards of 1,000 deliveries in the hospital, but not one death from any septic cause, nor even one case that gave us anxiety.

I hope that I have not wearied you with these minute details. I have entered into them to show that puerperal fever can be prevented, and that the cause is generally something so obvious that when it is discovered we are surprised at our own blindness in having overlooked it. It is generally held that septic infection is uncommon in private practice, but this I believe to be a dangerous mistake, and the same precautions are as absolutely necessary in private as in hospitals.

If in these remarks I have been able to show the inadvisability of separating our Section into two, and the advantages that would result from the union of obstetric and gynecological hospitals, my purpose has been accomplished.

STROUD URBAN AND RURAL (Population, 9,725 and 28,485). BISLEY (Population, 5,331).—These districts, which are under the supervision of Dr. T. Partridge, enjoyed comparative immunity from epidemic diseases during 1890; but Dr. Partridge's reports contain abundant evidence of good and useful sanitary work. Scarlet fever became somewhat troublesome in the rural districts, and necessitated the closing of schools in two villages, but no deaths were caused by this disease. The chief factors in the zymotic death-rates were diarrhœa and whooping cough. Special attention was given to the surroundings of each case of diphtheria and typhoid fever, most of them being found in connection with defective sanitary arrangements. One or two cases of the former disease were, doubtless, the result of contagion. The death-rates from all causes were as follows: Stroud Urban District, 16.2 per 1,000; Stroud Rural District, 14.7 per 1,000; Bisley, 15.0 per 1,000.