as infringing the sacred principle of local self-government. As things are at present, the arrangements for the isolation of infectious diseases in many parts of the country can only be described as an administrative chaos. Many towns of considerable size are still without proper accommodation of any kind for such cases, while in others such provision as has been made is utterly inadequate. Moreover, owing to the division of authority and the absence of efficient control by local centres, administrative anomalies—in the highest degree absurd, if they were not deplorable—have arisen. This is explained by the fact that the rural sanitary authorities have, in some instances, erected hospitals for their own use, while the towns in the district have not shown the same foresight. Thus we find hospitals situated near to large populations, the most important sections of which are excluded from the use of them. In this way it has happened that a hospital admirably adapted for the reception of infectious cases has stood empty when almost at its doors were patients of the very kind it was designed to accommodate, who, owing to the intricate machinery of local self-government which is a part of our existing constitution, could not be admitted within its walls. I need not dwell on the inconvenience and injury to individuals and the dangers to the community at large that may arise from administrative pedantry of this kind. The transference of the power of regulating the provision of isolation hospital accommodation to the county councils is the only means of remedying the state of things which I have here attempted to describe, and of saving the unnecessary suffering to patients and the expense to ratepayers entailed thereby.

Lines of Future Evolution.—With regard to the future evolution of sanitary administration, there is not much that can be profitably said at the present moment. The important questions now floating in the legislative atmosphere as to the advisability of transferring vaccination powers to the county councils, and as to the formation of sufficiently large areas for administrative purposes, must be left for solution till the opinions of men of light and leading on these subjects have had time to condense into practical conviction, and to infiltrate the minds of sanitary authorities. It may be said in general terms, however, that it is desirable that future evolution should be in the direction of greater simplicity in organisation, and a nearer approach to uniformity in the size of sanitary districts and in the methods of administration than exists at present. Here let me remind you that the question of "whole service" or other restrictions must obviously depend on considerations of administrative areas. The absurdity of considering them apart, as Parliament has done this session, is best illustrated by taking an extreme case, one of the tiniest and most impracticable areas. For example, Ham, near Richmond, in Surrey—the "umbrageous Ham" of the author of the *Seasons*—is a separate sanitary administrative area, with its own sanitary staff. The population is only 1,400, that of a mere hamlet. Now, if there could be found a medical officer willing to sacrifice his "whole time" and actually to reside on the spot, there would still be difficulty as to the emolument. This amounts to £21 per annum! Of course the remedy for this is the creation of workable districts for sanitary purposes within counties, and until that has been done it is premature to adopt the whole-service formula. I would not, however, be understood as advocating a cut and dry system, in which the country should be parcelled out into trim parterres like a magnified Dutch garden, governed by authorities acting with the rigid uniformity of mechanical puppets. There is something very distasteful to the English mind in anything that savours of centralisation or the restraint of administrative individualism by cast-iron rules; but, on the other hand, I think, most thoughtful sanitarians agree in considering that the present freedom of local authorities from all control is sometimes rather apt to degenerate into administrative licence, with serious consequences to those immediately affected thereby, and to the community at large.

In conclusion, while thanking you for your kind attention to these somewhat desultory remarks, I would express a hope that no one will think the subject of this address unimportant, though it is too likely that I may have failed to make it attractive. The details I have touched on are the framework of a vast living structure, all the parts of which should work together for the common good. The perfecting of this struc-

ture, the clothing of the dry bones of sanitary administration with the flesh of scientific knowledge and well-directed practical action, should be the aim of all legislators and authorities, and of all practitioners of curative, not less than of preventive, medicine.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF SURGERY.

At the Annual Meeting of the British Medical Association, held in Bournemouth, July, 1891.

By JOHN WARD COUSINS, M.B.LOND., F.R.C.S.,

Senior Surgeon to the Royal Portsmouth Hospital, and Portsmouth and South Hants Eye and Ear Infirmary.

RECENT ADVANCES IN THE TREATMENT OF TUBERCULOUS DISEASES OF THE JOINTS.

Historical Notes.—During the past thirty years very remarkable development has taken place in all departments of our art, and although I think the expansion of our knowledge in connection with diseases of the joints has been less brilliant than in many other directions, still it has not been less radical as regards the system of treatment. Until the time of the late Sir Benjamin Brodie confusion and uncertainty surrounded this class of disorders, but by the labours of this distinguished surgeon many of these diseases were disentangled from each other, and the affections of the articulating apparatus, which had been grouped together under the common designation of "white swelling," were clinically separated. In the fourth edition of his work on *Diseases of the Joints*, published in the year 1836, he graphically described tuberculous disease of the bones as commencing by the deposition of a transparent material in the cancelli, and afterwards undergoing transformation into a yellow cheesy substance. He drew a correct picture of the slow progress of the inflammation, the caries of the bone, the recurrent centres of suppuration, and the final termination either in imperfect ankylosis or complete disorganisation of the joint, associated too often with pulmonary disease or some other visceral affection.

Brodie and Liston investigated the morbid alterations in articular cartilage and synovial membrane. They regarded the cartilages as prone to primary alterations of structure, and they initiated the doctrine that suppuration was a rare result of their primary ulceration, and that it took place only in the advanced stages of the disease, with caries of bone and destructive changes in the synovial membrane. They regarded the presence of capillaries as an essential condition of the inflammatory process, and they laboured to demonstrate the vascularity of cartilage in disease. The pathological doubts and difficulties of these eminent surgeons occupied the attention of many of their followers, and their clinical observations were marked by clearness and accuracy, but their minds were prevented from forming correct interpretations of their facts by their strange prejudices and imperfect views of the healthy and morbid processes.

Some of the problems surrounding this structure were, however, at length solved by the labours of Redfern, Goodsir, and Rainey. These observers clearly demonstrated the changes which took place in cartilage by the perverted activity of disease. They asserted that it remained entirely non-vascular, and that its disorganisation was always accompanied by changes in the size and form of the corpuscles, and the softening and breaking up of the intercellular substance. During the process of repair they discovered that fibrous tissue had formed, and that into this new substance vessels were projected from the vascular system of the bone and synovial membrane, and thus the difficulties concerning the vascularity of inflamed cartilage were cleared away.

The Old Pathology of Tubercle.—The loss of substance in articular cartilage long occupied a prominent place in the

pathology of joint disease, and observations on the wear and tear of this texture in the joints of men and animals were utilised in support of a mechanical theory of incipient disease. At length, however, these narrow doctrines languished under the expanding power of broader views. The belief soon became general that joint disease could have an origin in any of the joint structures, and that most frequently the error commenced in the bone or the synovial membrane. In a large proportion of chronic affections the morbid changes were found to be the results of a peculiar inflammation, attended by the deposition of a semi-transparent exudation, which had a tendency to unhealthy suppuration, and to undergo fatty transformation. Pathologists diligently studied the scrofulous affections of bone, and for a long period of years the essential element in the process was described as a deposit of tubercle—a formation of low type which sooner or later excited destructive changes in the surrounding structures. In one case a synovitis issued in a gelatinous degeneration which ultimately extended to the bone and cartilage; in another case, the deposition of tubercle in the cancellous structure caused an expansion of the bone which insidiously progressed towards softening and caseous changes, inflammation of the fibrous capsule of the joint, and hopeless destruction by suppuration and caries.

The Scrofulous Diathesis and Tubercle.—Now, the universally accepted theory of all these changes was the unfortunate possession of a bodily or constitutional condition which was styled the strumous cachexia. This was explained as an altered direction of the normal nutrition of the system, which could make itself visible on slight provocation in any of the structures or organs of the body. Scrofula was regarded as an inherited or acquired constitutional condition, which might at any moment assume the appearance of a distinct tuberculous disease. On the other hand, tuberculosis was always manifested by a wider departure from normal nutrition, and presented new and lowly-organised formations which were deposited somewhere in the organism in the course of chronic strumous inflammation. Virchow described scrofula as a disease of the lymphatic system, and tubercle as a new product derived from the lymphatic elements of connective tissue.

Universally, then, tuberculosis was defined as an inheritance very variable in individuals; at the same time, outside influences were recognised as powerful factors in kindling the smouldering flame into activity. For many years the nature of tubercle received profound investigation, and on every hand the results of research tended to establish the time-honoured theory. The microscope failed to detect in it a specific element. All the histological constituents were in turn considered the essential ingredient in the deposit. Cells and granules, epithelial elements, and cells of giant form, were searched for some characteristic quality. Some of them appeared embedded in a fine homogeneous stroma, while others were either shrivelled or drowned in particles of oil, or else buried in molecular matter. The morbid changes in bone were searched over and over again, but in the process of degeneration and caseation only negative elements were discovered. In one part the osseous structure was observed undergoing destruction, in another the osteoblasts were at work forming masses of protecting bone. With one voice the microscopists declare that they could find no specific element, and that tubercle was so indefinite in structure that it could be recognised by negative rather than by positive characters.

Now these were the opinions which for more than half a century were confidently taught in all the schools. Surely the experience of our day is sufficient to convince us that there is no part of our pathology which may not be transformed, and no theory which may not be pulverised by progress. These doctrines were considered orthodox for many years, and by the profound investigations of many eminent men all difficulty appeared to have passed away. Tubercle had been examined by the best observers in the civilised world; thousands of clever eyes had gazed at it with intense persistency and curiosity, and with a remarkable unanimity they pronounced the opinion that it really contained nothing but cells.

The Modern Pathology.—In the year 1882 the whole pathology of tuberculosis underwent a great evolution by the complete demonstration of the life-history of the tubercle

bacillus. The disease occurring in any tissue of an organism must now be regarded as a specific disorder, the bacilli as the direct cause of all the morbid changes, and their presence as the distinctive sign of the disease.

Now it is only by drawing a comparison between the old and the new pathology that we can really decipher the magnitude of the revolution in our conceptions. We no longer fight about the primary seat of the disease, for it is now certain that it may commence in any of the tissues of a joint. At the onset it may be synovial or osseous, and when the origin is in the latter structure it may select its surface or its substance. Strumous disease of the bones and joints is the same disease as tuberculous disease of the bones and joints; for in all these affections a specific deposit can be detected. Tubercle bacilli are never found but in this special product, and wherever the tuberculous tissue is discovered this microbe has obtained a resting-place. Their number may vary in the diseases of different structures, and also at different periods of the same disorder; still, when only a few are present they will be found by careful scrutiny.

The new pathology of tubercle is sustained by a mass of evidence derived from microscopic research, the artificial cultivation of the parasite, and experimental inoculation, so that the old notions which had long surrounded the disease have been numbered with the things of the past. The long-accepted causes, too, have been dislodged from their position, and are rightly grouped as morbid tendencies. The inheritance of constitutional peculiarities, the liability to chronic inflammations, and the susceptibility to external influences are thus regarded as essential conditions which help the microbe to establish itself within the body. Still, these factors are not less potent because the specific character of tuberculosis has been recognised. Hereditary proclivities and physical peculiarities of structure have not fallen into insignificance, but in their new position it will be possible to better estimate their potency.

Hereditary Transmission.—As regards the old doctrine of hereditary transmission, must we abandon it altogether? How is it possible for the disease to be conveyed from the parent to the embryo unless the spores of the bacilli are themselves actually transmitted? The living particles have in themselves no penetrating power, still their vitality is certainly intensified by contact with living tissues. As a general rule they make an entrance into the body by the mucous surfaces, and then their diffusion is secure through the ever flowing streams of blood and lymph. We know that they find their way into the bones and joints, and there appears nothing likely to prevent them getting entangled with equal ease in the placental structures. I believe, however, that there is at least some evidence in support of the assertion that the microbe can pass freely from the mother to the foetus.¹ The great discovery of Koch may have around it many problems yet to be unravelled. It is, however, a fact of history that by his profound research he brought to light from the microscopic elements of tubercle a living atom which no human eye had seen before. Some have been so generous as to call it a lucky hit, but let us remember that the accident happened to a man of untiring energy and prodigious power of mental concentration, and these are qualities as essential for success in science as the full activity of the higher intellectual forces. I regard this power of mental concentration as the noblest element of genius. Do not the histories of men illustrious in science exhibit a passion amid their solitary labours? Every line of eternal truth that has been added to our knowledge, every new fact that has been brought to light from the deep secrets of the universe, every noble success in the onward march of science, every triumph achieved over the mysteries of the natural world are the offspring of unceasing devotion.

Arrest Possible in the Early Stage.—The modern pathology of tuberculosis has already exercised a salutary influence over surgical treatment and the progress of conservative surgery. Its distinct recognition as a specific and infectious disorder, in all its various manifestations, has placed the hope of arrest on a very different basis to that which it previously occupied under the old constitutional theory. During the early stages of joint disease the morbid action is often localised, and,

¹ Birch-Hirschfeld and Schmöll, *Beiträge zur path. Anat. und zur allg. Path.*, 1891, p. 429.

therefore, arrest is at least possible. There can be no reason why a joint or a bone should not recover, and the tuberculous infiltration atrophy, and ultimately shrivel into a fibrous scar just like a similar deposit in the apex of a lung. If the number of bacilli found by microscopic examination in a part are to be taken as any indication of the activity of the disease, then repeated observations are favourable to the conclusion that incipient disease in the articulating apparatus is more hopefully situated than incipient disease in the pulmonary tissue.

It is true that the early arrest of tuberculosis has long been the result aimed at by treatment. Half a century ago Brodie described the slowness of the process. The cure in the advanced stages, he said, began when the sinuses closed and the oedema of the limb subsided, but the morbid changes in the joint generally terminated in more or less ankylosis caused by the destruction of the articulating surfaces. The diseased limb should be kept in a state of perfect quietude, not that this alone would restore the bones to a healthy condition, but it would do much to prevent the inflammation extending to other structures. He recommended constitutional treatment, residence at the seaside, nutritious diet, and exposure to fresh air.

We now restrict the term tuberculosis to those local and general changes which are directly caused by the irritation of a specific bacillus. Under favourable conditions isolated patches of tuberculous formation may undergo atrophy, and the infiltration around them cease to extend. This arrest is the result of the timely death of the micro-parasite, for surely by its destruction alone the morbid action in the tissues can be localised, and the individual delivered from the danger of a diffused tuberculosis. What, then, constitutes the grave difference between this curative process and the insidious progress of the disorder? Does it depend upon the strength of the inoculation and the number and vitality of the bacilli; or is it due to the intensity and persistency of the predisposing conditions or other factors in the causation which may be vital, or chemical, or structural peculiarities of blood and tissue? In the light of new facts I think we must admit that the case is really dependent upon the complete destruction of the specific cause within the affected structures. At the same time, experience teaches us that arrest is possible, and that this fortunate issue may be obtained by the healthful influence of sunlight, pure air, and good food upon the whole organism. I wish some philanthropic millionaire, for the sake of humanity and science, would try the experiment of submitting five hundred scrofulous children living in the slums of our large centres, and suffering from incipient joint disease, to the renovating power of good food in combination with seaside purity and brightness for at least eighteen months, and I am sure some of them would by these potent remedies, and the application of a simple splint, escape from more serious surgical treatment.

Value of Expectant Measures.—In the management of joint disease we must take care to select the right moment for surgical interference. It is often an anxious question to decide when expectant measures ought to be abandoned. In the early stages the diagnosis may be open to question. Sometimes the threatening symptoms may be traced to a traumatic cause. There may be nothing in the local condition to indicate its specific character, and the general conformation and nutrition may not point to any predisposition to tuberculous inflammation. The affected joint may be swollen and occasionally painful and tender. Its outline may be altered and the normal movements impaired, and these slight indications may be marked by a persistent tendency to recurrence. In some cases the morbid action appears to have subsided, and the joint structures to have regained their healthy condition, but this favourable quiescence may be disturbed by the slightest injury, and any trivial accident may rouse again the dormant malady. Occasionally we obtain evidence of arrest many years after the favourable issue has occurred. I have seen several cases in which slight shortening of a limb has not been recognised until puberty, the period of life when the skeleton is in a condition of active development. The osseous deficiency is the result of latent mischief near the growing line, and the remnant of bygone epiphyseal inflammation which happened during early life.

The late Hugh Owen Thomas.—A few years ago strumous

children labouring under chronic joint disease were kept in bed for many months, but now expectant treatment can be carried out more hopefully by combining rest and protection of the limb with fresh air and exercise. The management of incipient joint disease has fortunately undergone a silent revolution through the mechanical genius of the late Hugh Owen Thomas, of Liverpool. His surgical appliances are admirably adapted for taking off concussion, arresting friction, and imparting support and protection without pressure; at the same time they are so simple in construction that the patient can adjust them without assistance. Thomas's splints have been utilised by surgeons in all parts of the world, and the name of our old colleague will long have an honourable place in the surgical records of our times.

Tuberculin.—When, a few months since, the celebrated announcement reached us that a new remedy had been discovered which possessed the remarkable power of causing the necrosis of living tuberculous tissue, a new method of arrest appeared probable in recent cases, and material improvement in others of greater severity which would prepare them for surgical treatment. It is not my intention, however, to invade the arena of my distinguished colleague, Mr. Watson Cheyne, who has promised to-day to tell us the results of his elaborate investigations with tuberculin, but I feel bound to mention that although I have used the remedy in many cases of joint and bone disease without decided benefit, it will still receive from me a full and impartial trial. It is certain that it contains an agent which is capable, even in almost an infinitesimal dose, of exciting active changes in the body containing any traces of active tuberculosis. In the chemical aspects of microbial life, and in the complex bodies which are formed by the artificial cultivation of the bacilli themselves, we discern the direction from which we may anticipate future discoveries. I regard the search for a remedy amid the growth of these living particles as a splendid effort to reduce the magnitude of a world-wide pestilence, for which, up to the present moment, no really scientific treatment has ever been propounded. The great German investigator has not yet finished his work; let us patiently wait for his results, and keep ourselves free from prejudice, hoping that his daring assault upon the most deadly of diseases may ultimately be crowned with success.

Surgical Treatment of the Past.—The recognition of the local character of tuberculosis in diseases of the joints has opened up the high road to many of the recent advances in surgical treatment. Sixty years ago these disorders were described by Sir Benjamin Brodie as having their origin in the cancellous structure of the bones, or as a consequence of inflammation extending from the synovial membrane to the osseous tissue. Morbid action, he said, commenced sometimes in one and sometimes in another texture, and in the advanced stages all the structures of the joint became involved in the disease. At that time, however, the surgical treatment was really little else than splints and plasters, incisions and punctures, and at length amputation as a final remedy. The constitutional theory of tuberculosis blocked the road of progress.

Fergusson and Butcher.—Between the years 1830 and 1840, Syme in Scotland, and Liston in London, revived with energy the resection of joints for disease—an operation which had been occasionally performed by British surgeons half a century before. But it was not until Sir William Fergusson had successfully practised it that excision became an established manipulation on all the joints, and professional prejudices were vanquished. Only a few months since, Richard Butcher, of Dublin, passed away after a long and brilliant career. He laboured, too, in his day to resuscitate the practice of excision, and devised the well known saw especially for joint operations, and this ingenious instrument has certainly assisted the progress of conservative surgery.

Early Efforts.—Until the last few years early operations in joint disease had scarcely received any attention. In 1878, Volkmann, the inventor of the cutting spoon, performed several partial excisions in recent cases of hip-joint disease, but his results did not stimulate him to further trials. Since the year of the pathological revolution (1882) many favourable cases have been recorded, and I feel confident the practice, although still regarded by some as experimental, has already saved many limbs from graver operations.

Advantages of Early Operation.—Whenever the indications

for surgical interference are clear, early operation must be attended with many advantages. The risk of delay is always in proportion to the progressive and obstinate character of the disease, and timely aid will often prevent its extension, and at the same time deliver the patient from the danger of deep infection and the development of secondary tuberculous centres. The only hope of cure must depend upon the complete removal of the diseased tissue, and the facility with which this can be accomplished rests entirely upon the extent of the local mischief. The preservation, too, of useful mobility in the joint may be anticipated when the morbid process is well localised, so that the manipulation involves only a limited excision of synovial membrane and a partial division of the fibrous capsule.

Partial Arthrectomy.—For the successful performance of a partial arthrectomy there must be clear evidence of a localised deposit. In some cases children have exhibited very little pain or lameness, but the joint has been in some part swollen, with the capsule thickened and the bones enlarged, but without any indication of softening or suppuration. By a well directed operation, near the neck of the femur or the head of the tibia, search has been made for a spot of tuberculous infiltration, with the result that a carious cavity has been found and small sequestra successfully removed. Surely these are examples of the enormous gain obtained by early operation.² If the latent disease had not been detected and cut out, the issue must have been irreparable injury of the articulation in every instance, and a more serious manipulation under conditions far less favourable.

Another recommendation for early arthrectomy is the little danger that attends the operation. By ordinary surgical precaution the risk may be fairly considered trifling, even when a portion of bone has to be resected. After a full incision in the most convenient position for exploration and carefully defining the disease, the infiltrated tissue must be excised with the scissors or cutting spoon, and the cavity thoroughly flushed with hot water. To insure rapid union, the surface should be then dried, and the wound closed with deep and superficial sutures. The limb must be kept at rest until the healing process is complete. Up to the present time a great many partial arthrectomies have been performed in this country by different surgeons, but I am not prepared to state the exact proportion of their permanent successes; and it appears to me that a report of the results of early operations from many hospitals would prove at the present time a very valuable record.

Advancing Local Tuberculosis.—The articular cartilages are seldom the seat of primary disease, for as a general rule the morbid process has its origin either in the synovial structure or the articular extremity of the bone. When the osseous tissue is the seat of a tuberculous infiltration the evidence of its existence is often wanting until softening occurs within it, and inflammatory reaction takes place around it. As soon as these infective changes reach the synovial membrane they extend to all the structures of the joint. Sometimes they make their way through the superficial cancelli to the outer layer of the bone, and then superficial caries and slow suppuration are the result. At another time the morbid process advances in the direction of the articular cartilage, softening and erosion of this structure follow, and then inflammatory changes which issue, unless checked by surgical treatment, in chronic abscess, caries of bone, imperfect arrest of the disease, and finally ankylosis. Now in all these forms of advancing tuberculous disease, surgery offers the only scientific method of treatment, and we can safely repeat our incisions, scoopings, scrapings, and cleanings, until the disorder is eradicated and a useful joint preserved.

Sudden Infection of the Joint.—But instead of a slow disorganisation, the tuberculous centre may be suddenly discharged into the capsule, diffusing the infective material over the whole synovial surface, and kindling suppurative inflammation with great rapidity. A few months since I performed arthrectomy on a child for acute infection of the knee-joint. In 1889 the little patient was under my care, in the Royal Portsmouth Hospital, labouring under a small sub-periosteal abscess over the head of the tibia, close to the reflection of the synovial membrane. The swelling was freely incised,

and a considerable deposit of caseous material cleaned out with the spoon. The bone was roughened perilously near the articular edge, and the parents were specially warned of the danger. Soon after the child left the hospital the swelling slowly recurred without either pain or lameness. In the month of July last she hurt her knee during a game of play. Acute pain immediately followed the accident, attended with rapid swelling of the joint, and fever. Three days after she was readmitted to the hospital under my care. The knee was at once freely opened on both sides, and the inner incision was carried through the abscess cavity over the head of the tibia. The capsule contained about three ounces of a turbulent fluid with many flakes. The infiltrated synovial membrane was freely excised with scissors, the joint thoroughly cleansed, and the limb carefully placed on a back splint. Irrigation was continued for a week. The child was discharged quite well in January. The movements of the knee are now normal, a result due to immediate treatment and the limited injury of the synovial membrane. (Figs. 1 and 2.)

FIG. 1.

FIG. 2.

Complete Arthrectomy.—Permit me now to offer a few remarks on the surgical treatment of more advanced cases in which the morbid process is too extensive for any partial operation. In the performance of complete arthrectomy a free division of the ligaments and capsule is necessary for the exploration of all the recesses of the articulation and the excision of deep infiltrations of the synovial and osseous structures, so that the preservation of only a limited mobility must be anticipated. The whole of the pulpy granulation tissue must be dissected off, and the ligaments and cartilages carefully scraped. It is absolutely necessary to remove every particle of the diseased synovial membrane, and all tuberculous foci in the bones must also be cleanly cut out with the gouge. Care must be taken to prevent any remnants of the infective tissue being left behind on the raw surfaces, and the accidental reinoculation of the disease through the medium of the fresh incisions. I regard the method advocated by Mr. Arthur Barker, of flushing the seat of operation with hot water, to be the best way of carrying out these important precautions, and for the rapid performance of this part of the operation his ingenious scoop and irrigator will be found of great practical utility. The operation of arthrectomy of the hip can be readily performed by the anterior and vertical incision and division of the neck of the femur with the saw, and then the excision of the infiltrated tissues. After a simple protective treatment for a few weeks, and the application of a Thomas's splint during convalescence, the results are often very satisfactory.

Surgical Treatment in the Advanced Stage of Hip Disease.—With reference to the old method of operating in the advanced stages of the disease, after suppuration has been proceeding for months and sinuses have long riddled the soft parts, and when the unfortunate patient has been exhausted by a general tuberculosis, I sincerely hope it will soon be cast into the shade for ever by the light of modern progress. It is my experience that these distressing cases recover more frequently by simple measures, consisting of free incision, scooping, irrigation, and drainage, and that the ultimate results are more satisfactory than those which follow the practice of severe and dangerous operations.

Complete Arthrectomy of the Knee-joint.—In performing complete arthrectomy of the knee-joint I consider the old horse-shoe incision, and an oblique division of the ligamentum patellæ, better than any other method for obtaining free access to the interior of the cavity. The practice of lifting up the tuberosity of the tibia instead of division of the tendon may be found very useful in some cases. It is important to clean the lateral and crucial ligaments carefully, and to avoid damaging the cartilages and articulating surfaces.

Successful cases reported in the BRITISH MEDICAL JOURNAL by Messrs. Watson Cheyne, A. Barker, and Charters Symonds.

Sometimes carious bone can be removed with the gouge outside the capsule of the joint. It must always be our object to secure the complete extirpation of the diseased structures, and to preserve as far as possible the mobility of the articulation. Fortunately the limb is not shortened, and the development of the bones is scarcely impaired. The articulating surfaces are in a great measure preserved, and the operation, when compared with resection of the joint, is attended with less risk of life. After arthrectomy I have seen a sound and useful limb with movement to the extent of 45° , and the patient capable of standing and walking many hours every day without fatigue. So good a result can only be anticipated under favourable conditions and early surgical interference. I have received from my colleagues in different places very varying reports of their successes. Sometimes they claim firm ankylosis, but at other times their operations have issued in weak limbs and joints, tending to serious flexion and displacement. Many of them were, no doubt, too late arthrectomies, and the careful excision of the bony surfaces would have been followed by firmer and better results.

The Old Operation of Excision.—With reference to my own practice, I prefer at once complete resection in all cases marked with old sinuses and the carious remnants of old infiltrations. My method is to freely open the joint and to remove as thin a layer of bone as possible from all the articulating surfaces. The gouge is used for cleaning out any infiltrations, and then the patella is divided vertically with the saw. After cleaning every recess in the capsule, the bones are jammed together as tightly as possible, the periosteum is carefully sutured, and the wound closed, except at the extreme ends of the incision. Now, I do not presume for a moment that my results are exceptionally good, but I submit sketches from photographs of some of my patients upon whom operations were performed many years ago (Figs. 3, 4,

FIG. 3

FIG. 4

FIG. 5

FIG. 6

FIG. 7

5, 6, and 7). I have often questioned them with reference to the inconvenience of the ankylosis, and, without exception, they have regarded it with indifference. One patient volunteered the remark that the only annoyance her stiff limb caused her arose from the necessity, when sitting in company, of crossing her legs.

Conservative Surgery in the Advanced Stages of Tuberculous Disease.—I am no advocate for any attempt at conservative surgery in the advanced stages of the disease, not from the dread of deepening the general and local infection from re-inoculation through the seat of operation, but rather from the feebleness of the vital power which has been slowly undermined by prolonged suffering and suppuration. The intensity of the disease has been quickened by a chronic septic condition of the system, and it is the danger of its sudden aggravation, even with every possible precaution, which adds to the risk of surgical interference. The micro-organisms of septicæmia, or their poisonous products, exert a marked influence over the course of tuberculosis, for their presence reduces the resisting power of the system, and helps on the local and constitutional spread of the disease.

But it is not only septic infection in its various forms that we have to combat, but every other kind of acute dis-

order of the blood. Children especially are liable to be attacked with any of the infectious fevers: and these are all attended with peculiar risks. Measles, from its disturbing influence over nutrition, is prominent among the group for its power of rekindling tuberculous inflammation.

Occasionally, however, I think that my experience has seemed to point in the opposite direction, and that an accidental blood storm has exercised a remarkable effect on the course of the malady. A young woman, who for many years had suffered from old disease of the knee, came under my care in consequence of acute inflammation of the joint, which she attributed to injury. Prior to this event she had undergone a good deal of surgical interference. Some years since I did a plastic operation for her, and in 1887 I performed abdominal section for pelvic suppuration and chronic peritonitis. A few days after her admission to the hospital her distress was so great and the constitutional disorder so acute, that I decided to amputate the leg. She was, however, I think very fortunately attacked with severe erysipelas of the head and face, and after a few weeks' dangerous illness her recovery all round was so remarkable, and the joint symptoms so much reduced, that I did an excision. The wound healed with rapidity, and she is now earning her livelihood on two legs instead of one.

Now, notwithstanding the opinion I have expressed concerning the risk of late operations, exceptions occasionally occur in practice when interference appears to be the better course. There are certainly cases of chronic tuberculous joint disease marked by secondary centres and general infection, and even aggravated by a chronic septic condition of the system, in which the vitality of the tissues and the residue of constitutional vigour appear sufficient to warrant an effort in the direction of conservative surgery.

The accompanying drawings (Figs. 8 and 9) are two examples. Fig. 8 shows the back and side of the elbow of a

FIG. 8

FIG. 9

FIG.

girl taken a few months after excision of the joint. At the time of the operation she was labouring under partially arrested hip disease with still one discharging sinus, and tuberculous deposit in many parts of the glandular system. She is now able to follow any occupation. Fig. 9 represents the elbow of a young woman, an orphan, with strong tuberculous history, upon whom I did a similar operation on a joint riddled with sinuses and surrounded with infiltrations. She had distinct physical indications of pulmonary complication, old disease of the opposite knee, and a persistently elevated temperature. She has now recovered with a useful arm, and is enjoying apparently good health. I am quite aware that these may prove only temporary successes, but it is our duty to look always on the hopeful side. I am, however, quite certain that their future prospects have been improved by surgical interference.

In conclusion, I have endeavoured to indicate some of the improvements which have taken place in the treatment of joint affections during the last few years, and to trace this progress to the influence of recent advances in pathology. Our science is destined to progress: before us there are great possibilities. It is true that to-day the pathway of scientific enterprise is still rugged and thorny, but obstruction is often a kindly foe, and prejudices are the secret friends of progress—like the rocky banks in the river's bed, which only display the resistless force of the flowing stream.

In reviewing the progress of the past we must not give way to wonder and surprise, but rather calmly cultivate a spirit of confidence and anticipation, and be ready to accept fresh light from whatever quarter it may fall upon us. May we, like Sir Benjamin Brodie, desire, above all things, the attainment of the truth, and be ready to pulverise our most cherished convictions, and to cast them away for ever, when truth and progress demand of us the sacrifice.