excision of too short a piece, permitting reunion of the cut ends. It is stated by Hütter¹³ that experiments show that not less than 5 inches must be removed to prevent reunion with certainty. Although I am inclined to think that this is an excessive length, there seems to be little doubt that a considerable gap between the divided ends can be bridged over, but I am convinced that in many of the cases in which a return of pain after neurectomy is ascribed to reunion, the result is in reality due to the occurrence of neuritis in the stump of the nerve, chiefly because the wound was not treated antiseptically. Another reason for a want of permanent relief after neurectomy is, I believe, because the nerve is not fully freed in the bony canal. If the neuritis does not frequently start in such a place, it is, I am sure, often very intense there, and the free removal of nerve in the canal, or the removal of bone so as to widen the aperture and prevent pressure, is essential. Then, too, if the neuralgia has persisted long, it is necessary to go far back, and a minor operation on the anterior portion of a nerve trunk may be followed by a recurrence of the pain within a short time. I believe it is necessary to divide the nerve as near its origin as possible, so as to sever it where it is still healthy.

It is exceedingly difficult to get statistics as to the duration of relief after operations for this affection. Many cases are reported as cured after a few weeks or months, but there is nothing to show that the relief lasted even for a year. In many cases it is known to recur within that time, but I think that if the principles I have laid down are followed, recurrence will occur far less often than it has hitherto.

The occasions on which I have obtained what may be regarded as absolute cure or have observed recurrence of pain are grouped together in the accompanying table, which has been constructed so as to exhibit at a glance not merely the direct effect of the operation, but also the surgical details of the procedure, and the success or failure of the same.

The conclusions I would draw from this table are, that as soon as drugs and electricity have definitely proved unequal to the task of controlling the pain, the branch of nerve affected should be excised. The rapidity with which the wound heals and the absence of a noticeable scar deprive the procedure of obvious drawbacks, while the genuine nature of the relief it affords, in contrast to other methods, is shown by the fact that patients once operated upon will hardly wait to hear of other treatment if some other branch become affected.

PRELIMINARY NOTE ON A NEW CHROMOGENIC MICRO-ORGANISM FOUND IN THE VESICLES OF HERPES LABIALIS. "BACILLUS VIRIDANS."

By WM. ST. CLAIR SYMMERS, M.B.

[From the Pathological Laboratory of the University of Aberdeen.]

Source of the Micro-organism.—It was found in great numbers in the lymph of herpetic vesicles which occurred on the lip of a boy suffering from typical acute croupous pneumonia. The lymph was gathered on the fifth day of the disease, the vesicles being then two days old.

Biological Observations.—The microbe grows luxuriantly on many nutrient media: on some it elaborates a beautiful pea-green pigment which permeates the medium, while the microbe itself is, when seen in mass, uncoloured, and has a greyish-white appearance. Light has no effect upon its growth or chromogenic powers. It flourishes well at ordinary temperatures, but grows somewhat faster when kept at a temperature of 100° F. It is aerobic to a certain extent, inasmuch as it flourishes best in contact with atmospheric oxygen, but still it is facultatively an anaërobie, and can grow, though slowly, in the absence of air—for example, in an atmosphere of pure hydrogen. Although, however, the microbe grows abundantly in absence of air, it cannot then produce its peculiar pigment.

Among other experiments the following was arranged. The microbe was inoculated on the sloping surface of a tube of Koch's nutrient gelatine, the whole being immersed in hydrogen gas, and care, of course, being taken that the air in the test tube was replaced by hydrogen. The gelatine slowly became liquefied, the whitish organismal mass sank to the

bottom of the now fluid gelatine, which remained a perfectly clear limpid amber in tint. Upon removing the test tube from the hydrogen, and thus readmitting air, the solution soon took on the usual green colour. From this and analogous experiments it was concluded, first, that the organism could live and vegetate apart from oxygen; secondly, that for the full production of the pigment oxygen is absolutely necessary; and, thirdly, that the green pigment is not secreted as such, but, in all probability, as an immature substance which easily gives rise to it on contact with oxygen.

On the other hand, the microbe on some media grows most luxuriantly, and, notwithstanding the presence of oxygen, does not produce pigment; for example, on cocoanut, on potato, etc., it grows readily, but produces no pigment. Again, the usual media in which the pigment appears may, by the addition of an antiseptic, such as creasote, be rendered unsuitable for the elaboration of the pigment, while still affording a rich nutriment for the microbe. Thus, if 1 per cent. of creasote be added to bouillon the medium soon becomes cloudy after inoculation with the microbe, but no pigment appears; the cloudiness deepens into milky opacity in time, the fluid teems with well developed microbes, but the pigment is not present even in the faintest degree. These are merely temporary aberrations in its physiological activity, nor has it yet been possible, though many attempts have been made, to obtain the organism disassociated permanently from its chromogenic attribute. As soon as the organism has been placed in proper surroundings the pigment-forming power reasserts itself. The proper surroundings consist, then, in a suitable nutrient medium, such as Koch's gelatine, in the presence of free oxygen, and in the absence of such restraining influences as antiseptics. Given these conditions the microbe, no matter through how many generations its power of pigment formation has lain dormant, at once rises to its old activity and produces the green as beautiful as ever.

In the accompanying lists A and B are to be found the media which are best suited for the elaboration of the pigment, and also those which do not offer such occasion. It is understood, in both cases, that abundance of free oxygen has been supplied.

A. Pigment Formed.—On agar-agar (neutral), on agar-agar plus 6 per cent. glycerine, on agar agar (alkaline), on Koch's gelatine, on Koch's gelatine (alkaline), on bouillon, on blood serum, on milk, on cocoanut water, on cocoanut water and gelatine, on tuberculous sputum, on fish broth, on Wassermann's fluid,¹ on Wassermann's fluid plus 1 per cent. of milk sugar (very good), on Beyerinck's medium.²

B. No Pigment Formed.—On agar-agar (very acid), on Koch's gelatine (very acid), on potato (both acid and alkaline), on beetroot, on turnip, on carrot, on parsnip, on cocoanut, on urine.

Morphology.—The microbe occurs either as a rod or thread form. The threads are either in true or pseudo-filaments, that is either as long solid filaments or as strepto-bacteria. The filaments are mostly solid, without septae; they are sometimes very long, with tapering ends, often twist back upon themselves, curving in various directions, and are flexible. Neither cocci nor spirals have as yet been seen, although attempts have been made to obtain such forms artificially after the manner of Charrin, with bacillus pyocyaneus.³ Free spores do not occur as a rule, but by influencing the growth of the bacillus by means of antiseptic or other poisonous substances, cylinders filled with spore-like bodies have been obtained. Dumb-bell and drumstick forms have also been found in great quantity, the drumstick forms being similar to those of the bacillus of tetanus. The fact of the above being true spores is perhaps not beyond doubt. It may be mentioned that no green-producing organism hitherto described has been observed to form spores, the only exception being the bacillus pyocyaneus, in which Professor Ferrari, of Catania,⁴ has, contrary to many observers, described their formation.

The Pigment.—I have not been able to separate this in a pure form as yet. It is, as seen in solution of melted gelatine, dichroic, green by reflected, and straw yellow by transmitted, light. It is difficult to see it by gas light. When several weeks old it changes colour, gradually becoming like burnt sienna, and ultimately assuming a crimson tint in addition to the sienna colour. Both mineral and organic acids decolorise it without destroying it; on neutralising with an alkali the green colour returns. Boiling has no effect upon it. Liquor ammoniæ intensifies the normal green colour; and in media which do not appear to contain the pigment it

¹ *Annales de l'Institut Pasteur*, 1887, p. 591.

² SUPPLEMENT TO BRITISH MEDICAL JOURNAL, January 10th, 1891.

³ Charrin, *La maladie pyocyaneuse*, Paris, 1889.

⁴ *L'Ateneo Medico Parmense*, 1889, page 115.

¹³ *Grundriss der Chir.*, i, 144.

can be at once made evident, if present, by adding liquor ammoniæ. It is insoluble in chloroform, thus showing that it is not pyocyanin. Sulphate of ammonium and acetic acid precipitate it from its solutions along with the albuminoid substances which are rendered insoluble by this salt. The pigment appears to be identical with that of the bacillus virescens of Frick, but this organism is quite distinct from the one described by him.

Classification of Organism.—According to the latest investigator on the subject—those of Frick, of Zürich⁵—there are only a few such organisms that can be recognised with certainty. These he gives in a tabular form after the manner of Lusenbergl—

Bacillus "A."	pyocyanus	Bacillus fluorescens liquefaciens.	Bacillus viridis pallescens.
Bacillus "B."	pyocyanus	Bacillus virescens.	Bacillus iris.
		Bacillus fluorescens.	

Of these there are only two with which the bacillus herein described could be confounded, namely, bacillus fluorescens liquefaciens or bacillus virescens. From the bacillus virescens—the subject of an exhaustive investigation by Frick—it can easily be differentiated by the fact that the bacillus viridans liquefies gelatine, whereas Frick's bacillus has no such action. There are several other distinguishing features which need not be mentioned here. As regards the bacillus fluorescens liquefaciens, it never forms threads, never grows out into filaments, is purely aerobic, spores are not observed, it thrives badly at higher temperatures (100° F.), and liquefies gelatine slowly, in all of which it differs markedly from the bacillus of this communication.

Pathogenesis.—The organism appears to have a pathogenic action on rabbits and white rats; patches of alopecia and ulcerations have been observed on rabbits when the microbe was injected subcutaneously. These experiments were performed on six rabbits only, and, until further investigation has been undertaken, no conclusions are warranted.

ABSTRACT OF PAPER ON SENILE CHANGES IN BONES AND SOME SENILE DISEASES OF THE OSSEOUS SYSTEM.

Delivered to open a Discussion on the Subject in the Section of Pathology at the Annual Meeting of the British Medical Association, held at Bournemouth, July, 1891.

By F. S. EVE, F.R.C.S.,

Assistant Surgeon London Hospital; Surgeon (Out-patient) Evelina Hospital; late Pathological Curator Royal College of Surgeons.

(Concluded from page 1195.)

I CANNOT leave the subject of mollities ossium without mentioning the beneficial effects thereon resulting from removal of the ovaries. The point of departure for this proceeding was the case of a woman in whom a cure of the bone disease took place after Porro's operation performed by Fehling for pelvic deformity. This surgeon and others have now treated some twenty cases of mollities by oöphorectomy with great success. The ages of the patients ranged between 28 and 51 years, and they were mostly multiparæ. The affection had, as a rule, existed several years, in one case for thirteen. Fehling thinks that the cause of the osteomalacia must be sought in a pathological hyperactivity of the function of the ovary.

The disease is said to have been experimentally induced by semistarvation in a pregnant dog.¹⁰ Possibly it is connected in some manner with the blood-forming properties of the marrow, for it attacks chiefly those bones in which this is most active—namely, the pelvis, vertebræ, ribs, and skull, and its common occurrence during pregnancy, when great demands are made on nutrition, lends some colour to this supposition.

⁵ Vide Virchow's *Archiv für pathologische Anatomie und Physiologie*, 1889, p. 266.

¹⁰ Stilling and v. Mering, *Arch. f. med. Wissen.*, p. 803, 1889.

A condition of the bones nearly resembling mollities is occasionally met with in cancer, itself a disease of advancing life. In illustration of this reference may be made to a case I had the opportunity of investigating. The patient, a woman, aged 40, was affected with a very extensive soft carcinoma of the omentum and peritoneum with secondary

Fig. 3.

growths in the liver, the lumbar and mesenteric glands. The cancellous bones, especially the ribs and ilium, were so soft that they could be cut with a knife. The calvarium was slightly softened but not thickened. In the femur the medullary cavity was enlarged, and its upper half filled with soft red, the lower with fatty, medulla. The body was much emaciated and deeply jaundiced.

The high degree of atrophy of the bones is very perceptible to the naked eye on holding microscopic sections of rib, ilium, and femur up to the light. All the cancellous tissue of the rib (Fig. 4) is seen to have been removed, only a

Fig. 4.—Slightly magnified.

very thin wall of bone remaining. The cancellous tissue of the ilium forms a very wide-meshed network. The femur is less affected. The minute changes (see Fig. 5) do not differ in kind, but only in degree, from those found in true mollities, but the areas of decalcification on the outer margins of the trabeculæ are not so extensive or widely spread. The marrow appeared similar; there was little fatty degeneration and no evidence of cancerous growth. The fact that the vertebræ, ilium, ribs, and skull were more affected than the long bones is another point of resemblance between this condition and mollities.

[In conclusion the author expressed his indebtedness to the President of the Royal College of Surgeons and Professor Stewart for the loan of many specimens with which the paper