BRITISH MEDICAL JOURNAL

LONDON SATURDAY FEBRUARY 6 1943

THE TREATMENT OF CEREBRAL CONTUSION*

BY

LAMBERT ROGERS, M.Sc., F.R.C.S.

Surgeon Captain, R.N.V.R.

Professor of Surgery, University of Wales; Surgical Consultant, R.N. and E.M.S.

Cases of cerebral contusion following so-called "closed" head injuries are usually admitted to beds in the general surgical wards of a hospital, and two tendencies may be noticed in regard to them: first, like fracture cases in the same wards, they are not infrequently regarded rather as necessary evils and relegated to the care of the house-surgeon; secondly, treatment is usually of a symptomatic character—e.g., for restlessness—without due consideration for the complex pathological processes which result from the injury to the brain and are involved in its recovery. It is not surprising, therefore, to find these cases either just left recumbent, or else subjected to various methods of treatment, which for the most part may be described as of an expectant character, consisting merely of rest in bed and efforts to palliate complications as they appear.

What, then, should we do for a patient rendered unconscious by a severe head injury and who may or may not present an obvious lesion such as a scalp wound, for which local treatment is required? Although such cases are of frequent occurrence it is true to say of them that, first because of the extreme variability in the severity of head injuries and of their effects, rarely, if ever, do any two have identical cerebral lesions; and, secondly, even after a prolonged period of unconsciousness some of them make remarkable recoveries, even if given little or no treatment at all—e.g., just left to lie on a bunk, as happened in sailing-ships in olden days. Obvious difficulties, therefore, are encountered in testing the efficacy of any particular form of treatment for such cases. The fact remains, however, that many patients who recover from the immediate effects of closed head injuries do present sequelae in the form of the so-called post-concussional state. We may accordingly ask whether it is not possible to lessen the incidence or altogether avoid this post-concussional state by adopting prophylactic measures in the form of early treatment. It is my belief that it is possible, and the object of this paper is to record experiences with a method of treatment which I contend is indicated on theoretical, experimental, and clinical grounds, and which appears to promote the recovery of these patients, with a singular freedom from complications or after-effects.

If, following a particular form of treatment, convalescence is smoother than it is otherwise known to be, and if there is a consistent absence of sequelae known to occur from time to time in cases not so treated, it is suggested that we have a strong argument in favour of that treatment—provided always, of course, that the treatment itself does not add to the mortality of the condition.

Testing the Value of Treatment

What methods have we for testing the value of a particular form of treatment? The answer is not as simple as it might seem to be at first sight. It would appear that all we had to do was carry out direct experiment—i.e., test the remedy; but it has already been pointed out that many patients recover whether a particular form of treatment is given or not, and they may do so even despite it. Next, since no two cases of head injury—which term implies a complex and variable

pathology—are quite alike, it is difficult to obtain controls by which to effect anything like exact comparison. Lastly, while a certain amount of information is to be gained from animal experiments, there are obvious fallacies in transferring the results of such experiments to man or in attempting to draw conclusions from clinical observations made on animals. To test the evidence, therefore, the most scientific course will be to adopt a combination of methods such as the following:

(i) To examine the argument for the particular remedy by careful consideration of the premisses on which it is based and the conclusions to be derived therefrom.

(ii) To review the experimental evidence bearing on the argument.

(iii) To examine the results of a series of treated cases, followed over some years, particularly noting the incidence of complications known to occur from time to time with other methods of treatment.

(iv) To review a few cases in some detail while, so far as possible, making allowance for the qualifications already mentioned.

It is first necessary to indicate the type of treatment. Briefly, this consists of continued gentle dehydration, instituted with the object of withdrawing cerebrospinal fluid from the intracranial cavity and so providing potential space into which brain-swelling can take place, thereby avoiding compression of its blood vessels.

I. The Argument

The argument in favour of this form of treatment is briefly the following: Bruised tissue, whether that comprising the brain or any other organ or structure, must be permitted to swell if it is to recover completely. If by confinement within a limited space swelling is prevented there must be a reduction in blood supply owing to the compression of the vessels supplying the damaged tissue, and this reduction is detrimental to complete recovery.

While to-day the Monroe-Kellie doctrine may not be wholly acceptable in the form in which it was originally propounded, it is probably true to say that with certain reservations the data on which it is founded are true and that in general it holds good. There is a relatively fixed total volume of the cranial contents, but with a capacity for change in any one of the three chief elements comprising it—viz., the brain itself, the circulating blood, and the cerebrospinal fluid. The most practical method of providing for swelling of the bruised brain is to reduce the amount of the cerebrospinal fluid within the skull. This may be conveniently done by dehydration.

If tissue-swelling is allowed to take place after injury the subsequent recovery of the tissue is striking; if, however, swelling is restricted or prevented altogether, a varying degree of permanent damage follows. This important principle of pathology is exemplified when an organ readily capable of expansion—like the kidney—is compared in its reaction to injury with an organ restricted by its capsule from expanding—like the testis. The renal capsule is thin and readily distensible; the testicular capsule (tunica albuginea) is tough, fibrous, and comparatively inelastic. The bruised kidney swells, recovers, and functions normally; the bruised testis cannot swell, and subsequently atrophies. The cerebral envelopes are even more

* Based on a lecture delivered at the Royal Naval Hospital, Plymouth, Jan. 7, 1943.

rigid than the tunica albuginea, and the only way in which some variation can be given to their limits of constriction, other than by craniotomy (with its attendant risks and disadvantages), is by reducing the volume of the cerebrospinal fluid surrounding the brain, the fluid itself being incompressible. As Wilfred Trotter (1929) has pointed out, a simple bruise, of no ultimate importance to an organ with a yielding capsule, is a serious matter with the brain, which must be allowed to swell if its circulation is not to be materially interfered with. If this swelling, which is the normal reaction to injury, is restricted, the blood supply to the bruised region is diminished, and the ensuing ischaemia may well result in cortical atrophy with associated ventricular distension or become manifest by symptoms such as those which comprise the post-concussional state. This post-concussional state may conceivably represent incompletely resolved cerebral contusion or subsequent atrophy consequent upon failure to provide for brain-swelling following injury.

Brain-swelling.—In the case of the brain, injury produces swelling very readily and to an extreme degree. Thus, Dandy (1941) writes: "Trauma that would cause no concern to other harder tissue produces severe swelling of the brain." Some idea of the extent of this swelling and the rapidity with which it occurs can be gathered from cases of cerebral tumour which have undergone operative treatment. In this connexion the same author writes: "The large cavity created by the tumour's removal may be entirely obliterated in less than half an hour, and with no gross haemorrhage."

One of the arguments advanced against the routine use of dehydration to permit brain-swelling to occur in cases of closed head injuries is that from time to time the pressure of the cerebrospinal fluid in the lumbar pond of patients unconscious as a result of a severe head injury has been found to be low. This, however, does not invalidate the thesis or contraindicate dehydration—first, because experimentally (White *et al.*, 1942) no correlation has been found to exist between cerebrospinal fluid pressures and the degree of cerebral swelling (e.g., as produced during anoxia); and, secondly, because variations in the pressure of the fluid in the lumbar pond take place so frequently that an isolated observation that at one particular time it is low is probably of little significance. Finally, it is reasonable to assume that such pressure differences as occur during brain-swelling are chiefly of a local character and may require intracellular observations to detect them; or, as White (1942) has recently suggested, in head injuries there is often concomitant anoxia of the brain due to depression of the respiratory centre, and this anoxia may considerably reduce the secretory activity of the choroid plexus.

Still further evidence that the findings at lumbar puncture are not necessarily an indication of the intracranial fluid state has been found at operation; thus Bathe Rawling (1918) wrote in connexion with subtemporal decompression which he was at that time performing for cases of suspected cerebral oedema: "In one of my last cases four punctures carried out immediately previous to operation showed a great diminution in the amount of cerebrospinal fluid; and, in the operation performed immediately afterwards, the opening of the dura showed a most typical condition of cerebral oedema with very great excess of fluid."

II. Experimental Evidence

Experimentally it should be possible to ascertain whether brain-swelling takes place following head injury: White (1942) has found a very constant swelling of the brain of cats, even after the animals have had mild concussion—e.g., the brain, instead of being 10 to 11% smaller than the intracranial chamber, was found to be only 7 to 8% smaller. Surgeons (including myself) who have operated on cases of closed head injury within the first few hours have commented on the swollen condition of the brain. Furthermore, it is a widely held view, and probably true, that when a compound fracture with its immediate decompressive effect has occurred convalescence is easier and freer from complications than when a comparable type of injury has produced cerebral contusion alone.

Turning next to the other side of the problem, Weed and McKibben (1919) constantly found histological changes in the brains of animals given intravenous injections of hypertonic or hypotonic solutions if by keeping the skull closed the brain volume was not allowed to alter. On the contrary, however,

when the brain volume was allowed to alter by opening the skull with a trephine these histological changes did not take place.

Provision for Brain-swelling.—If brain-swelling which is the normal reaction to injury is allowed to take place the blood supply remains relatively unimpaired and resolution and recovery are complete. If the swelling is restricted local blood supply is impeded and may be followed by changes which are evident clinically as the "post-concussional state." Theoretically, provision may be made for brain-swelling by performing craniotomy or by withdrawing the incompressible cerebrospinal fluid from the skull; for, as Dandy (1933) has pointed out, "space compensation is the principal function of the cerebrospinal fluid." This fluid-withdrawal may be either by lumbar punctures or by dehydration, or it may be effected by a combination of both methods. In practice it has been found that craniotomy is unnecessary and undesirable and that dehydration is the procedure of choice, probably because this is a continuous and gentle process, whereas repeated lumbar punctures result in a rapid to-and-fro swing of cerebrospinal fluid production and thus are only occasionally employed to augment dehydration or for diagnostic reasons.

Dehydration by Magnesium Sulphate.—The question arises whether it is possible by any simple measures to produce the continued gentle dehydration previously advocated, and the reply is that this can be effected by instillations of magnesium sulphate into the rectum at regular intervals. Cushing and Foley (1920) showed that 20 to 30 c.cm. of a saturated solution of sodium chloride introduced into the duodenum or the rectum of the cat produced a maximal fall of cerebrospinal fluid pressure precisely comparable to that which occurred when the solution was given intravenously. They pointed out, moreover, that when the salt is given by the alimentary route the intracranial changes which are produced are not only not attended by disturbances of pulse, respiration, or blood pressure, but that possibly, also, alteration of the cellular elements of the blood is avoided. Temple Fay (1924) found that magnesium sulphate instilled into the rectum had advantages over sodium chloride in causing less discomfort and being almost twice as efficient in its effects on the cerebrospinal fluid. Being non-dialysable and therefore not absorbed into the blood stream, magnesium sulphate does not produce a secondary wave of tissue oedema with an increase in intracranial pressure as does the sodium salt. Writing of the rectal use of magnesium sulphate in the University Hospital in Philadelphia, he states: "The beneficial use of this salt in cranial injuries has been well established in this clinic." This has also been our experience at Cardiff. Its use in the treatment of cerebral oedema has also been endorsed by Donald Munro (1938) and others.

The details of treatment are briefly as follows: (1) The patient is propped up in bed with the head raised to reduce intracranial venous congestion, and for the same reason it is necessary to make sure there is no constriction of the neck veins by tight pyjama collars or other garments. (2) Retention enemas of 6 oz. of 30% magnesium sulphate solution in water are given every six hours. These must be very gently and slowly administered so as to be retained. (3) Fluid intake by mouth is restricted and a certain degree of thirst is a criterion of effective dehydration.

III. Examination of Results

In 1929* this practice was begun in the surgical unit at the Cardiff Royal Infirmary, and has consistently been followed since in all cases of head injury severe enough to be admitted (Lambert Rogers, 1935, 1937, 1938a, 1938b). The majority of these patients were in a state of profound or prolonged unconsciousness. In the years between 1929 and 1940 550 cases were treated. Fractures of the skull were detected in 133 of these, but there is reason to believe that many others had basal fractures not revealed by x-ray examination. Of these patients 25 died, the majority soon after admission.

An absence of both immediate and late complications has been noticed. Traumatic delirium (cerebral irritation) was

* The idea followed Wilfred Trotter's teaching of the pathology of brain injury, and I had thought we were the first to institute the practice, but have since read that this treatment was in operation in 1921 on the neurosurgical service of the late Dr. Charles H. Frazier (Temple Fay, 1935).

infrequent, and when it did occur was only of a mild character. The absence of the noisy, shouting, restless type of patient with head injury, previously so often seen, was a welcome change which has often been commented upon by those taking part in the dehydration treatment. Furthermore, if focal signs arose in the course of treatment these were considered to indicate craniotomy, and were found at operation to be caused by blood-clot and not oedema, which in some cases previous to the dehydration regime had been revealed when these cases were operated upon because of the onset of focal signs. Lastly, although a follow-up clinic is maintained in connexion with the surgical unit, not one of these 525 cases has required readmission because of the so-called post-concussional state or sequelae of any kind. It is tempting to doubt, therefore, whether there is such a condition as the post-concussional state if brain-swelling is permitted to take place at the time of injury, and whether in fact this state, as has already been suggested, does not represent incomplete resolution the result of failure to institute effective initial treatment. It is perhaps pertinent to remark that during the period under consideration cases of so-called post-concussional state were admitted from other clinics where dehydration had not been carried out.

IV. Review of Some Cases

The following details of a few cases exemplify the result of this treatment.

Case 1 : Fracture of base of middle fossa extending into right parieto-occipital region of vault. Cerebral contusion and laceration. Spastic paraplegia, right brachial monoplegia, left facial palsy, cyanosis, coma three weeks, cerebral irritation, bilateral suppurative otitis. Dehydration 15 weeks. Recovery complete.—A physician aged 53 was admitted after a motor accident, a fast-travelling lorry having collided with the side of the taxi-cab in the back seat of which he was sitting. On admission he was cyanosed and in profound coma; respirations were deep and noisy; he was bleeding from both ears and nostrils. There was a superficial scalp wound on the forehead and a haematoma over the occiput. There was a slight right facial palsy; the right arm was spastic, the left flaccid; there was spasticity of both legs, and the plantar reflexes were extensor. Death appeared to be imminent. He was propped up and dehydration begun. Lumbar puncture two days after admission showed a pressure of only 100 mm. water and red opalescent fluid; 20 c.cm. were withdrawn, and this was repeated 2 days later. Systolic blood pressure, 164 mm. Hg. Radiographs showed a "vertical fissure in the right occipito-parietal region." At this time he was groaning unceasingly, and was rather restless and still profoundly unconscious; the right arm was paralysed, but he was moving his left arm and both legs. The facial palsy had increased slightly. Three weeks later he developed suppurative otitis media in both ears. He was irrational but recovering consciousness, though unable to recognize an old friend. Later still the plantar reflexes became flexor; he was less excitable and had no headache, but complained of tingling in his right arm. Blood pressure, 130 mm. Hg. Dehydration was continued for a period of 15 weeks. Recovery was complete, and he returned to active consulting practice. For the seven years which have since elapsed he has worked consistently, is apparently normal in all respects, and is at present working harder than ever.

Case 2 : Linear closed fracture. Cerebral contusion, coma, delirium, right monoplegia soon proceeding to hemiplegia, middle meningeal haemorrhage. Bone flap; removal of 83 g. of extradural blood-clot. Dehydration 1 week. Recovery complete.—A farmer aged 33 was involved in a brawl in a local inn and hit with a ploughshare on the left side of his head. He was a big muscular man, over 6 ft. tall, and on admission was unconscious and very restless. Pupils reacted very sluggishly to light and both plantar reflexes were extensor. Dehydration was begun on admission. He remained unconscious for three days, but gradually became more docile as he was dehydrated. He then developed a right monoplegia which soon progressed to a hemiplegia. The systolic blood pressure was 90 mm. Hg. A linear fracture traversed the left parietal and squamous temporal bones. A left parietal bone flap was turned down and 83 g. of extradural blood-clot removed. He made a complete recovery, power returning first in the leg and then in the arm.

Case 3 : Closed head injury—probable basal fracture—unconscious 48 hours. No dehydration; intense headache, bradycardia; semi-conscious and very restless when seen. Dehydration 14 days. Rapid improvement and recovery.—An Army officer aged 35 fell from a lorry, struck his head, and was admitted unconscious to a cottage hospital. He was kept lying in bed for five days, and when then seen was semi-conscious, restless, vomiting, and complaining of intense headache. The pulse rate was 56; the cerebrospinal fluid was intimately mixed with blood. He was propped up and dehydration begun. The change was rapid, his headache being relieved, the

vomiting ceasing, and restlessness subsiding. The matron of the cottage hospital volunteered the remark that the magnesium sulphate instillations relieved his headache. He was dehydrated for a fortnight, and was discharged symptom-free.

Case 4 : Closed fracture, right parietal, following fall of 20 ft. Unconscious 2 hours, subsequently drowsy. Dehydration 5 days. Recovery complete.—A dockyard labourer aged 22 had an epileptic fit and fell from some staging about 20 ft. He was admitted to hospital, remained unconscious two hours, and was subsequently drowsy. He remembered feeling he was going to have a fit and then later finding himself in hospital. He was dehydrated for five days. The medical officer reported that he became less drowsy after the dehydration and made a complete recovery.

Case 5 : Closed fissured fracture, left parietal, following fall from lorry. Unconscious for few minutes. Wound of pinna. Dehydration 3 days. Recovery complete.—A marine aged 35 fell out of the back of a lorry and was unconscious for a few minutes only. On admission to hospital he had headache, was drowsy, and desired only to sleep; his cerebration was slow. He was dehydrated for 3 days, and the medical officer noted: "Patient benefited from dehydration. It relieved headache and he became brighter."

Case 6 : ? Fracture in parieto-occipital region. Closed head injury. Fall off back of lorry; unconscious 15 hours. Intense headache. Dehydration 7 days. Recovery complete.—A pensioner yeoman of signals aged 42 fell off the back of a lorry and was admitted unconscious. He remained so for 15 hours, and, although when he regained consciousness he was mentally alert, he was depressed and had intense headache. Medical officer noted: "Patient obtained benefit from dehydration. Headaches were relieved."

Case 7 : Closed injury. Patient found lying unconscious beside bicycle in road. Abrasions of face, hands, and leg. Dehydration 18 days. Recovery complete.—An A.T.S. officer aged 30 was admitted unconscious to hospital, having been found in the road beside her bicycle. She remained unconscious for 6 days and was then unresponsive and drowsy, and complained of intense headache. The optic disks showed slight papilloedema, and the pulse was consistently below 60. The electro-encephalogram showed no abnormality in any area, but she was not made to overbreathe. When she was first seen either brain-swelling or a subdural haematoma was suspected. Remarkable changes followed dehydration. Once this was effectively instituted she made an uninterrupted and rapid recovery, and was discharged symptom-free.

Case 8 : Closed injury following fall off lorry; semi-conscious, traumatic delirium, hemiplegia. Dehydration 3 weeks. Recovery.—A soldier aged 25 fell out of a lorry on to the road and was admitted to a local hospital in a semi-conscious condition. He became very restless and irritable, and was noticed to have a hemiplegia. When seen 48 hours afterwards he was still only semi-conscious, very restless, irrational, irritable, and hemiplegic. The right plantar reflex was extensor, the pulse between 50 and 60. Lumbar puncture showed blood intimately mixed with the cerebrospinal fluid. Following dehydration he made a rapid recovery, his symptoms disappearing from the time of its institution.

Not once, but many times recently, I have had the experience of finding a young Service patient with a head injury lying flat in bed, unconscious, irritable, and restless, and within a few hours of propping him up, doing a lumbar puncture, and beginning dehydration therapy, seen him become a conscious rational being once more. I attribute such changes to the recognition of the importance of brain-swelling and the provision thus made for it.

While there have been difficulties in getting this treatment accepted as a routine measure in this country and few British neurosurgeons, with the exception of Wylie McKissock (1942), have openly supported it, there have not been lacking elsewhere those who have upheld the claims made for it. In the United States Foster Kennedy and Wortis (1932) have advocated dehydration treatment, and Temple Fay (1937) claims not only a reduction in mortality of somewhere about 25% in head injury cases so treated but that there is a remarkable improvement from the standpoint of subsequent economical and social adjustments. He writes: "If its [the brain's] tissue can be maintained alive during the crisis, a prompt return of function will follow to a far greater degree than has formerly been recognized. . . . What surgeon would permit a patient with a bed-sore to continue pressure on the area and expect the devitalized anaemic tissue to recover its function? Yet many . . . permit a brain to become involved in a relentless hydraulic compression, squeezing the blood out of this vital and important tissue." On the Continent, Decker (1938) of Lausanne has upheld its value, claiming a lowering of mortality and improvement in results since its adoption, and there have been others who from time to time have spoken or written in favour of it.

But for the most part these have been "voices crying in the wilderness." Anything like routine appears to be abhorrent to some surgeons, yet the reaction to injury must at least be always in the same direction, and if bruised brain always reacts by swelling, treatment surely should always be directed to providing for the swelling to take place.

Conclusion

To sum up, then, we may inquire whether the Monroe-Kellie doctrine does hold in regard to the reaction of the bruised brain to injury. Is the encapsulation of the brain by cranium, meninges, and cerebrospinal fluid of a comparatively rigid and inelastic character like the tunica albuginea of the testis, as has already been suggested, or does it by some physiological mechanism readily accommodate to cerebral swelling as does the thin elastic renal capsule to swelling of the bruised kidney? If, as it seems to me, the encapsulation behaves in a comparatively rigid and relatively inelastic manner, there can be no questioning the desirability of directing our treatment of the bruised brain towards an attempt to render the cerebral encapsulation more accommodating and thus to maintain the blood supply to the contused area by permitting it to swell in response to the law of its reaction to injury. It is further contended that the best method of doing this is by continued gentle dehydration to facilitate brain-swelling and so prevent the reduction in the blood supply to the damaged areas of the contused brain.

There is some reason to think that our forefathers may have had some inkling of the benefits of dehydration. Ambroise Paré over 300 years ago advocated restriction of fluids, purgation, and sweating for cases of head injury (Turnbull, 1939); and the prescribing of calomel and salts and otherwise purging of patients with fracture of the base of the skull, which was common practice early in this century, was probably introduced quite empirically, but was nevertheless essentially sound.

Perhaps in addition to the striking clinical improvement which is so often observed as soon as this treatment is instituted, the absence of sequelae in a long series of cases is the best criterion on which to base its claims. It is simple—so simple that it tends to be neglected or perhaps rather scornfully regarded, as Naaman was inclined to treat the advice of Elisha (ii Kings v). The Syrian captain was angered and disappointed that the prophet had not come down and performed some miracle. In these cases of severe closed head injury with profound unconsciousness it would be so much more dramatic to perform wide decompressive craniotomy, but the simpler treatment is so effective that, like bathing in the Jordan, it suffices. To those who are still not convinced by the arguments here put forward, the figures presented, or the cases instanced, one may quote what was once graphically said of some numerical evidence: "An emaciated child suffering from hunger is a more impressive sight than any number of mortality statistics." A few cases treated by this method will prove more convincing than any theoretical considerations.

Summary

Prolonged unconsciousness following a severe head injury, particularly when blood is intimately mixed with the cerebrospinal fluid, would indicate that the brain has been bruised or torn. Evidence is presented that the brain reacts to such injury by attempting to increase its volume.

No two head injuries are exactly comparable, and it is well known that remarkable and complete recoveries are often made from extensive injuries to the skull and brain. It is difficult, therefore, to assess the value of any particular type of treatment. The absence of sequelae following the consistent application of any particular form of treatment, however, may be urged in its favour.

Dehydration is a method of permitting brain-swelling to take place after injury, and the evidence in favour of its application is here examined. It should be effected by rectal magnesium sulphate and not by intravenous methods, which probably produce undesirable reactions.

Smoothness of convalescence and an absence of sequelae—e.g., the so-called post-concussional state—have been noticed after treatment of a large series of cases by continued gentle dehydration.

It is suggested that the post-concussional state may, in many cases at least, be the result of failure to introduce effective treatment in the early stages and represent a condition of unresolved contusion, it being maintained that resolution takes place readily and completely if the requisite condition for reparation is provided. It is suggested that this is the provision of space so that brain-swelling can occur and is not restricted by the cerebrospinal envelopes, with consequent impairment of the cerebral blood supply. Provision for such swelling is made by sustained dehydration, which reduces the incompressible water content of the skull.

REFERENCES

- Cushing, H., and Foley, F. E. B. (1920). *Proc. Soc. exp. Biol.*, N.Y., 17, 217.
 Dandy, W. E. (1933). *J. Amer. med. Ass.*, 101, 772.
 — (1941). *Dean Lewis's System of Surgery*, vol. 12.
 Decker, P. (1938). *Helv. med. Acta*, 5, 808.
 Fay, Temple (1924). *J. Amer. med. Ass.*, 82, 766.
 — (1935). *Ann. Surg.*, 101, 76.
 — (1937). *Surg. Clin. North Amer.*, 17, 1661.
 Kennedy, Foster, and Wortis, S. B. (1932). *J. Amer. med. Ass.*, 98, 1352.
 McKissock, Wylie (1942). *Proc. roy. Soc. Med.*, 35, 757.
 Munro, D. (1938). *Cranio-cerebral Injuries*, New York.
 Rawling, L. Bathe (1918). *British Medical Journal*, 1, 499.
 Rogers, Lambert (1935). *J. roy. nav. med. Service*, 21, 2.
 — (1937). *J. int. Chir.*, 3, 109.
 — (1938a). *British Medical Journal*, 1, 685.
 — (1938b). *Ibid.*, 1, 743.
 Trotter, W. (1929). *Nature*, 123, 533.
 Turnbull, F. (1939). *Canad. med. Ass. J.*, 40, 579.
 Weed, L. H., and McKibben, P. S. (1919). *Amer. J. Physiol.*, 48, 550.
 White, J. C. Personal communication, 25/8/42.
 — Verlot, M., Silversone, B., and Beecher, H. K. (1942). *Arch. Surg.*, 44, 1

THE NATURE OF THE RHEUMATIC CHILD

BY

DOUGLAS HUBBLE, M.D.Lond.

Assistant Physician to the Derbyshire Royal Infirmary; Physician to the Derbyshire Hospital for Sick Children

(Concluded from p. 125)

Aetiological Factors in the Rheumatic State

It should be generally conceded that statistical inquiry has a more limited value in such a disorder as the rheumatism of childhood than it has in many other diseases. Diagnosis is more than usually difficult, so that the outlook, the experience, and the bias of the individual clinician have to be assessed. The conclusions, too, are valid only for the time and place in which they are made. This may be true in some degree of all statistical investigations, but it is most true of diseases that are disorders of personality and environment. So it should surprise no one that careful investigations by experienced observers at different times and in different places arrive at contrary conclusions with regard to the aetiological factors of rheumatism. The influences of climate, of heredity, of poverty and its accompaniments, are still keenly debated. It may be said of climate that its influence is equivocal; the disease is certainly not limited to northern latitudes and temperate climates as was once taught (Swift, 1940; Fernando, 1939), and it seems probable that the higher incidence of rheumatism in cold than in hot climates depends on the increased liability to respiratory infections in these areas. In England the geographic distribution of the disease cannot be shown to depend on climatic factors (Morris and Titmuss, 1942).

It is generally admitted that the disease shows a well-marked familial incidence, but in Great Britain most observers agree that this depends more on exposure to similar environmental stresses than on an inherited quality. In America, observers are inclined to attach more importance to the influence of heredity, but there are many facts (some of which are dealt with below) which suggest that the heritable factor cannot be of dominant importance. There are, however, three factors—age, social grouping, and the haemolytic streptococcus—about which opinion may be more dogmatically stated. Of these factors it may be said that unless an individual is between the ages of 3 and 15 it is uncommon for him to develop the disorder for the first time; that if he belongs to the wealthier classes he will with great probability escape it altogether, and that, if he does develop it, then at some phase of the illness, but not always at the start, the presence of the haemolytic streptococcus can be demonstrated. These factors deserve more attention here.