MENINGEAL REACTION IN A CASE OF GLANDULAR FEVER

BY

J. SHAFAR, M.D., M.R.C.P., D.P.H.

AND

J. C. R. WEIR, M.B., Ch.B.

Medical Officers, Mearns Kirk E.M.S. Hospital, Renfrewshire

During the last six months numerous examples of a clinical syndrome which is characterized by enlargement of the lymphatic glands have been observed among members of the Senior Service. The groups of glands in the posterior triangles of the neck and in the axillae have been typically involved in a discrete, painless, and only slightly, if at all, tender swelling; the axillary masses attained the size of a small tangerine, whereas the glands in the neck remained relatively small, being palpable as numerous shotty lumps. The usual presenting features were headache of a persistent and severe type, stiffness of the back of the neck, and general malaise. Epistaxis was a not infrequent accompaniment. More recently, abdominal symptoms consisting of nausea, vomiting, and abdominal pain have predominated; the abdominal pain had led to a mistaken diagnosis of sub-acute appendicitis on several occasions by the medical officer submitting the case, a diagnosis which seemed further supported by the marked tenderness over the right iliac fossa. Distinguishing signs in such instances were the presence of a mass of enlarged glands in both iliac fossae and tenderness in these areas. The other sets of glands involved in the malady were the mediastinal and inguinal groups. The former did not produce any specific symptoms, but were demonstrable by radiography.

The disorder has a benign course, and, apart from perhaps a mild pyrexia in the early stages of the condition, no upset of the temperature or pulse rate occurred. Within one to four weeks the patient was free of symptoms, although the glandular swellings required a much longer period before disappearing. Splenomegaly was rarely encountered. The infectiousness of the disorder was made apparent by the incidence of new cases in the same ward and by its development among a number of the medical and nursing attendants.

Differential counts of the white cells of the blood were carried out, and in practically all a deviation from the normal was demonstrable. The polymorphs totalled some 50%, the remainder being made up of small and large lymphocytes and an average of 10% of monocytes. The Paul-Bunnell reaction was negative with but few exceptions.

The relation of the disorder to glandular fever (infective mononucleosis) requires consideration. The general behaviour of the condition would well fit in with such a diagnosis, but the exceptional exhibition of heterophile antibodies must occasion some doubt. Prof. Paul (1941) draws attention to the discrepancies of this reaction. In his own series of 50 cases he obtained 90% positives, whereas he quotes Erf (1936-7), who found that not more than 40% of cases yielded a positive result. Paul rightly argues that the percentage must be influenced by the diagnostic criteria and by the frequency with which the serological tests are made during the disease. The latter surmise is borne out by the example reported by Himsworth (1940), in which the reaction was negative in the early stages of the illness but subsequently became positive 11 weeks after the onset of symptoms. Further observations are necessary before it can be asserted whether distinct categories can be segregated from infectious mononucleosis on the basis of the presence of agglutinins in the patient's serum to the red blood corpuscles of the sheep, or their absence. What has been convincing, however, is the clear-cut clinical picture and course presented by our cases, which now number almost 200. The similarities to infectious mononucleosis are so many that until further evidence is forthcoming regarding the existence of separate and distinct varieties of "glandular fever" we feel justified in including our cases under this diagnostic title.

Complications have been very few. Jaundice was seen on three occasions. Since it was of an obstructive type, and since catarrhal jaundice had been so prevalent in the wards, the coexistence of these two maladies could not be excluded. A mild bronchitis and nasopharyngitis were not uncommon. A skin rash was seen in 4 instances; but it differed individually in appearance, and in all cases was evanescent.

The case presented here is the sole example of a neurological disorder encountered in the series.

Case Report

The patient, a man aged 28, was admitted to hospital on June 17, 1942, with generalized body pains, malaise, and sore throat of a month's duration. His only significant illness in the past was an attack of pneumonia 11 years ago; he was subject to recurrent attacks of tonsillitis.

Examination showed an unhealthy and congested state of the tonsils and nasopharynx, enlarged discrete and painless lymphatic glands in the neck, axillae, and inguinal regions, and scattered rhonchi in the chest. No clinical evidence of disease was discovered in the other systems. The temperature and pulse rate were normal. There was little alteration in his condition until June 22, when he then complained of severe frontal headache, and in the early hours of the next morning became delirious and restless. Examination revealed rigidity of the neck, photophobia, generalized hyperaesthesia, and a positive Kernig sign. No other objective abnormal signs were found in the central nervous system or in the optic fundi. The temperature had risen to 101° F. and the pulse rate to 100. Chloral and bromide were administered with good effect. In the afternoon of June 23 the above findings were more pronounced and the temperature had risen to 103° F. Lumbar puncture was performed, milky fluid being obtained. On analysis this showed a centrifugized purulent deposit, gross increase in cellular content, and Pandy test ++; there were no organisms in the direct smear or on culture. A differential count of the cells of the C.S.F. showed: polymorphs 87%, large lymphocytes 3%, small lymphocytes 6%, monocytes 1%, endothelial cells 3%. He was placed on sulphapyridine, which was discontinued after four days. On the day after the lumbar puncture a marked improvement in his condition was observed. His pulse and temperature were at normal levels and his delirium and incoherence had disappeared; the frontal headache and the nuchal rigidity were, however, still present. A further lumbar puncture was carried out on June 24; the fluid on this occasion was opalescent and contained a slight purulent deposit. The Pandy test was positive, the Wassermann reaction negative, and a large excess of cells were found (polymorphs 70%, large lymphocytes 4%, small lymphocytes 6%, monocytes 11%, endothelial cells 9%); again organisms were not demonstrable.

The patient continued to improve, and three days from the onset of his meningeal reaction his condition was most satisfactory: the nuchal rigidity and headaches were gone and his temperature had remained normal. The C.S.F. at this date showed a fibrinous pellicle, but was otherwise gin-clear; the Pandy test gave a very faint positive reaction, and the cells were few in number and mainly lymphocytic in type. Culture of the fluid failed to grow any organisms. The results of an examination of the C.S.F. on June 27 were: fluid clear, no pellicle on standing, Pandy test ±, W.R. negative, and but a few cells of the large lymphocytic class. On the 30th C.S.F. analysis gave a result identical with that obtained on the 27th. The patient's progress was satisfactory and he was symptom-free—although the lymphatic glands were still palpable, but smaller—until July 1, when he developed a fresh tonsillitis with secondary enlargement of the draining lymph glands in the neck and a pyrexia of 101° F. A throat swab showed diplococci in short chains, while Vincent's or Klebs-Loeffler's organisms were not found. The inflammatory reaction of the throat cleared up in three days. No further developments occurred and, apart from the persisting but steadily diminishing enlargement of the glands, the patient appeared to have made a complete recovery. He was discharged for a period of convalescence, following which tonsillectomy was recommended.

Differential white cell counts of the blood gave the following results:

	June 18	July 27
Polymorphs %	79.1	51
Large lymphocytes %	12.7	28
Small lymphocytes %	0.8	11
Monocytes %	6.8	7
Eosinophils %	0.5	3

A blood count (July 3) showed: red cells 5,200,000, Hb 90% white cells 14,500. The blood W.R. (June 27) was negative. Two Paul-Bunnell reactions (June 19 and July 7) were negative.

A guinea-pig was inoculated with C.S.F. on June 24 and suffered no ill effects; necropsy performed on July 28 showed no evidence of disease.

Discussion

Uncomplicated infectious mononucleosis invariably has a benign outcome, and sequelae of any significance are rare. Nephritis is not an uncommon event in the course of the disorder, but the renal function remains unimpaired and rapid recovery ensues. Haemorrhagic phenomena, pulmonary features, and jaundice are other complications, and attention has recently been attracted to involvement of the nervous system. Bernstein (1940) details the neurological manifestations encountered prior to the date of his monograph. Peripheral cranial nerves are rarely affected, but persistent anosmia and facial palsy have been recorded. Paresis of the extra-ocular muscles (Fledelius, 1935) and photophobia and nystagmus associated with poor articulation (Marshall, 1939) have also been reported. An extraordinary example of glandular fever with a strongly positive Paul-Bunnell reaction, in which a paralysis of the serratus magnus muscle developed during the third week of the illness and which cleared up in about four months, is described by Richardson (1942). The meninges appear to be the commonest site of affection, and Bernstein states that in at least 12 cases of glandular fever the clinical picture of serous meningitis has been found. Often the meningeal features antedate other manifestations of the disease. Nuchal rigidity, positive Kernig and Babinski signs, convulsions, and mental changes have been present in certain of such instances. Bilateral optic neuritis (Gsell, 1937) appeared in one case. The cerebrospinal fluid in the meningeal syndromes accompanying infective mononucleosis usually contains an increased protein and cellular content (chiefly lymphocytic in type), but examples have been reported in which the fluid has been normal in spite of unmistakable clinical signs of meningeal irritation (Huber, 1938). Huber also mentions a case in which the pleocytosis attained a level of 1,000 cells per c.mm. The condition of the spinal fluid in uncomplicated glandular fever is of some interest, but is neither constant nor characteristic. Excess of cells and positive Pandy reactions have been mentioned, but the sugar content and colloidal gold curves remain normal, and the Wassermann reaction is negative even in the presence of a false positive in the blood. Culture of the fluid has proved negative except on four occasions when *Listerella* were isolated (Schmidt and Nyfeldt, 1938).

A recent communication (Landes, Reich, and Perlow, 1941) deals with a case of infectious mononucleosis the early symptoms of which simulated those of encephalitis. Of particular importance with regard to our own case is the report by Thelander and Shaw (1941). They described 3 cases of neurological complications. The first was that of a meningitic picture in the course of infective mononucleosis; the peculiarity noted here was the fact that the cerebrospinal fluid, which at the onset contained 630 cells per c.mm., at the end of four months continued to demonstrate a pleocytosis consisting of 82 cells per c.mm. The Paul-Bunnell reaction was strongly positive. The two other cases occurred in sisters, one of whom, aged 3½, was suddenly affected by fever, pain in the back of the neck, and a convulsion. No glandular or splenic enlargement appeared during the illness. The C.S.F. contained 40 lymphocytes per c.mm., and the blood showed a white cell count of 17,000 per c.mm., with the polymorphs accounting for 69% of the total. Encephalitis of unknown aetiology was diagnosed. Four days later the other sister, 2½ years of age, became ill with similar symptoms. The cells in the C.S.F. amounted to 70 per c.mm., and were chiefly lymphocytic in type. The total leucocyte count of the blood was 45,700 per c.mm., and a differential count revealed that 81% of these were lymphocytes. Again no enlarged glands or spleen could be found, and the Paul-Bunnell reaction was negative. The authors consider that the findings in the last two cases point either to a diagnosis of infectious mononucleosis or to a hitherto undescribed pathological syndrome, and refer to their large experience with benign lymphocytic meningitis and encephalitis, in neither of which conditions had they seen cases in which the blood picture or the results of serological tests suggested mononucleosis. The third of Thelander and Shaw's cases and the case reported here present many features in common. The large numbers of cases in our series, referred to above, showing similar signs and symptoms and running a similar course, would appear to support our contention that the group, if not glandular fever (as manifested by the demonstration of heterophile antibodies), has a common aetiology and

forms a clinical syndrome, of which the present case is unique in the manifestation of a meningeal reaction.

We wish to express our indebtedness to Dr. J. A. Wilson, medical superintendent, who kindly carried out the laboratory investigations, and to Dr. A. M. Scott for his interest and advice.

REFERENCES

- Bernstein, A. (1940). *Medicine*, **19**, 85.
 Erf, L. A. (1936-7). *J. Mt. Sinai Hosp.*, N.Y., **3**, 113 (quoted by Paul).
 Fledelius, M. (1935). *Acta ophthalmol.*, **13**, 150.
 Gsell, O. (1937). *Dtsch. med. Wschr.*, **53**, 1759.
 Himsworth, H. P. (1940). *Lancet*, **1**, 1082.
 Huber, W. (1938). *Schweiz. med. Wschr.*, **68**, 892.
 Landes, R., Reich, J. P., and Perlow, S. (1941). *J. Amer. med. Ass.*, **116**, 2482.
 Marshall, E. A. (1939). *Amer. J. clin. Path.*, **9**, 298.
 Paul, J. R. (1941). *Lancet*, **1**, 584.
 Richardson, J. S. (1942). *Ibid.*, **1**, 618.
 Schmidt, V., and Nyfeldt, A. (1938). *Acta oto-laryng.*, **26**, 680.
 Thelander, H. E., and Shaw, E. B. (1941). *Amer. J. Dis. Child.*, **61**, 1131.

THE RADIOGRAPHIC POSTERO-MEDIAL BORDER OF THE LUNG

OR THE LINEAR THORACIC PARASPINAL SHADOW

BY

JAMES F. BRAILSFORD, M.D., Ph.D., F.R.C.P.

Radiological Demonstrator in Living Anatomy, University of Birmingham

Under the latter title an editorial in *Radiology*, Aug., 1942, p. 229, draws attention to "a slender line of demarcation which is often seen in antero-posterior or sagittal roentgenograms of the bony thorax and upper abdomen. This line lies on the left side of the lower two-thirds of the thoracic spine and sometimes continues as far down as the plane of the first two lumbar segments. The shadow is frequently an enigma to radiologists and other clinicians viewing such roentgenograms. It is not visible on all films or projections of this portion of the body, but is observed with such frequency that it must be the result of variation in the course and position of a normal structure situated therein." Then follows a discussion in which it is inferred that it is not bony, not due to unilateral spinal ligaments, descending aorta, or inferior vena cava, "but it might be an enlarged and elongated hemiazygos vein." "This editorial is written with the suggestion that this finding be referred to as the left thoracic paraspinal shadow and that its frequency be noted by various radiological observers. In this manner its true incidence may be ascertained and its apparent nature confirmed."

An Important Diagnostic Sign

This radiographic line of demarcation is of great importance in the diagnosis of lesions involving the lower dorsal and upper lumbar spine. It is often seen on the antero-posterior radiographs of the dorso-lumbar area on the left side, but it is seen, though less often, on the right side as well. It extends upwards sometimes to the level of the 4th dorsal vertebra, but only as far as the reflection of the pleura on to the diaphragm, for it marks the postero-medial border of the lung. As will be seen from the accompanying sketch of a cross-section of the body at the 9th dorsal level, the postero-medial border of the left lung, which has been more thickly outlined for emphasis, lies in a direct sagittal line for two or more inches. For the whole of this distance it is separated from the vertebral body by the pleura, fascia, and some areolar tissue. On a plane with the anterior surface of the body it is displaced laterally by the descending aorta and more anteriorly by the heart. On the right side the medial border of the lung extends partly over the anterior surface of the vertebral body and is displaced laterally by the oesophagus and the heart. Antero-posterior radiographs of this area will therefore show the postero-medial boundary of the lung as a straight line lateral to the outline of the vertebra, because on the lung side of the boundary the tissue is relatively radiotransparent, whereas on the vertebral side the tissue is much more radio-opaque. The descending aorta, being more opaque than the lung, will also be seen as a linear shadow lateral to the postero-medial border of the lung where it is superimposed over the postero-medial aspect of that organ. On the right side, because of the shelving margin of the postero-medial aspect of the