in general, instead of the predominant typical anabolic and hypertrophic reaction (still preserved only in adrenals and heart), with toxic doses katabolic and atrophic changes prevailed, resulting in some cases in the death of the animals.

Effect of Hypervitaminosis

The administration of the extra amounts of vitamins produced no considerable effects, if any (compare group 2 with group 1, Table I; and groups 2 and 6 with groups 1 and 5, Table II). Hypervitaminosis B and C (Table II) produced in male rats a small, consistent, but statistically not significant increase of seminal vesicles, liver, and fat deposition. Vitamin B also produced greater gain in body weight and improvement in external appearance and behaviour (the rats appeared to be more lively and, when handled, felt firmer and stronger, and had thicker and more glossy fur than the control groups "Co-V"). Histologically no definite effect of these doses of vitamins could be found on the structure of the thyroid gland, except perhaps a slightly depressing effect of halibut and cod-liver oils. This could easily be explained by the extra amount of iodine given in these oils (Mellanby, 1934; Sampson and Korenchevsky, 1932).

Effect of Hypervitaminosis on Hyperthyroidism

The effect of hypervitaminosis on non-toxic hyperthyroidism (group 4 in Table I, and groups 4 and 8 in Table II) was different from that on toxic hyperthyroidism (groups 3 and 4, Table III). In rats kept on our *normal diet and receiving non-toxic* doses of thyroid hormone, vitamins A and D (in the doses given) did not at all affect the specific reaction of organs and tissues to this hormone, while vitamins B and C produced a slight decrease in the hypertrophy of liver and kidneys. In rats kept on a *minimal normal* amount of vitamins B (Table III) and injected with *toxic* doses of thyroxine, the simultaneous administration of extra amounts of vitamins B was followed by a gain instead of a loss in body weight and the partial restoration of specific anabolic changes instead of a pathological (atrophic and almost exclusively katabolic) reaction of the organs to thyroid hormone. In such animals the atrophic changes were decreased in seminal vesicles, prostate, and liver and replaced by hypertrophy in spleen, while in kidneys hypertrophy became more pronounced. The fact that some typical reactions to thyroxine were not prevented in such animals indicates that in vitamin-treated rats suffering from thyrotoxicosis all the improvements described both by previous workers and by ourselves (including gain instead of loss in body weight) could be explained by the making good of the relative vitamin deficiency produced by, and complicating, the thyrotoxic effects. Histologically the thyroid glands were in the same resting condition as in the rats treated with thyroid hormone alone.

The results of our experiments therefore indicate that for the investigation of antagonistic effects neither toxic doses of thyroxine, which greatly raise the vitamin requirements and produce non-specific changes, nor diets with inadequate levels of vitamin content appear to be suitable.

Conclusions

In the therapeutical application of thyroid hormone its katabolic properties in raising metabolism and especially in reducing body fat are often taken most into consideration.

The results of our experiments (as judged by changes in weights and histological structure of organs) and those of some previous workers emphasize that thyroid hormone also possesses anabolic properties, which can lead to better development of such vital organs as liver, kidneys, spleen, heart, and adrenals in both sexes, and ovaries in females.

As shown especially in our experiments on females, it is possible, using moderate doses of thyroid hormone, to reduce the body fat to about one-fifth without affecting appreciably the gain in body weight, and at the same time preserving the stimulating anabolic effects of this hormone on the development of the vital organs mentioned above.

Thus from small or moderate doses of thyroid hormone favourable therapeutical results can be expected, but when large doses are given the useful effects disappear and the hormone becomes atypically toxic and dangerous.

The useful effects of thyroid hormone cannot be normally and fully operative, unless liberal amounts of vitamins are given simultaneously, because of the increased requirements

in vitamins which arise during conditions accompanied by increased metabolism (such as those produced by thyroid hormone).

Therefore the best possible therapeutical results can be expected from simultaneous administration of thyroid hormone with extra amounts of vitamins.

Since both thyroid hormone and male sex hormones possess hepato-, nephro-, and cardiostrophic properties, and thyroid hormone also produces ovario-, spleno-, and adrenostrophic effects, the possibility of therapeutical application of thyroid hormone in cautious doses with vitamins, or together with sex hormones, should be investigated in some suitable diseases of liver, kidneys, and heart, or when an increased activity of ovaries, adrenals, or spleen is desirable.

The results of our experiments and those of some previous workers do not suggest, as has been stated by certain authors, any direct antagonistic effects of vitamins on thyroid hormone; but the data indicate that some beneficial results of the administration of vitamins in some cases of experimental and clinical hyperthyroidism were probably due chiefly to removal of the relative vitamin deficiency present in these cases.

Grants from the Medical Research Council and the Lister Institute have enabled us to carry out this work, and to them we offer our thanks. We are also grateful to the Ella Sachs Plotz Foundation for a grant from which some of the technical expenses were met. We are indebted to Messrs. Roche Products Ltd. for the generous supply of vitamins, and to Messrs. Parke, Davis and Co. for the liberal supply of desiccated thyroid powder.

REFERENCES

- Abelin, I. (1933). *Hoppe-Seyl. Z.*, **217**, 109.
 Cowgill, G. R. (1934). *The Vitamin B Requirement of Man*, p. 65, New Haven.
 — and Palmieri, M. L. (1933). *Amer. J. Physiol.*, **105**, 146.
 Demole, V., and Ippen, F. (1935). *Hoppe-Seyl. Z.*, **235**, 226.
 Drill, V. A. (1938). *Amer. J. Physiol.*, **122**, 486.
 Epstein, A. A. (1922). *Amer. J. med. Sci.*, **163**, 167.
 Hall, K., and Korenchevsky, V. (1938). *J. Physiol.*, **91**, 365.
 Hanel, P. (1937). *Klin. Wschr.*, **16**, 1105.
 Hendricks, W. A. (1933). *Amer. J. Physiol.*, **105**, 678.
 Himwich, H. E., Goldfarb, W., and Cowgill, G. R. (1932). *Ibid.*, **99**, 689.
 Korenchevsky, V., and Ross, M. A. (1940). *British Medical Journal*, **1**, 645.
 — and Hall, K. (1941). *Biochem. J.*, **35**, 726.
 —, Burbank, R. C., and Cohen, J. (1941). *British Medical Journal*, **1**, 396.
 Mellanby, E. (1934). *Nutrition and Disease*, London.
 Mlinko, Z. (1938). *Hoppe-Seyl. Z.*, **256**, 42.
 Mosonyi, J. (1935). *Ibid.*, **237**, 173.
 Nespor, E. (1936). *C. r. Soc. Biol. Paris*, **122**, 427.
 Parhon, C. I., and Werner, G. (1935). *Ibid.*, **118**, 1659.
 Plaut, F., and Bülow, M. (1935). *Klin. Wschr.*, **14**, 1318.
 Sampson, M. M., and Korenchevsky, V. (1932). *Biochem. J.*, **26**, 1322.
 Sure, B., and Buchanan, K. S. (1937). *J. Nutr.*, **13**, 521.
 — and Theis, R. M. (1938). *Proc. Soc. exp. Biol. N.Y.*, **37**, 646.
 Thaddeus, S. (1938). *Arch. exp. Path. Pharmac.*, **190**, 227.
 Wohl, M. G., and Feldman, J. B. (1938). *Trans. 3rd International Golter Conference*, p. 460, Portland, Oregon.

THE INCREASING IMPORTANCE OF ENDOCRINOLOGY IN GENERAL MEDICINE*

BY

S. L. SIMPSON, M.D., F.R.C.P.

Physician, Willesden General Hospital, with charge of Diabetic Clinic; Physician, Princess Louise Kensington Hospital for Children; Consulting Physician, Soho Hospital for Women

In this address I hope to be able to indicate clearly that the "increasing importance of endocrinology in general medicine" is a reality and is of interest to the physician, surgeon, and gynaecologist, and therefore to the general practitioner.

Adrenal-Pituitary Disorders

Cushing's Syndrome.—I shall first consider a syndrome which has been brought into prominence in the last decade by the observation of Harvey Cushing (1932), but which had already been recognized and described as the "fat bearded diabetic woman" by Achard and Thiers. The eponymous title was suggested by Parkes Weber, who had himself previously given a magnificent clinical description of the disorder. Cushing's invaluable contribution was to draw attention to the possibility of a basophil adenoma of the anterior pituitary being a primary cause, comparable to the eosinophil adenoma of acromegaly. It is now recognized that other primary causes exist—namely, tumours of the adrenal cortex, of the ovary, and of the thymus

* Delivered as the Annual Oration to the Leeds Jewish Hospital Medical Society, Nov. 29, 1942.

—but Crooke (1935) has shown that a constant change in all these pathological entities is hyalinization of the basophil cells of the anterior pituitary. Cushing described his syndrome as follows:

"(1) A rapidly acquired, peculiarly disposed, and usually painful adiposity (in one instance representing a 40% gain in weight), confined to the face, neck, and trunk, the extremities being spared; (2) a tendency to become round-shouldered (kyphotic), even to the point of measurable loss of height associated with lumbos-spinal pains; (3) a sexual dystrophy shown by early amenorrhoea in the females and ultimate impotence in the males; (4) an alteration in normal hirsuties shown by a tendency to hypertrichosis of face and trunk in all the females as well as the pre-adolescent males and possibly the reverse in adult males; (5) a dusky or plethoric appearance of the skin with purplish lineae atrophicae; (6) vascular hypertension; (7) a tendency to erythraemia; (8) variable backaches, abdominal pains, fatigability, and ultimate extreme weakness."

This description gives an indication of the protean manifestations of an endocrinopathy, but it is more important to realize that a presenting symptom may overshadow the rest of the picture. Thus the disorder may be diagnosed as essential hypertension, nephritis, malignant nephrosclerosis, cerebral tumour, diabetes mellitus, polycythaemia, idiopathic adiposity, or sterility in the female or impotence in the male. It is also important to remember that an idiopathic adiposity may be the precursor of the complete syndrome, which may manifest itself only at the climacteric or in the course of a pregnancy. Not all patients developing Cushing's syndrome are sterile or without menstrual bleeding, although the latter tends to be scanty (Simpson, 1936).

The differentiation of the pathological varieties of Cushing's syndrome is helped by biological assay, and in 1934 and 1936 my colleagues and I recorded a high androgen excretion in virilism. I suggested the origin of androgens from the adrenal cortex and the significance of urinary 24-hourly values above 300 international units as indicating an adrenal cortical tumour. Work in this field was fruitfully developed by R. K. Callow (1938a, 1938b), with whom I collaborated on the clinical side over a number of years, especially in the demonstration of urinary androgens after bilateral ovariectomy.

Phaeochromocytoma of Adrenal Medulla.—A phaeochromocytomatous tumour of the adrenal medulla consists of adrenaline-secreting chromaffin cells, which discharge into the blood paroxysmally, though possibly also in constant small quantities. Hypertension, with periodic exacerbations to very high levels—e.g., over 300 mm. Hg, with the diastolic pressure rising to a relatively smaller extent—is the major manifestation. These vasomotor crises are due to a discharge of adrenaline into the blood stream; and other features are pallor, coldness and cyanosis of the extremities, profuse perspiration, trembling, agonizing headaches, painful constricting sensations in the chest with difficulty in breathing, angina-like pains, cramps of the calf muscles, mydriasis, albuminuria, and pyrexia. Ischaemic nephritis may also occur. The presenting features of the disorder, however, may be attacks of cyanosis, coughing and blood-tinged sputum, and pulmonary oedema; blurred vision, papilloedema, retinal haemorrhages, exophthalmos, mydriasis, and cephalalgia; epigastric pain, nausea and vomiting; an enlarged thyroid, with tachycardia; albuminuria and hypertension; attacks of coma due to hypertensive encephalopathy; hypersensitivity to heat associated with a considerably raised basal metabolic rate. The following mistaken diagnoses have been made: neurosis, neurocirculatory asthenia, thyrotoxicosis, diabetes mellitus (but severe diabetes mellitus may coexist), essential hypertension, malignant nephrosclerosis, pulmonary oedema, cerebral tumour, acute abdomen, and angina. The chief cause of such mistaken or incomplete diagnoses is the failure to realize that such a syndrome exists.

Sexual Precocity.—It is well known that sexual precocity may be associated with neoplasms of the endocrine glands, especially of the adrenals. It is less well known that the primary lesion may be a cerebral neoplasm in the neighbourhood of the third ventricle, producing its effects via the hypothalamic-pituitary mechanism, and it is probable that the classical sexual precocity associated with pineal tumours is also produced through this mechanism and not through an absence or excess of pineal gland secretion. There are sometimes, although by

no means always, interesting points of clinical differentiation, but the essential point I wish to make is the overlapping of the endocrinological and neurological field. We have a similar overlapping, not well recognized, in the condition of gigantism, which does not necessarily present itself as a pathological entity, but as a variety of normality. Thus recently a mother brought her daughter of 14 to see me because the girl was easily tired and she thought it was due to her growing too rapidly. Examination of the fundi did not show the optic atrophy associated with a pituitary tumour, but the papilloedema of increased intracranial pressure, the latter confirmed by radiographs. The probability is that we are dealing with a third-ventricle tumour, and ventriculograms are being taken for further indications.

Addison's Disease.—This disorder is well known as a clinical entity, and interest has been intensified by the discovery of a potent cortical extract. But the protean manifestations of the disorder, and the fact that it is comparatively rare, lead to mistaken diagnoses, especially in fair people in whom pigmentation is minimal or absent, or in whom it is identical with that of leucoderma. Addison described one such case, and I have had three cases under my care, one ending fatally and showing atrophy of the adrenals at necropsy. I have seen a case of Addison's disease for the first time at necropsy, after the patient had been operated on because of an erroneous diagnosis of perforated gastric ulcer, the original disorder being quite unsuspected, although typical pigmentation in moderate degree was visible at the necropsy. Severe abdominal pain may be a feature of Addison's disease. Asthenia and psychoneurosis are other erroneous diagnoses met with. On the other hand, the label of Addison's disease may be incorrectly given. Thus I have had cases labelled Addison's disease in which the diagnosis was pregnancy with functional vomiting, carcinomatosis, pellagra, anorexia nervosa, and chronic arsenical poisoning. The Cutler or Kepler test of sodium and chlorine concentration in the blood and urine is valuable not so much for confirming Addison's disease as for excluding this diagnosis in a variety of disorders in which it may be suspected, often with some reason.

Simmonds's Cachexia.—The characteristics of this disorder, due to insufficiency of anterior pituitary function, are anorexia, wasting, weakness, hypotension, bradycardia, and loss of primary and secondary sexual qualities. The disorder as a complete syndrome is a rarity, but as an incomplete syndrome following pregnancy, especially with haemorrhage or infection at parturition, it is in my experience by no means uncommon, and is often missed entirely or mistaken for myxoedema. On the other hand, the condition of anorexia nervosa, which is primarily a psychoneurosis but secondarily a vitamin-deficiency and hormone-deficiency disorder, is often mistaken for Simmonds's cachexia. Clinically, the restless energy, the persistence of sexual hair, and a generalized downy growth of hair characterize anorexia nervosa, and further differentiation may be obtainable from urinary androgen assay and insulin-sensitivity curves.

Diabetes Mellitus

I refer only briefly to this condition because it is familiar to all of you, but it illustrates quite forcibly the thesis that endocrine disorders may present themselves as non-endocrine disorders in various branches of medicine. Thus it is now platitudinous to point out that the diabetic may first appear as suffering from boils, carbuncles, general pruritus, pruritus vulvae, dermatitis, xanthomatosis, neuritis, endarteritis with ulceration and gangrene of an extremity, cataract, retinitis, amenorrhoea, impotence, or infantilism, and such widespread manifestations are equally true of many other endocrinopathies.

I would suggest here that diabetes should not be arbitrarily excluded from the field of endocrinology because of the mathematical calculation of dietary values, and that those who have specialized in the field of diabetes could very appropriately extend their interests to the whole field of endocrinology. Not only has Young (1937) stirred the experimental and clinical world by the production of permanent pancreatic diabetes by injections of a pituitary diabetogenic hormone, but it has also been shown (Lukens and Dohan, 1942) that the development of this pancreatic diabetes can be prevented by insulin or diet, and can even be abolished or cured in the early stages. There is little doubt that this mechanism of producing a type

of diabetes operates clinically in man, and it is well known that diabetes may complicate acromegaly, Cushing's syndrome, thyrotoxicosis, and other endocrine disorders. It is less well known that it may coexist with Addison's disease (Simpson, 1932), or that the latter may be superimposed upon pre-existing diabetes. I have seen two such cases, one through the courtesy of Dr. R. D. Lawrence. A small number are recorded in the literature (McCullagh, 1942), and the involvement of the suprarenals may remain unsuspected. Essentially the diabetic tolerance improves and the insulin dosage is progressively lowered, but clinically the patients get thinner and weaker. Pigmentation may not be conspicuous. A study of sodium chloride metabolism confirms the diagnosis.

The Gonads

The field of gynaecology and obstetrics is saturated with, and I think overwhelmed by, the vast amount of animal experimental endocrinology and therapeutic indications deduced therefrom. Several books have appeared on the endocrinological aspects of, or endocrine therapy in, gynaecology. Unfortunately species differences often invalidate application of the results of animal experiments to man, and, on the whole, endocrine therapy in this field has been tantalizingly promising but somewhat disappointing. Potentially it remains a very fruitful source of further study. Neither should one limit attention to the more obviously pathological—such as the endocrine concomitants—or aetiological factors in eclampsia. Physiological aberrants are equally important. Thus the increasing adiposity that follows each pregnancy in those with an unstable endocrine constitution is almost certainly associated with failure of involution of the physiological pituitary and adrenal hyperplasia, and the persistence of adrenal androgen secretion is often indicated by an accompanying mild but definite hirsutism.

The brilliant results of oestrogen therapy in the vasomotor and psychoneurotic disturbances of the climacteric are too well known to bear repetition, but it is not generally realized that such disturbances may persist for ten or more years after the menopause or may return after a long interval of freedom, or may supervene for the first time ten years after the menopause, just as they may occur in the late forties in a woman who has been completely ovariectomized in the second or third decade of life.

In the male the treatment of sterility by gonadotrophic hormone or small doses of androgens is far from satisfactory, but the treatment of hypogonadism in the surgical castrate or eunuchoid is dramatic. Also, excellent physical and mental improvement has occurred in a group of cases showing juvenilism in the presence of apparently normal genital function. Impotence due to organic hypogonadism always responds to androgenic therapy, but psychological impotence calls for psychological treatment.

Parathyroids

Hyperparathyroidism, due usually to an adenoma of one of the parathyroid glands, is primarily an endocrine disorder, but its manifestations are in the main skeletal, the general decalcification and rarefaction of bones resulting in bony deformities, spontaneous fractures, and variable bone pains. Previous to the discovery of the function of the parathyroids and knowledge of the high serum calcium, low serum phosphorus, and negative calcium balance in this condition, the disorder was often confused with osteitis deformans and more localized forms of rarefying osteitis. I have recently seen a case in a woman of 40 who first had a localized bony swelling of the tibia removed, the histological report being osteoclastoma. It was only some months later that the underlying condition was recognized, and a parathyroid tumour sought for, found, and removed. This patient also had multiple renal calculi (calcium deposits)—which reminds one that these patients may first present themselves to the urogenital surgeon with renal colic or haematuria; or, if the calcium deposition is generalized (calcinosis), there may be severe renal insufficiency and uraemia. In such cases the inadequate excretion of phosphorus blurs the chemical picture.

Hypoparathyroidism appears to be more common in Denmark than in this country. It may be due to atrophy or degeneration of the parathyroids or may follow thyroidectomy for thyrotoxicosis in a small percentage of cases, interference with

blood supply being apparently at least as important as inadvertent removal of parathyroid glands. In addition to the characteristic tetany (associated with a low serum calcium and high serum phosphorus) there may be various cerebral and mental disturbances, with epileptiform convulsions, changes in character, hallucinations, and dementia. It is known that patients have been treated as idiopathic epileptics for years (as has been the case with hypoglycaemic attacks resulting from pancreatic islet tumours). Hypoparathyroidism may also present itself in the first place as a trophic disturbance of the skin, hair, nails, teeth, or eye lenses (cataract). Thinning or loss of hair, impetigo herpetiformis, acrodermatitis continua, psoriasis pustulosa, and various ill-defined eczematous lesions have been found associated with hypoparathyroidism, and have apparently responded to calcium and calciferol therapy. Lachmann (1941) states that migraine, cardiac pain with palpitations, and gastro-intestinal spasm severe enough to lead to laparotomy may be presenting features in unsuspected hypoparathyroidism.

Thyroid Gland

The diagnosis of thyrotoxicosis, primary or secondary, is relatively easy, yet it is surprising how often the condition is missed, even when the manifestations are characteristic, and especially when the presenting feature is not obviously endocrine. The cardiologist's awareness of thyrotoxicosis as one cause of tachycardia, auricular fibrillation, or angina-like pain is now taken for granted, and more recently attention has been drawn to the possibility of a small group among hypertensive cases being thyrotoxic in origin. It is perhaps less well realized that anginal pains, cardiac insufficiency, and electrocardiographic changes may have myxoedema as their background. The pulse is usually slow in myxoedema, but may be rapid in the presence of cardiac insufficiency. The capillaries also show characteristic narrowing and absence of tortuosity.

Gastro-intestinal symptoms may be severe in thyrotoxicosis, and I have seen a necropsy on one patient who was operated on for a "query perforated gastric ulcer," the diagnosis at operation having been changed to acute pancreatitis, the pancreas being very congested. I have also seen a necropsy on a patient who was admitted in semi-coma with the diagnosis of "query intracranial haemorrhage, query concussion, with cerebral irritability," when the patient was really in the terminal stages of a thyrotoxicosis crisis. I have seen several cases of "idiopathic wasting, query malignancy" in older people with unsuspected secondary thyrotoxicosis who recovered a few stone in weight following thyroidectomy.

Asthenia is commonly present in thyrotoxicosis, and rarely myasthenia gravis may coexist, but thyrotoxicosis may be wrongly diagnosed as myasthenia gravis or progressive muscular atrophy. Neurocirculatory asthenia may be associated with enlargement of the thyroid gland, but it is coincidental, and an erroneous diagnosis of thyrotoxicosis has led to a useless thyroidectomy.

Psychoneuroses and the Psychoses

Endocrinopathies are more often associated with psychoneuroses than with psychoses, but the latter may coexist with, or be precipitated by, endocrine disorders—e.g., manic-depressive disorders, paranoia, delusional insanity in thyrotoxicosis, myxoedema, chronic hypoparathyroidism, chronic hypoglycaemia, and the climacteric. Hemphill and Reiss (1940) recognize a type of involutional melancholia associated with adiposity and scanty or absent menstruation which they claim responds to oestrogen therapy; and a hypo-ovarian-hypoadrenal type which is said to respond to corticotrophic hormone. A cautious approach to the problem is called for, and a critical evaluation of cause and effect is essential, but it is interesting to note that many years ago Mott drew attention to the degeneration of the testes in dementia praecox.

It is my opinion that all endocrinopathies, constitutional or more obviously pathological, influence behaviour pattern, and that the association of mild or severe psychoneuroses with untreated endocrinopathies is very frequent. This is obvious with thyrotoxicosis and the climacteric, for example, but I have seen a transformation of personality follow adequately treated Addison's disease, eunuchoidism, and even hyperparathyroidism. The important point is that a group of psychoneuroses and psychoses coming to the psychiatrist in

the first case may have an underlying endocrinopathy as an aetiological factor.

Endocrines and Ophthalmology

In 1938 Sir Walter Langdon-Brown gave a lecture to the Oxford Ophthalmological Congress reviewing the ocular manifestations of endocrinopathies as they present themselves to the ophthalmologist. Parathyroid, pineal, pituitary, thyroid, adrenal medullary, gonad, and pancreatic disorders may all show ophthalmological signs, and these may for a time be the presenting feature.

Endocrines and Cancer

Experiments in small animals, such as mice, indicate that oestrogens in large doses over long periods may induce neoplastic growth in genital and other organs. Thus Allen (1942) records cancer of cervix uteri (mice), uterine fibromyomas (guinea-pigs), mammary cancer (mice), interstitial cell tumours of the testis (mice), adenomatous tumours of the pituitary (rats and mice), tumours of the suprarenal cortex (mice), and osteogenetic transformations of bones including tumours (mice). Such neoplastic changes are not found in monkeys, and there is no clinical evidence of the endocrines being responsible for malignant changes in the genital or other organs in man.

Endocrine Therapeutics in Non-endocrine Disorders

Hormones have been used with success in a number of non-endocrine disorders. Thus insulin is anabolic in some thin people, and recently Sargent (1942) has found it useful in a group of neurotics who have lost much weight and have broken down after a long period of stress, showing "a peculiar mixture of anxiety, hysterical, and reactive depressive symptoms." Insulin has also been used in the vomiting of pregnancy and in thyrotoxicosis. Insulin-shock therapy is apparently more beneficial than electric-shock therapy in early schizophrenia. It is claimed that cortical extract has proved of value in surgical shock and shock from burns, the blood chemistry in both instances having some similarity with that of Addison's disease. Apparently the corticosterone carbohydrate influencing factor is more important than the desoxycorticosterone factor. In infections the adrenal cortex loses lipid and undergoes congestion and degeneration. Cortin has therefore been advocated for the acute stages of a variety of infections, especially malignant diphtheria, and for post-infection debility. It is also said to be of value in meningococcal septicaemia of infants, as is illustrated in the Waterhouse-Friderichsen syndrome. Oestradiol appears to be of value in a group of menopausal rheumatoid arthritics. Testosterone I have found of value (Simpson, 1938) for its anabolic nitrogen-retaining effect in hyperthyroidism associated with much wasting. It has also been used for menorrhagia and mastitis. Both testosterone and stilboestrol are effective in suppressing lactation. Adrenaline is of course specific in asthma. Pitressin has been used for chronic pyelitis, and in obstetrics.

Conclusions

I think we must conclude from the above review that the manifestations of endocrine diseases extend throughout the whole domain of medicine, including in that term surgery and gynaecology and the various "special" branches of medicine; that endocrine disorders do not necessarily appear as complete syndromes; and that unless an awareness of the possibilities is present they may remain unrecognized as endocrinopathies when the presenting feature is not obviously endocrine. In the opposite direction one sees a tendency to ascribe the origin of obscure chronic disease, especially of a functional character, to endocrine disorders which are not really present, and the necessity for a knowledge of existing methods of differentiation and the development of more specific chemical, metabolic, and biological tests.

Endocrinology, however, viewed from any angle, is no longer merely a lusty infant. It has grown to maturity, and is at least as well founded on scientific data as many other older accepted specialties. I doubt whether it would gain from the founding of a special hospital for its study—rather would it tend to flourish as special centres or clinics in large hospitals, remembering that such centres for accumulating clinical material from large areas are essential for progress in this field. Such centres or clinics, however, must not be

auxiliary adjuncts in the charge of assistants, created merely to give an air of modernity or progress. Endocrinology is one of the most difficult specialties. Its practice, and even more so its teaching, require wide general clinical experience and judgment, as well as personal experience and continuous familiarity with the enormous and ever-widening field of experimental research and a specialized knowledge of endocrinopathies, which can only be obtained by those who have had special facilities and opportunities for their study. An endocrine clinic should be an important integral part of a hospital, and should serve as a model for modern methods of investigation, research, and therapy in disease.

REFERENCES

- Allen, E. (1942). *Endocrinology*, 30, 942.
 Callow, R. K. (1938a). *Proc. roy. Soc. Med.*, 31, 841.
 — (1938b). *Lancet*, 1, 435.
 Crooke, A. C. (1935). *J. Path. Bact.*, 41, 339.
 Cushing, H. (1932). *Johns Hopk. Hosp. Bull.*, 50, 137.
 Hemphill, R. E., and Reiss, M. (1940). *J. ment. Sci.*, 86, 1065.
 Lachmann, A. (1941). *Acta med. scand.*, Suppl., 121, 1.
 Langdon-Brown, W. (1938). *Trans. ophthalm. Soc.*, 57, 454.
 Lukens, F. D. W., and Dohan, F. C. (1942). *Endocrinology*, 30, 175.
 McCullagh, E. P. (1942). *Clev. clin. Quart.*, 9, 123.
 Sargent, W. (1942). *British Medical Journal*, 2, 574.
 Simpson, S. L. (1932). *Quart. J. Med.*, n.s., 1, 99.
 — (1934). *Proc. roy. Soc. Med.*, 27, 383.
 — (1936). *Trans. med. Soc. Lond.*, 59, 36.
 — (1938). *Post-grad. med. J.*, 14, 144.
 — de Fremerly, P., and Macbeth, A. (1936). *Endocrinology*, 20, 363.
 Young, F. G. (1937). *Lancet*, 2, 372.

USE OF AMPHETAMINAE SULPHAS IN FACILITATING ELECTRICALLY INDUCED CONVULSIONS

BY

K. C. BAILEY, M.D., D.P.M.

Deputy Medical Superintendent, St. James Hospital, Portsmouth

The use of electricity for inducing convulsions in the treatment of mental disorders has revealed a significant factor in connexion with the electrical stimulus. The electrical convulsion threshold has been found to vary, so that not only do successive applications of electricity require increasing strengths of current, but also there is a wide variation in the initial effective amount necessary. In some patients, more noticeably those suffering from certain types of disorder, a very high threshold either obtains from the outset or rapidly develops. In view of the possibility that electricity, particularly in high strengths, may not be altogether harmless, it seems desirable to discover methods by which the electrical convulsion threshold may be lowered. This to some extent may be achieved by increasing the interval between successive treatments, but at least in the cases exhibiting a high initial threshold recourse cannot be had to this method. Moreover, the period of the treatment would be much prolonged. Other ways of achieving the same objective are suggested by the fact that certain drugs have a stimulating effect on the cerebral cortex, and, owing to the results obtained with amphetaminae sulphas (benzedrine sulphate), experiment in the use of this drug was thought likely to be effective, and was therefore undertaken. Before the results of this procedure are examined it is profitable to consider the theoretical basis of this ascertained variation in the electrical convulsion threshold.

So complex has the nomenclature in psychiatry become, and so difficult is it to fit cases into definite disease entities, that it is often more useful to consider symptom-complexes. Particularly is this so in relation to therapeutic measures.

Variations in Cerebral Activity in Psychotic Disorders

Certain types of psychotic disorders exhibit the symptom of diminution of motor activity. This occurs particularly in depression and in schizophrenia. It may be of any degree from lassitude and a disinclination to do anything to apathy and even stupor. When such a prominent feature presents