cause of the heightened threshold in certain types of cases was raised. Assuming that the diminished motor activity associated with depression is due to increased cerebral inhibition, in contrast with that which occurs in schizophrenic disorders, a heightened electrical convulsion threshold might reasonably be expected to result from this inhibition. Now, it is shown by this investigation that cases of depression generally have a higher threshold than cases of other disorders. Investigation also showed that the threshold increased with age, and as depression itself is more commonly found in older age groups there are two possibilities: (a) that certain forms of depression raised the threshold *per se*, or (b) that age alone has this result. It is impossible to say definitely which factor operates the more strongly, or, indeed, whether the age factor is not really the determinant of the psychosis.

Amphetaminæ sulphas, being a known cortical stimulant, was utilized in an attempt to lower the electrical convulsion threshold; in this it was found to be effective, and smaller strengths of electricity could be used.

As regards the mode of action, considerations were discussed showing the likelihood that it exerts its effect centrally rather than by any accompanying peripheral changes, such as the heightened blood pressure. The possibility of the achievement of the central action by a process of "substrate competition," as suggested by Gaddum in the case of ephedrine, was advanced.

In two cases there was some post-convulsion excitement. This was never very great, and may not have been due to the drug, since such excitement has been present after the convulsion when the drug was not employed. No other undesirable effects were observed from the use of the drug: even in the patient with a blood pressure of 210/120 no harm appeared to result with the doses given.

I am indebted to Dr. Thomas Beaton, medical superintendent, St. James Hospital, Portsmouth, for his kind permission to publish this investigation and for his help and interest in it.

REFERENCES

- Beecher, H. K., McDonough, F. K., and Forbes, A. (1938). *J. Neurophysiol.*, **1**, 324.
 Cerletti, U., and Bini, L. (1938). *Boll. Accad. Med. Roma*, **64**, 36.
 Gaddum, J. H. (1938). *British Medical Journal*, **1**, 713.
 Hemphill, R. E., and Walter, W. G. (1941). *Lancet*, **1**, 446.
 Krause, W. (1936). Quoted in *Zbl. Physiologie*, **102**, 38.
 McGregor, J. S. (1941). M.D. thesis.
 McKail, R. A., Obrador, S., and Wilson, W. C. (1941). *J. Physiol.*, **99**, 312.
 Miller, F. R. (1937). *Ibid.*, **91**, 212.
 — Stavraky, G. W., and Wootton, G. A. (1940). *J. Neurophysiol.*, **3**, 131.
 Pavlov, I. (1941). *Lectures on Conditioned Reflexes*, vol. 2, London.
 Walter, W. G. (1942). *J. ment. Dis.*, **88**, 306.
 Williams, D., and Ritchie Russell, W. (1941). *Lancet*, **1**, 476.

ACUTE DISSEMINATED LUPUS ERYTHEMATOSUS: RECOVERY WITH SULPHANILAMIDE

BY

J. E. G. PEARSON, D.M., M.R.C.P.

Hon. Physician to the Connaught Hospital; Assistant Physician
in the E.M.S.

The following case is reported because few records have been found of successful treatment of acute lupus erythematosus with sulphanimide.

Case History

The patient, a married woman, first attended the Middlesex Hospital Dermatological Department in 1933, at the age of 31. She showed lesions of chronic lupus erythematosus on the cheeks, hands, and sternal region which had developed after sunbathing 4½ years previously. During this period she had suffered from toxic hepatitis and anaemia following arsenical treatment for her skin complaint. There was no family or personal history of tuberculosis. Constant supervision and treatment at the Middlesex Hospital included several courses of quinine iodobismuthate intramuscularly, quinine and stovarsol tablets by mouth, and injections of gold thiosulphate, which had to be discontinued owing to a febrile reaction with a rash after the fourth injection. With this treatment her skin condition was kept well under control.

On July 25, 1940, the patient was admitted as an emergency to an E.M.S. hospital. A few days previously she had been

taken suddenly ill with acute dissemination of her rash. On admission her temperature was 100.4°, pulse 90, and respirations 20. There was a diffuse patchy erythema, affecting chiefly the cheeks, nose, upper lip, neck, and limbs, some of the lesions showing scaling; the finger-joints were swollen, stiff, and painful; and there was generalized lymphadenopathy, most marked in the cervical glands. In the first 8 days 27 g. of sulphapyridine were administered, and though she felt better, her temperature, which remained swinging, had risen to 103°, and she had more extensive joint pains. Many original skin lesions had faded, but numerous others had appeared, some being discoid in nature.

On the ninth day a macular rash, becoming confluent in places, started on the neck, thighs, and trunk, and rapidly became diffuse: although it was thought to be an erythematous phase of her lupus, the possibility of drug rash was considered, and as the sulphapyridine had evidently had no beneficial effect it was discontinued. A blood culture taken on the fifth day proved sterile apart from a contaminating *Staph. albus*, and a blood count showed red cells 4.5 millions, Hb 86%, leucocytes 7,500 (neutrophils 63%, lymphocytes 37%).

Her condition rapidly deteriorated, and on the twelfth day she was very drowsy, her cerebraation was slow, and the temperature reached its peak of 105.4° and remained between 104 and 105° for 36 hours. Pulse 110-124; respirations 24-28. She became completely irrational and delirious, and the tip of the spleen was easily palpated. The position was desperate, and it was decided to attack a possible streptococcal infection anew.

A drip blood transfusion of 1 pint initiated intravenous therapy, and during the next 8 days she received intravenously 21 pints of glucose-saline, 35.5 g. of soluseptasine, and 50 c.cm. of pentnucleotide, with daily intramuscular injections of 6,000 units of antistreptococcal serum. She became rational again, but was very weak, and a stomach tube for feeding purposes was necessary for a further 10 days. During the four weeks after the start of her intravenous treatment she received the following: 35 g. of soluseptasine (first week) and 93 g. of sulphanimide (last three weeks); 225 c.cm. of pentnucleotide intravenously or intramuscularly; and 144,000 units of antistreptococcal serum intramuscularly. Apart from a few chronic lesions her rash faded completely; her arthropathy disappeared; her haemoglobin fell from 92% to 57%; daily white cell counts showed an almost constant leucopenia, the lowest being 2,500; and almost all her hair fell out. When a rise of temperature to 101°, with a macular drug rash on the trunk, occurred at the end of this period, a blood transfusion of two pints appeared to do good, and a steady recovery followed, although a mild attack of rubella caused a slight scare. The only possible septic focus discovered during her stay was pyorrhoea affecting five teeth in the lower jaw. Extraction was purposely postponed until July, 1941, when it was performed uneventfully.

In Dec., 1941, a tuberculin patch test proved negative, but within five days she had a recurrence of her arthropathy and lymphadenopathy. These subsided under sulphanimide treatment without any spread of her skin lesions, but further tuberculin tests were considered unjustifiable. Apart from this setback she has remained in apparently good health since her acute attack.

Discussion

Barber (1941) considered that chronic lupus erythematosus can be produced by reaction either to the tubercle bacillus or to a streptococcus; Kelvin (1941) implied that tuberculous toxin can be held responsible for all cases. It is difficult to believe that the latter theory accounts for many of the acute forms of the disease. Sequeira (1927) certainly said that evidence of tuberculosis, including acute miliary and glandular types, could be found in about two-thirds of the patients, but also stated that "the possibility of a non-tuberculous origin of some cases must not be overlooked." Roxburgh (1933) went further in attributing five deaths to streptococcal septicaemia. This termination with or without acute endocarditis has come to be recognized, and although it cannot be taken to prove that the streptococcus is the cause of the whole clinical picture, it is at least suggestive, and is presumably the reason why the sulphonamides were first given a therapeutic trial in this condition. Success has been reported in a small proportion of the patients so treated in America by Weiner (1940), Wile and Holman (1940), and Hopkins (1941), sulphanimide being the compound chiefly used. In four of the fatal cases on which necropsies were performed tuberculosis was conspicuous by its absence, while a variety of acute lesions were encountered, including pericarditis, pleurisy, splenitis, and glomerulo-tubular nephritis.

Although the aetiology of the condition is not clear, further trial of sulphanimide therapy is urged for the acute forms of lupus erythematosus without clinical evidence of tuberculosis. In view of the leucopenia accompanying the condition, the simultaneous administration of pentnucleotide seems to be indicated.

I wish to thank Dr. Henry MacCormac for his encouragement, and for permission to publish this case, for which I am also grateful to Surg. Rear-Admiral Whiteside, medical superintendent.

REFERENCES

- Barber, H. W. (1941). *Brit. J. Derm. Syph.*, **53**, 1, 33.
Hopkins, H. H. (1941). *Arch. Derm. Syph.*, Chicago, **43**, 833.
Kelvin, J. (1941). *Lancet*, **2**, 597.
Roxburgh, A. C. (1933). *Brit. J. Derm. Syph.*, **45**, 95.
Sequeira, J. H. (1927). *Diseases of the Skin*, 4th ed., London.
Weiner, A. L. (1940). *Arch. Derm. Syph.*, Chicago, **41**, 534.
Wile, U. J., and Holman, H. H. (1940). *Ibid.*, **42**, 1059.

Medical Memoranda

An Improved Pappenheim Stain for Detection of Gonococci in Smears of Pus

In 1938 Sandiford described a malachite green-pyronine counterstain for use in Gram's method. He stated that with it pus cells were bluish green, gonococci bright red, and Gram-positive cocci purple-black. It was in fact a modification of the original Pappenheim and Saathof stain (Pappenheim, 1899), which consisted of methyl green, pyronine, ethyl alcohol, phenol, and glycerol. Walton (1936) used the Pappenheim formula recommended by Conn (1929) and got good results in the examination of smears for gonococci; but Sandiford (1937) could not reproduce them, possibly because he used the original Pappenheim formula. Sandiford found that bacteria and cells and their nuclei were not well differentiated on account of a violet coloration, and in 1938 suggested that this coloration was a metachromatic effect due to the presence of methyl violet as an impurity in methyl green. He therefore substituted malachite green.

According to Conn, methyl green is largely used in stain technology for its metachromatic effect, and is actually crystal violet into which a seventh methyl group has been introduced. This seventh methyl group is very loosely attached and thus there is always some methyl violet present, either because it is not all converted into the higher homologue (methyl green) or because it has broken down. In order to obtain pure methyl green Conn suggested that the commercial dye should be shaken up with chloroform, in which substance methyl violet is soluble.

Walton (1938-9) increased the stability of Pappenheim's stain by incorporating in it methyl alcohol instead of ethyl alcohol. He held that methyl alcohol by virtue of its extra CH₃ groups prevented the seventh methyl group of methyl green from dissociating itself.

THE NEW METHOD

In preliminary work at the Liverpool laboratories it was found that Pappenheim's stain deteriorated within a week even after the substitution of methyl alcohol, and gonococci stained more of a violet colour than a red. It was decided, therefore, to shake up methyl green with chloroform before incorporating it in the Pappenheim stain; this resulted in stabilization of the stain. In the method now used a 1% aqueous solution of methyl green is shaken in a separating funnel for a few minutes with about one-third of its volume of chloroform. The chloroform becomes deep violet and is discarded after separation. A modified Pappenheim solution is then prepared which has the following composition:

Methyl green (Gurr), chloroform-treated, 1% ..	100 c.cm.
Methyl alcohol, absolute	10 c.cm.
Pyronine G (Gurr)	0.6 g.
Phenol	1.0 g.
Glycerol	20 c.cm.

The solution is allowed to mature for 24 hours and is then filtered into a dropping-bottle before use. The optimum staining period is 1/2 to 1 minute.

If used as a counterstain in Gram's method for smears of pus, Gram-positive organisms stain normally and gonococci brilliant red. Most other Gram-negative organisms are a less intense red. Nuclei of the leucocytes are bright blue, but the cytoplasm of the polymorphonuclears behaves differently from that of the mononuclears. The cytoplasm of the latter is red, whereas that of the former is unstained. Endothelial cells have blue nuclei and a cytoplasm of a variable pink.

Sandiford found that the phenol in Pappenheim's solution, normally 2 g. in 130 c.cm., tended to decolorize Gram-positive bacteria after more than 2 minutes, and therefore omitted it from his modification. This fact was confirmed by me, and 1 g. of phenol was substituted. With this amount the decolorizing action was minimized, although it was of little importance because the staining

period was much shorter than that used by Sandiford. The staining intensity of Pappenheim's solution is lessened if phenol is omitted.

In a comparison of Sandiford's modification of Pappenheim's stain with that described by me it was found that gonococci were more distinctly red by the latter and easier to detect against the clear unstained background of the polymorphonuclear cells. Moreover, the contrast of the brilliant-red gonococci and the bright-blue nuclei by my modification was greater than that by Sandiford's, with which gonococci were red and the pus cells green with bluish-green nuclei.

The modified Pappenheim stain here described was primarily intended for use in the examination of smears for gonococci, either directly or as a counterstain, but it is very suitable for blood films, particularly in the differentiation of the mononuclear leucocytes. The stain is also excellent for cerebrospinal fluid containing meningococci, and any amount of pyronine from 0.5 to 1 g. will yield good results.

The keeping quality of the modified stain has not yet been determined, but it is longer than 3 months and the solution is effective after Sandiford's has begun to deteriorate.

MAXWELL M. BARRITT, M.Sc., D. Bact., L.M.S.S.A.
Assistant (Acting City) Bacteriologist, Liverpool Corporation.

REFERENCES

- Conn, H. J. (1929). *Biological Stains*, pp. 109, 119, New York.
Pappenheim, A. (1899). *Virchows Arch.*, **157**, 19.
Sandiford, B. R. (1937). *J. Path. Bact.*, **45**, 467.
— (1938). *British Medical Journal*, **1**, 1155.
Walton, S. T. (1936). *J. Bact.*, **31**, 61 (abstract).
— (1938-9). *J. lab. clin. Med.*, **24**, 1308.

The Vitamin C Saturation Test

This test, as modified by Harris and Abbasy (*Lancet*, 1937, **2**, 1429), affords useful and reliable information concerning the vitamin C reserves of men. During my year as lieutenant in the R.A.M.C. I carried it out on 1,200 soldiers according to a scheme drawn up by the Directorate. I performed several thousand titrations, and in the first tour each involved the usual simple calculations, taking into account the period of retention, the volume of the sample of urine, the burette reading, and the variable strength of the reagent—2:6-dichlorophenol-indophenol. These were tedious. The following is the procedure followed in the second tour; it is described here because it saved so much time both in the titrations and calculations.

As before, the men were dosed with 0.75 g. of ascorbic acid in water at 7.30 a.m. They emptied their bladders at 10.30 a.m., and a two-hour period of retention was adhered to rigorously. This is better than three hours, which in winter is too long. The samples were collected at 12.30 p.m. in numbered 2-lb. jam-jars, and instead of measuring the volume, which for 100 samples took about 1½ hours, each sample was made up to 500 or 1,000 ml. with tap-water. While an assistant did this, 10 ml. was pipetted off from each and run into a 6-oz. medicine bottle, numbered on shoulder and on cork. These bottles had previously received 1 ml. of glacial acetic acid.

This method obviates the addition of large quantities (30 to 50 ml.) of reagent in titrating the more concentrated urines. By taking 2.20 ml. of acidified sample for titration (namely, 2 ml. of urine), using a pipette graduated to the tip, with reagent adjusted so that 1 ml. was equivalent to 0.1 mg. of the pure vitamin, convenient readings were obtained. These were around 0.2 ml. for the initial low values of the first day, rising finally to about 6 to 8 ml. Saturation was taken to be when 50 mg. was excreted in a 3-hour period, or correspondingly for two hours. It is convenient to use the numbered corks to identify the 2-ml. portions removed to test-tubes for titration.

The reagent was adjusted to correspond to 0.1 mg. of vitamin, using 100-mg. portions weighed before the tour (not a bought tablet or a 1-g. tube as sold—these are not prepared as standards, and do not justify the implicit faith in their exactness that one sometimes finds). Tap-water had to be used, but this introduced no appreciable error. The standard must be titrated immediately after preparation.

The method appears to be entirely reliable, and will differentiate groups of men living under slightly different conditions. This may best be appreciated by plotting the number of daily doses required to saturate on the vertical axis and the number of men saturated by that day on the horizontal axis; namely, the sum of those saturated on first, second, and third days—say 60%—would be plotted at 60 for third day. Another way of assessing is to plot for each group the number unsaturated on some selected day; this shows up deficiencies. Thus, for example (an entirely hypothetical case), suppose one group of men had 5% remaining unsaturated after six doses and another had 50%, the difference could be strikingly shown in a graph. If, however, the two groups were close together they might best be differentiated by choosing a smaller number of doses for a comparison. Thus two groups which gave the same percentage of unsaturated after, say, five doses might be separated by one of them having a larger number unsaturated after four doses.

Marine Biological Laboratory,
Plymouth.

W. R. G. ATKINS, Sc.D., F.R.S.