Feb. 25.—Considerable vomiting, projectile in nature, without nausea and preceded by bouts of severe headache. C.N.S. completely normal.

Feb. 27.—Lumbar puncture performed and 8 c.cm. of clear fluid withdrawn, not under pressure.

From this date the patient's condition improved rapidly; the vomiting ceased, headache disappeared, and the central nervous signs remained normal. A further lumbar puncture was performed on March 12, 1940. There were no abnormal characteristics about this fluid, and, since the patient was by now completely recovered, he was sent to a convalescent home and thence on sick leave. There were no noticeable sequelae.

TABLE II.—C.S.F. Results in Case II

Date	Protein (mg./100 c.cm.)	Cells per c.mm.	Lymphocytes	Sugar (mg./100 c.cm.)	Chlorides (mg./100 c.cm.)	Organisms	Lange Gold Curve
23/2/40	70	20	++	40	710	Nil T.B. not found	0022221000
27/2/40	40	8	++	—	—	Nil	0011000000

Wassermann reaction negative

The diagnosis of rubella, as in Case I, does not seem open to reasonable doubt. Meningism was not present at any stage, and the central nervous signs and symptoms were those of an encephalitis with resulting increase in cerebral pressure.

Commentary

Complications of any variety during an attack of rubella are very rare, and involvement of the central nervous system is the least common of these. The cases of this type so far reported include manifestations of meningitis, encephalomyelitis, polyneuritis, and ascending paralysis of the Landry variety. At least two of these cases were fatal.

The necropsy on the 18-year-old girl described by Taylor (1937) revealed the typical picture of an acute to subacute encephalomyelitis with demyelination most pronounced in the perivascular regions, as may be found in encephalitis associated with the exanthemata, such as measles, small-pox, and mumps. According to Russell Brain (1940) the cause of this type of encephalitis is still unknown. The former view that it is due to the virus of the exanthem in question is in disfavour owing to failure to reproduce experimentally the pathological picture by means of the virus. An alternative opinion, that the condition is brought about by another virus common to all patients and normally latent but activated by the exanthem, would explain the similarity of pathological features in all cases and also the fact that encephalitis of this nature may arise with apparent spontaneity unpreceded by an infection. No such virus has, however, so far been demonstrated. Finley (1937) suggests that the encephalitis may be an allergic reaction in the brain following the general eruption. This is in accordance with the finding of Rivers and Schwentker (1935) that a demyelinating encephalomyelitis could be produced in monkeys by repeated injection of brain material.

I am grateful to Surg. Cmdr. Mussen, R.N., for bringing Case II to my notice, and for his advice in the compilation of this account; also to Surg. Rear-Admiral Bradbury, R.N., and Surg. Capt. Parker, R.N., for permission to publish it.

REFERENCES

- Brain, W. Russell (1940). *Diseases of the Nervous System*, London.
Finley, K. H. (1937). *Arch. Neurol. Psychiat.*, Chicago, 37, 505.
Rivers, T. M., and Schwentker, F. F. (1935). *J. exp. Med.*, 61, 689.
Taylor, R. J. (1937). *Med. J. Austral.*, 2, 604.

In a discussion on bacteriaemic staphylococcal infection, Champ Lyons (*Surg. Gynec. Obstet.*, 1942, 74, 41) observes that the importance of thrombophlebitis in the production of metastatic abscesses was recognized by John Hunter. He quotes Topley to the effect that in a bacteriaemic staphylococcal infection "the cocci are carried from the original focus of infection in detached fragments of an infected thrombus." Lyons presents evidence in support of this view, and believes that pulmonary thrombophlebitis as a factor in the perpetuation of a staphylococcal bacteraemia may be likened to a left-sided endocarditis.

PLANNING FOR THE TREATMENT OF HEAD INJURIES*

BY

HUGH CAIRNS, D.M., F.R.C.S.

Brigadier, Consultant Neurosurgeon; Nuffield Professor
of Surgery, Oxford

The war has given impetus to the study of head injuries, and information is now available which should assist in the formulation of plans for post-war medical services.

I. Plans should be based on a Survey of Unselected Material

Before the war, with few notable exceptions, specialists responsible for teaching on head injuries saw an undue proportion of the severe cases and those in which compensation or litigation was pending. Although as long ago as 1932 Ritchie Russell had shown that the majority of patients with head injury made a satisfactory return to work within six months, treatment and prognosis of head injuries continued for the most part to be governed by pathological doctrines which implied, or were thought to imply, that bruising of the brain was an almost invariable accompaniment of even minor concussion, and that the bruising took a long time to resolve and might continue to produce symptoms for a number of years. Patients are still encountered who have been confined to bed for weeks or months after relatively trivial head injuries, and have been warned that because of brain damage they cannot expect to resume full normal activity for periods of months or a year.

There is still considerable controversy about the pathological processes in head injury, but in the special centres established since the outbreak of war comprehensive survey of head injuries is accumulating an impressive number of facts which show that the picture of the late results of blunt head injury† has been painted blacker than it really is. Thus, in a military hospital for head injuries, in a series of 237 acute cases of head injury in the fighting Services (not including fatal cases, but otherwise unselected), Symonds and Ritchie Russell (1943) found that 80% were successfully returned to duty, the majority within three months. Among civilians Eric Guttman (1943) obtained similar results in a consecutive series of 158 wage-earners admitted to the Radcliffe Infirmary, Oxford, suffering from head injury: 25% returned to work within three weeks, 50% in between three and nine weeks, and only 8 patients had not returned to work within six months. In Guttman's series there had been an initial hospital mortality of 4%.

These are the results in unselected hospital material from a mixed urban and rural area, at a time of "comparatively standardized and favourable social conditions, when everybody has the choice of well-paid jobs and most employers are ready to facilitate gradual return to work." In civilian head injuries, permanently incapacitating brain damage appears to be an uncommon event—among Guttman's 158 wage-earners there were only 2 patients so affected—and early return to work may be expected in the majority. The results suggest the need for revision of teaching on the subject of head injuries. They also suggest that where problems of compensation arise settlement should be made at an earlier date than has been customary in the past.

II. Where should Head Injury Patients be Treated?

Accident cases are usually taken to the nearest hospital, and it would be difficult to devise machinery for primary segregation of head cases in special centres. Guttman found that 30% of his head-injury patients had major injuries in other parts of the body, such as fractures of long bones, and this fact favours the current practice of sending all accidents to the general hospital. There is little doubt, however, that, in the general hospital, accident cases should be segregated in an accident service, as has been done at Birmingham and Oxford, under the care of surgeons who particularize in accident work and who have special facilities for resuscitation.

Since the majority of head injuries are evidently mild and, when treated correctly, uncomplicated, it would be reasonable

* Based on a paper read to a meeting of consultants of the combined Services at Reading, Nov. 28, 1942.

† Gunshot wounds of the head are not considered in this paper. They constitute a separate problem.

to expect that the *trained* accident surgeon should himself be able to see the majority of his head-injury cases through to the stage of return to full work, with the assistance of the general practitioner, the industrial doctor, and the social worker. The services of the nearest neurosurgeon would be sought in head injuries which were complicated by severe damage to the brain or skull and in those cases which were not recovering according to expectation. When fit to be moved such cases should be transferred to head-injury centres.

III. The Need for Statistics

Planning for head-injury centres (and for accident services) after the war would be assisted by more precise knowledge of the number, type, and distribution of accidents than seems to be available at present.

In Britain, road accidents of all types amount to about 230,000 a year, industrial accidents (exclusive of mines and certain other industries) to about 270,000 a year, and mining accidents to about 133,000 a year (Safety in Mines Research Board, 1937). I have not found any information about the number of accidents which occur at home and at play, but in the U.S.A. in 1940 the deaths from accidents in the home almost equalled those from motor vehicles (National Safety Council, Chicago, 1941).

These figures do not discriminate between the patient requiring in-patient hospital treatment and the man who is so mildly injured that he can be treated at home and is able to return to work within a few days. In 1935, of the 1,300,000 accident cases in this country, only 136,000 were treated as in-patients (Report on Rehabilitation, 1939). The percentage of head injuries may vary from one type of accident to another. Of accident cases admitted to the Radcliffe Infirmary—mainly road accidents—30% showed head injury. On the other hand, in occupational accidents in the U.S.A., assessed from compensation data, head injuries amounted to no more than 7% . in British mines in 1935 the proportion of head injuries was also 7% (Safety in Mines Research Board, 1937).

Clearly, more exact information is wanted, particularly in regard to patients requiring in-patient hospital treatment. It is reasonably certain, however, that there are many more head injuries each year than could be treated by the neurosurgeons. Their function should be, along with neurologists and psychiatrists, to treat the cases presenting special difficulties.

IV. Rehabilitation

The accident surgeon, or someone on his staff, should have a clear understanding of the problems of treatment and rehabilitation of head-injury patients. Operation is often required for the repair of scalp wounds, but for other purposes only in a limited number of cases—according to Rowbotham (1942) in about 5%. Neglect of a meticulous surgical technique in closing the scalp may wreck the result in otherwise favourable cases, as Botterell and Jefferson (1942) have pointed out. In the cases of Symonds and Russell (1943) and Eric Guttman (1943) the methods of treatment were simple and did not include any dehydration. The patient was allowed to get up gradually soon after he had recovered consciousness. He was examined thoroughly to see that there was no residual brain damage, and his personality was assessed by a study of his past, social, and family histories. An important part of the treatment is reassurance of the patient and his relatives. In the cases from the fighting Services, occupational therapy and graduated physical training were employed; these patients had to be rendered fit for full duty in the Army or R.A.F. In the civilian cases, on the other hand, no such measures were used: the patient was discharged to his home soon after he was able to get about, with instructions to report at the out-patient department on a date a little before the time at which it was estimated he might be fit for work again. The patient was advised to take graduated exercise of his own choice, a letter was sent to his family doctor, and he was seen by appointment in the out-patient department from time to time, by those who had looked after him in the acute stage, until he was once more in his stride at work. Difficulties of a psychological or social character which had been elicited by the preliminary study of personality or at the subsequent out-patient visits were dealt with by explanation and reassurance, and by visits of the social worker (who attended all out-patient sessions) to the home or to the man's employers.

Guttman's work suggests that in the majority of head injuries in civilians the main factors in rehabilitation are the doctor's knowledge of the patient and of his social conditions, as well as of his injury, and a sustained interest in him until he is once more back at work. It really amounts to psychological management, but I have avoided using the word "psychological" lest it be thought that this process of rehabilitation requires as a routine the services of the psychological specialist or the use of elaborate psychological techniques. Handling of most patients along common-sense psychological lines should not be beyond the powers of the competent accident surgeon, who starts, moreover, with the initial advantage of having helped the patient through the acute stage of his illness. In limb injuries, though all the emphasis is placed on physical re-education (Report of the Interdepartmental Committee on Rehabilitation, 1943), I suspect that the psychological side is probably just as important as it is after head injuries, and that in protracted cases return to work is as often delayed by psychological factors as by the physical condition of the limb.*

If Guttman's work is confirmed there is clearly no need for institutional rehabilitation in the majority of cases of civilian head injury. Institutional rehabilitation would be required, however, for the severe or recalcitrant cases and for those whose home conditions were unfavourable. The proportion requiring institutional treatment might be expected to vary with the social conditions at the time.

Addendum

Recent planning for rehabilitation of disabled persons raises some doubts as to whether the problem of restoring the sick and injured to full work is clearly understood by the planners. According to the Beveridge Report (Beveridge, 1942), rehabilitation is "a new field" of remedial activity. The Tomlinson Report (Report of the Interdepartmental Committee on Rehabilitation, 1943) states that "rehabilitation in its strictly medical sense means the process of preventing or restoring the loss of muscle tone, restoring the full functions of the limbs, and maintaining the patient's general health and strength. (This is apart from special rehabilitation treatment required for particular diseases such as tuberculosis.)" The report recommends the institution of a diploma "to cover the special needs of rehabilitation." Some readers might conclude from these statements that rehabilitation is a new technique of treatment on physical lines, to be conducted by specialists.

There is nothing new about rehabilitation. For example, it has been practised systematically for over 50 years by those responsible for the care of the blind and to what high standard this work has been raised in this country may be seen in the British Council's fine documentary film *Out of the Night*. In each broad group of maladies the methods of after-care are to a considerable extent specialized. In the case of blindness this is obvious. After limb injuries the doctor who is responsible for getting the patient fit for work must be competent to judge whether the position of the bone and the union of the fracture are satisfactory; and in this group of cases physical re-education is important. After head injuries rehabilitation must always be partly based on thorough neurological examination; physical re-education, as has previously been pointed out (Cairns, 1942a, 1942b), plays a small part.

If there is one thing common to rehabilitation after all varieties of accident and disease it is the need for study and understanding of the personality of the patient and of his social circumstances, and this should be the duty of every doctor who treats a patient. The "new" things in rehabilitation are the more general realization by the medical profession of its responsibilities for convalescence of all patients, and the formal recognition of the part which a patient's personality plays in his readjustment after injury or disease. The importance of the social worker in rehabilitation is also becoming more generally accepted.

* The importance of the psychological factor in limb injuries was clearly recognized by the Delevingne Committee, who wrote (Report of Interdepartmental Committee on Rehabilitation, 1939, p. 52): "The rate and degree of recovery will depend not only on the nature of the treatment given for the physical injury but also on the individual mental 'make-up' of the injured man"; and again, "The witnesses generally were agreed that a psychological factor was present in varying degree in the majority of cases, and that this was often a dominant factor in retarding recovery."

It would be harmful if people came to think that rehabilitation consisted of physiotherapy or some other technical method of treatment, or that it was the particular province of any group of specialists. To set up a specialty of rehabilitation would be akin to establishing specialties for diagnosis or for therapy. In a national scheme all varieties of medical specialist would be involved, and the general practitioner and the industrial doctor would be more essential than any of them. The whole medical profession would be involved. In ancillary services the social worker would be at least as important as the masseur.

Extension of rehabilitation to all sick and injured persons would obviously require a large increase of social workers and doctors, whose training for this work would tend to become standardized. But in each group of diseases or injuries successful rehabilitation depends to some extent on special knowledge of the disabilities produced by the disease or injury in question, and of the reactions of patients to these disabilities. Trained experts would deal with the cases requiring institutional after-care, mostly in special centres, from which advances in knowledge of the social aspects of ill-health might be expected. But, if the experiences with head injuries are any guide, it would seem that much of the after-care of the nation's sick and injured should continue to be directed by the doctor who treats the patient in the acute stage of his illness. If in his student days he can be taught the after-care of patients in a practical way, and if he can be supplied with more precise knowledge of the social prognosis of diseases, and with adequate assistance from social services and from employers, this task should not be beyond him.

REFERENCES

- Beveridge, W. (1942). *Social Insurance and Allied Services*, H.M.S.O.
 Botterell, E. H., and Jefferson, G. (1942). *British Medical Journal*, 1, 781.
 Cairns, H. (1942a). *Proc. roy. Soc. Med.*, 35, 299.
 — (1942b). *Brit. J. phys. Med.*, n.s., 5, 84.
 Guttman, E. (1943). *British Medical Journal*, 1, 94.
 National Safety Council, Chicago (1941). *Accident Facts*.
 Report of Interdepartmental Committee on Rehabilitation of Persons Injured by Accidents (1939). H.M.S.O.
 Report of Interdepartmental Committee on Rehabilitation and Resettlement of Disabled Persons (1943). H.M.S.O.
 Rowbotham, G. F. (1942). *Acute Injuries of the Head*, R. and S. Livingstone, Edinburgh.
 Russell, W. R. (1932). *Brain*, 55, 549.
 Safety in Mines Research Board (1937). *Protective Equipment of Mine-workers*.
 Symonds, C. P., and Russell, W. R. (1943). *Lancet*, 1, 7.

REHABILITATION IN AN E.M.S. ORTHOPAEDIC UNIT

BY

S. ALAN S. MALKIN, M.B., B.S., F.R.C.S.Ed.

*Surgeon-in-Charge, Harlow Wood Orthopaedic Hospital,
Nottingham; Orthopaedic Surgeon, Mansfield and
District General Hospital*

AND

GILBERT PARKER, M.B., Ch.B., F.R.C.S.Ed.

*Hon. Assistant Surgeon, North Riding Infirmary, Middlesbrough;
Acting Assistant Surgeon, Harlow Wood Orthopaedic
Hospital, Nottingham*

The establishment of orthopaedic centres in the Emergency Medical Service stimulated those working in them to tackle the problem of rehabilitation, already carefully investigated by the Delevingne Committee. With few exceptions it had been the custom in civilian practice to discharge a patient when fit for light work and still maintain some supervision of his further recovery as an out-patient. In the case of Service patients (who have formed the majority of those treated in this centre) the disposal was not so easy, for discharge from hospital removed them completely from the supervision of the surgeon. The danger of too early discharge from surgical supervision is easily seen; a typical example will make the point clear. In civilian work it had been our practice to discharge meniscectomy cases from hospital two weeks after operation and to continue treatment as out-patients. Service patients retained for a further fortnight have usually acquired full knee movement, have a fair quadriceps and no effusion, and are able to walk

several miles without undue pain or fatigue. No doubt some of these men, if returned to light duties, would rehabilitate themselves, but many would sustain some minor trauma, develop effusion, and be sent back to hospital some months later with a history of recurrent effusion and marked quadriceps insufficiency. The problem of rehabilitating such patients is then much more difficult than if it had been dealt with in the immediate post-operative period.

Before discharge from hospital a patient should fall into one of the following groups:

(a) Fit to return to full duties in his original category.

(b) Fit to return to duty in a reduced category; usually cases which have progressed satisfactorily, but in which some residual disability remains—e.g., limitation of elbow movement after fractures in the region of the elbow-joint.

(c) Fit to be transferred to a Command convalescent depot and undergo graduated Army training. This includes cases in groups (a) and (b); it also includes patients who have improved beyond the limit of possible relapse but who are not quite fit for duty—e.g., lower-limb fracture patients who can walk and run easily and require only time and graduated use to complete their full recovery.

(d) Patients who are unfit for Service duties and are discharged from the Service. These fall into three groups: (i) Patients fit to return to their full pre-Service occupation; (ii) patients unfit to follow their pre-Service occupations, but fit to undergo training at a Ministry of Labour training establishment; (iii) severely disabled patients who require vocational training.

The majority of the cases so far treated have been in group (a), (b), or (c).

Organization

When the need for a rehabilitation annexe was appreciated we were fortunate in securing for our purpose a country house, within reasonable distance of the hospital yet far enough away to make a definite break in hospitalization. The house, capable of accommodating 50 patients, is situated in ample grounds, with two large lawns suitable for outdoor games and exercises, a large indoor riding school (now converted into a gymnasium), and an open-air swimming pool. Outbuildings have been converted into recreation rooms. As the work progressed and the value of removing ambulatory cases from a hospital atmosphere became obvious, the original annexe was found to be too small, and near-by B.R.C.S. auxiliary hospitals were utilized as subsidiary rehabilitation centres.

The rehabilitation annexe is staffed by a resident masseuse and P.T. instructor. Each auxiliary hospital has an Army P.T. instructor attached. An occupational-therapy hut has been established at the parent hospital and liaison with the annexes is made through regular visits of one of the staff.

Auxiliary Hospitals

The relation between the parent hospital and the auxiliary hospitals, and the place of the latter in the scheme of rehabilitation, will best be appreciated by following the progress of patients with fractures of the tibia and fibula from the hospital, through the auxiliary hospital to the rehabilitation annexe.

Rehabilitation, it has been stated, begins not in a special centre but in a hospital bed, and ends with a return to full pre-accident activity. All bed patients perform general and special bed exercises regularly. For those confined to bed with leg injuries occupational therapy of a remedial nature is not possible, but diversional work is started in bed. We are fortunate in the situation of our hospital in pleasant country surroundings; beds are pushed out of doors in good weather, and even if the patient's stay in bed is prolonged there is no need for him to see "only the stone walls of hospital wards" (Watson-Jones, 1942).

As soon as leg-plaster cases are considered suitable for walking a sole-plate is applied to the plaster. This consists of a wooden plate with heel and metatarsal bar, the heel and bar being shod with rubber. We prefer this to the Böhler iron, as a heel-and-toe gait (so rarely attained with a Böhler iron) is easily possible with this sole-plate. Patients walk first with the aid of crutches and then with two sticks. Later they may discard their sticks; but we are in no hurry for them to do this, and prefer that they should retain the sticks and walk with a normal gait rather than that they should discard them and