acute myeloid leukaemia (1), tuberculous meningitis (1). Table VIII shows the causes of death in the other cases, and, where applicable, the percentage mortality of the disease.

TABLE VIII.—Causes of 25 Deaths

Diphtheria (faucial)	4 (3%)
Diphtheria (cutaneous)	1
Acute primary pneumonia	4 (2.4%)
Malaria—blackwater fever	1 (0.13%)
Pulmonary tuberculosis	2
Bacillary dysentery	2 (0.10%)
Enteric group	6 (25%)
Acute infective hepatitis	1 (0.3%)
Streptococcal meningitis	1
Heat exhaustion	1
Encephalitis	1
Staphylococcal septicaemia	1

Conclusion

I have tried to condense eighteen months' work into a few thousand words, and have had, perforce, to appeal to figures and tables—always dull things; it is hoped that the main problems of Middle East medicine have been made clear.

We have not been working under conditions to which we were accustomed in peacetime. This hospital is built on desert which was uninhabitable three years ago; every possible amenity has been provided and we have ample safe water, a good hot-water system to the wards, electric lighting from a mains supply, and every reasonable drug and instrument. Three-quarters of the patients have been treated in tents and only the minority in the few hatted wards, conditions which would be impossible in England but are by no means ill adapted to this climate. I feel that the results of treatment could hardly have been bettered under our peacetime civilian conditions.

To practise in a country where tropical disease is rife is to get a glimpse into the future which is opening out before the advances of chemotherapy. While much of tropical specific therapy is old, much is new, and it has been a unique experience to treat bacillary dysentery with the newer sulphonamides. If from the cases treated here the ones amenable to specific chemotherapy are excluded—malaria, amoebiasis, bilharziasis, kala-azar, the avitaminoses, pneumonia, cerebrospinal meningitis, tonsillitis, and bacillary dysentery—there are only small groups of cases still remaining refractory. There is no specific treatment for enteric fever, or such virus diseases as acute infective hepatitis, coryza, and encephalitis or poliomyelitis. Sandfly fever is not important from the point of view of treatment, as it is so brief. Finally, I would again draw attention to the adverse effect of this climate on chronic chest diseases.

SPECIFIC IDENTIFICATION OF THE CHIEF PATHOGENIC CLOSTRIDIA OF GAS GANGRENE

BY

G. F. PETRIE, M.D.

AND

DOROTHY STEABEN, B.Sc.

The Lister Institute, Elstree, Herts

The isolation in a pure state and the identification of the chief anaerobes of gas gangrene—namely, *Cl. welchii*, *Cl. septicum*, and *Cl. oedematiens*—are everywhere recognized as being beset with peculiar difficulties, owing partly to the need of these bacteria for oxygen-free conditions of cultivation, and partly to their tendency to grow in cryptic association with other anaerobes in spite of prolonged efforts to obtain pure cultures. The method described below is likely to aid in their identification, since it depends upon the highly specific, lethal, and antigenic toxin which is characteristic of each of them. This method consists in growing the bacterial inoculum on the surface of agar to which an appropriate amount of one of the specific antitoxins has been added. During the growth of the colony toxin is produced, diffuses outwards from the

colony, and interacts with the homologous antitoxin to form a delicate but unmistakable precipitate resembling a halo round the colony.*

The method is identical with that which was found by one of us (Petrie, 1932) to serve as a differential test for rough and smooth strains of meningococcus, pneumococcus, and *B. dysenteriae* (Shiga) and to establish a clear distinction between group 1 strains (type 1 and type 3) and group 2 strains of meningococcus (cf. Branham, 1940). When, for example, a smooth strain of group 1 meningococcus is grown on an agar plate containing 5% of the homologous immune serum a precipitate is formed round the colonies which is composed of the polysaccharide complex (the somatic antigen) in union with the protein of the specific antibody, whereas in the anaerobic group under discussion the precipitate consists, in the final analysis, of a purely protein aggregate—namely, the toxin molecule in combination with a variable number of the specific molecules of the homologous antitoxin. Molecular aggregates of each kind, irrespective of the chemical nature of the reacting substances, become linked together to appear as a macroscopic precipitate in agar, as described below; in a fluid medium the product of the reaction takes the form of floccules. Soon after the publication of the earlier paper the author made a limited number of observations on the effect of seeding the Park-Williams No. 8 strain of *C. diphtheriae* on agar plates containing diphtheria antitoxin, when ring-like precipitates were noted round the colonies.

Technical Details of the Method

The method is essentially a simple one, and any difficulties that may be encountered are likely to be due to the strict necessity for providing a high degree of anaerobiosis in order to obtain adequate surface growths on agar plates; this proviso applies particularly to *Cl. oedematiens*.

The medium used is 2.5% agar containing the ingredients that are added for the routine production in this Institute of high-grade toxins of the three anaerobes. The basal medium is a horse-meat infusion broth (1 lb. to 1 litre of tap-water heated to 95° C. with 0.5% acetic acid, neutralized and filtered) together with 3.5% Evans peptone and 0.5% sodium chloride; pH adjusted to 7.8, 2.5% agar added, and the medium tubed in 17-cm. amounts and steamed or autoclaved. Before pouring the melted agar the following ingredients are added to it in sterile solution when the temperature has fallen to about 45° C.

For *Cl. welchii*: 2 c.cm. of a mixture containing 5 g. glucose in 100 c.cm. of a sulphated horse-muscle extract (Macfarlane and Knight, 1941), bringing the final concentration of the extract in the medium to 10% and of the glucose to 0.5%. For *Cl. oedematiens*: 2 c.cm. of the sulphated horse-muscle extract, but without glucose.† For *Cl. septicum*: 2 c.cm. of a mixture containing 5 g. glucose in 100 c.cm. heated normal horse serum (56° C. one hour) so that the final concentration of serum in the medium is 10% and of the glucose 0.5%.

Lastly, there is added at the same time 1 c.cm. of a dilution in broth of the appropriate antitoxin, which, when added to 17 c.cm. of agar and 2 c.cm. of extract or serum mixture, represents approximately 8 I.U. per c.cm. of the final medium. The special ingredients can be mixed to provide a sterile solution, and it is doubtless an advantage to add to each mixture an amount of *l*-ascorbic acid, which favours anaerobic growth by virtue of its being a reducing substance, to give a concentration of 0.025% in the final medium.‡

After the plates are poured they should be dried in an incubator at 37° C. for at least 5 hours in order to prevent motile organisms from filming over the surface. The plates may be spread in the usual way, or a heavy inoculation with a platinum loopful of a pure or reasonably pure culture may be implanted on the agar at any chosen site. The plates are examined after 24 or 48 hours' incubation at 37° C. anaerobically. If the resulting colonies are well grown but show no precipitate the plates should be placed at room tem-

* The Nagler reaction resulting from the diffusion of toxin from colonies of *Cl. welchii* on human serum agar plates is used to identify this bacterium (Hayward, 1941).

† There is reason to believe that for both *Cl. welchii* and *Cl. oedematiens* equally good results can be obtained by omitting the sulphated horse-muscle extract from the medium; the agar to be tubed should then be increased from 17 c.cm. to 19 c.cm.

‡ The special ingredients and antitoxin solutions should be kept preferably in a refrigerator, or otherwise in the coolest place available, and great care should be taken to preserve their sterility. Assistance will gladly be given in the technical application of the method if inquiries are addressed to the Lister Institute, Elstree, Herts.

perature aerobically and observed daily, because the haloes, especially those of *Cl. oedematiens*, may be slow in appearing.

Characteristic Features of the Precipitates

There is usually no difficulty in recognizing the halo against a black background with ordinary good daylight, but a simple magnifying-glass or even a binocular dissecting microscope of low magnification is helpful on occasion. At an early stage, when the zone of precipitate is narrow and of slight density, or when it is desired to light up the structural details of the halo to the best advantage, a simple improvised illuminator can be used. This consists of a black plyboard, $1 \times 1\frac{1}{2}$ ft., with a back-rest and a central opening about one inch in diameter, which may be fitted with an iris diaphragm; an electric light provided with a white reflecting shade is placed behind the opening in a position which allows only the reflected light to illuminate the desired portion of the plate. The best view is obtained by ensuring that the external surface of the agar plate is clean and by removing the cover.

When the precipitate has been laid down as an apparently uniform deposit and is left aerobically at room temperature, concentric (Liesegang) rings often appear. These are so characteristic that the specific nature of the precipitate is established beyond doubt (Figs. 1, 2, and 3). They form much more

or group 1 meningococcus is incorporated in the agar (Petrie, 1932). A striking feature of the rings in the anaerobic haloes is the fact that, though the margin of the colony may be

FIG. 3.—Shows rings of specific precipitate round a colony of *Cl. oedematiens* implanted on an agar plate containing *Cl. oedematiens* antitoxin.

FIG. 1.—Shows rings of specific precipitate round the colonies of *Cl. welchii* on an agar plate containing *Cl. welchii* antitoxin.

FIG. 2.—Shows rings of specific precipitate round the colonies of *Cl. septicum* on an agar plate containing *Cl. septicum* antitoxin.

quickly and more readily than the similar rings already described and illustrated in the haloes that are formed when an antibacterial serum against smooth strains of pneumococcus

wavy or irregular, the circular zones give the impression of having been drawn with compasses. Two or three colonies on a spread plate may be seen to be enclosed within a double ring, or a bar of growth may show a double contour in the precipitate extending from its margin.

The breadth and also the density of the rings in a particular halo often vary. Thus one of the best specimens so far obtained was observed after three weeks at room temperature round a colony (8×8 mm.) of *Cl. oedematiens* on agar which contained 75 I.U. of *Cl. oedematiens* antitoxin per c.cm. of the medium. Here there were four rings of precipitate separated by four clear zones. The innermost zone of precipitate was close to the colony but separated from it by a narrow clear space: this zone was narrow (0.5 mm. across) and of such a density as to appear white; then followed a clear zone of 0.5 mm.; then a zone of precipitate of 1 mm., showing a central groove as if about to split and of moderate density; then a clear zone of 1 mm.; next, a zone of precipitate of 1 mm. and of slight density; next, a clear zone of 2 mm.; and, lastly, a zone of precipitate of 1.5 mm. and with the least density of all. The measurements are approximate. A similar picture was given by a colony of *Cl. tetani* (type 1) when grown on a medium containing tetanus antitoxin.

The concentration of the antitoxin, expressed as I.U. per c.cm. of the medium, influences the extent and the density of the precipitate at an early stage of its development. Thus both an excess and also a suboptimal concentration of antitoxin may inhibit the reaction, in accordance with the fact that an excess of either of the reacting substances in the ordinary flocculation test brings about solution of the floccules. This no doubt explains why a narrow clear zone immediately adjoining the colony margin invariably intervenes between the colony and the precipitate, since the concentration of the toxin per unit-volume of medium is presumably highest in this region.

The most likely explanation, in our view, for the appearance of concentric zones is that the forces which combine to build up the framework of molecular aggregates as a uniform deposit to the naked eye are not in equilibrium, and that at a later stage stresses or strains in the structure result in condensation of the precipitate in the form of annular zones separated by clear zones. The remarkable symmetry of the concentric zones is evidence that the molecular aggregates which constitute the toxin-antitoxin precipitate within the semi-rigid matrix of the agar gel are disposed in an ordered arrangement and that the forces controlling them are in equilibrium. When the precipitate shows little tendency, even after a prolonged interval, to form multiple annular zones—as, for example, the diphtheria toxin-antitoxin halo—the density seems to increase from the

margin of the colony to the outer margin of the halo, which appears as a dense ring and which may be surrounded by an extremely tenuous precipitate of considerable breadth, detectable when photographed but not apparent to the naked eye. This variety in its final form also gives the impression of representing a stage of stable equilibrium. The technique may prove to be useful for the study of periodic structures in general and of the physico-chemical basis of the toxin-antitoxin reaction.

Optimal Conditions for Production of Specific Haloes

The need for using a medium which is favourable for toxin production has been indicated by finding that ordinary agar and peptic digest agar are less suitable than the media recommended here. A natural (unconcentrated) antitoxic serum appears to give better results than a salt-concentrated antitoxin or a purified antitoxin as obtained by the pepsin process. The concentration of the antitoxin in the medium, as already stated, is an important factor. In our experience an amount equivalent to 4 to 8 I.U. per c.cm. of the medium gives good results for each of the three strains of anaerobe tested by us, but a smaller amount may prove to be better adapted to less toxigenic strains met with in field work.

When the plates are removed from the anaerobic jar after 24 or 48 hours they may show no sign of a precipitate, but if kept at room temperature aerobically a halo will appear and Liesegang rings may develop. This observation merely proves that toxin had diffused from the colony before its interaction with antitoxin had become manifest as a precipitate. The presence of formalin vapour does not inhibit the development of the reaction.

General Observations on the Method

The ease with which the reaction is obtained is fortunately in the order of the relative frequency and importance, as causal agents of gas gangrene, of the three anaerobes—namely, *Cl. welchii*, *Cl. septicum*, and *Cl. oedematiens*—and may depend upon the rate and vigour of the surface growth of each species on agar plates. The precipitate formed by *Cl. welchii* can be seen after 24 hours' anaerobic incubation. Until experience of the method has been gained it is advisable to keep the plates, after adequate growth has been obtained anaerobically, for a week or 10 days at room temperature aerobically in order to trace the sequence of changes in the haloes.

The method has probably a considerable advantage over those used hitherto, since the association of other bacteria in the colony will not exert a deleterious influence upon the formation of the specific precipitate so long as the toxin of the clostridium remains unaffected by metabolites liberated by the contaminants. As is well known, much of the confusion in defining and specifying the characters of the pathogenic anaerobes by determining their biochemical activities was caused in the past by the presence of unsuspected bacterial symbionts in cultures that were apparently pure. Obviously the method will fail when non-toxigenic colonies or strains appear on the plates, but from a practical point of view in relation to gas gangrene this result is not of serious moment, and is indeed an advantage.

Alternative Applications of the Method

The reaction has also been obtained in a column of agar in a test-tube inoculated on the surface with a culture of *Cl. welchii*; the homologous antitoxin had been added to the agar. The precipitate showed three zones, one above the other, to a depth of 4 mm. below the surface. There can be no doubt that a Seitz filtrate of a broth toxin prepared from the anaerobes in question will give a specific precipitate when superposed on a column of agar containing the corresponding antitoxin. When the toxin molecules have diffused into the agar, precipitation will take place either uniformly or in zones (Brown, 1940).

When the requisite apparatus for obtaining surface growths on agar plates in strictly anaerobic conditions is not available, a modified technique of this kind is worth a trial for detecting toxins in mixed cultures, and we suggest the following procedure.

Employ suitable selective methods to encourage the growth of each of the anaerobes sought for in the primary material, such as wound exudate. Subculture into a separate meat-broth tube or small flask of the Robertson type, but modified, as indicated above, to favour toxin production of the corresponding anaerobe. Grow for

48 hours, filter through a Seitz disk, pipette the filtrate of each culture in a suitable amount on the top of a column of 2.5% saline agar containing 8 I.U. of the corresponding antitoxin per c.cm. of the medium, keep the preparation for one day or more at room temperature, and note whether a precipitate appears in due course. The addition of ascorbic acid to the fluid media during the enrichment process will favour both growth and toxin production.

This simplified procedure may not, however, always succeed, owing to destruction of the specific toxin by proteolysis, or from other adverse conditions induced by the accompanying contaminants. There has, as yet, been no opportunity of testing whether the method is a practicable one for determining the presence of one or more of the pathogenic anaerobes in the exudates of gas gangrene.

When the usual methods of obtaining effective anaerobiosis cannot be applied, the device recommended by Cruickshank and Macdonald (1942) may be adapted to our method and given a trial. These authors grow the inoculum on the surface of an ascorbic-acid-agar medium, superpose a sterile cellophane disk, and pour a layer of the same agar over the cellophane. The original account should be consulted for further details.

Specificity of the Reaction

There can be little doubt that the reaction is a highly specific one. Controls with normal horse serum and with the heterologous antitoxins have been negative. With our present knowledge of these toxins it can be asserted with confidence that when the test is positive the identification of the bacterium needs no support from any other differential character, such as its proteolytic or saccharolytic activities or its pathogenic action in susceptible experimental animals.

The Method as Applied to Investigation of Anaerobic Flora of Wounds and to Diagnosis of Gas Gangrene

The observations recorded here are based on one toxigenic strain of each of the three clostridia—namely, those which are used in this Institute for the routine production of the toxins intended for the immunization of horses. A limited number of comparative observations on halo production by the highly toxigenic Park-Williams No. 8 strain of *C. diphtheriae* and by weakly toxigenic strains—one gravis and two mitis cultures—when implanted on the same plate containing diphtheria antitoxin, showed that the gravis and mitis colonies produced haloes, although the precipitate took longer to form than that which was seen round the No. 8 colony; this result may be significant in the present connexion as indicating that the reaction has a fairly elastic range, provided that time is given for it to mature. Moreover, in a series of agar plates containing varying amounts of *oedematiens* antitoxin good haloes appeared within a range of which the extremes represented 7 and 70 I.U. per c.cm. of the medium, although in some of the plates the reaction was considerably delayed. If 8 I.U. of the antitoxin per c.cm. of the medium should prove to be an excessive amount, 4 units, 2 units, or 1 unit or less per c.cm. may readily be substituted by preparing successive dilutions of a portion of the antitoxin solution with an equal volume of sterile broth and adding 1 c.cm. of the dilution so prepared to the melted agar.

We hope that workers in Army and civilian laboratories who have opportunities of examining severe wounds will give the method a trial and thus help in defining its sphere of usefulness.

Summary

A method of identifying the three chief pathogenic clostridia met with in gas gangrene, by means of specific precipitation resulting from a toxin-antitoxin reaction round the colonies on agar plates, has been described. The appearances presented, and particularly the formation of Liesegang rings, are so characteristic as to leave no doubt of the identity of the associated bacterial species.

Our best thanks are due to Mr. J. Smiles of the National Institute for Medical Research for photographing agar plates illustrating the features of the specific precipitates.

REFERENCES

- Branham, S. E. (1940). *Bact. Rev.*, **4**, 59.
Brown, R. (1940). *Proc. Soc. exp. Biol.*, N.Y., **45**, 93.
Cruickshank, J., and Macdonald, A. (1942). *Mthly. Bull. E.P.H. Lab. Service*, Nov., p. 10.
Hayward, Nancy J. (1941). *British Medical Journal*, **1**, 811.
Macfarlane, M. G., and Knight, B. C. J. G. (1941). *Biochem. J.*, **35**, 884.
Petrie, G. F. (1932). *Brit. J. exp. Path.*, **13**, 380.