BRITISH MEDICAL JOURNAL

LONDON SATURDAY MAY 1 1943

THE INVESTIGATION OF HAEMOLYTIC TRANSFUSION REACTIONS*

BY

P. L. MOLLISON, M.B., M.R.C.P.

A haemolytic transfusion reaction may be defined as the occurrence of an abnormal destruction of the red cells of either the donor or the recipient immediately after transfusion. Following a normal compatible transfusion of fresh blood the donor's erythrocytes are destroyed only at a slow rate, approximately 1% of the total being eliminated each day; since this is (presumably) the rate of destruction of the recipient's own erythrocytes no signs of increased blood destruction are observed. When incompatible blood or blood that has been stored for an excessive length of time is transfused, however, the donor's erythrocytes are rapidly destroyed and signs of increased red cell destruction develop. Destruction of some of the recipient's own red cells may follow the transfusion of blood the plasma of which contains high-titre incompatible agglutinins.

When transfusions are given to persons affected with certain varieties of haemolytic anaemia the signs of haemolysis may temporarily become more pronounced. Such reactions may not be due to one of the causes already considered, and may be listed as rare varieties of haemolytic transfusion reactions.

Haemolytic reactions may be mimicked by the injection of haemolysed blood. This, by the above definition, cannot be considered as a true haemolytic reaction, since the destruction of the donor's erythrocytes precedes transfusion. Nevertheless, the signs produced may be indistinguishable from those of true haemolytic reactions. These various types of haemolytic reaction may be indistinguishable clinically. Each may give rise to jaundice, with or without haemoglobinuria, and each, if fatal, may produce the same post-mortem appearances. On the other hand, each variety may pass unnoticed unless special tests are applied. The purpose of this report is to discuss the pathological features of each type and to describe methods of differential diagnosis.

The establishment of the diagnosis falls conveniently into two parts: (1) the demonstration of the occurrence of increased blood destruction; and (2) the demonstration of the cause of the increased destruction.

The Signs of Increased Blood Destruction

The signs produced by increased blood destruction vary according to whether destruction is predominantly intravascular or extravascular (intracellular).

Intravascular Destruction.—Intravascular destruction results in the liberation of haemoglobin into the plasma. Knowledge of the rate of removal of free haemoglobin from the blood stream has been chiefly derived from experiments in which haemoglobin solutions have been injected intravenously into human subjects. The rate of elimination varies with the initial concentration, being more rapid at high concentrations. For instance, when the initial plasma level is approximately 200 to 300 mg. per 100 c.cm. 50% or more may be removed in three hours (Ottenberg and Fox, 1938), although the plasma may not be entirely cleared for from 8 to 12 hours (Ottenberg

and Fox, 1938; Gilligan *et al.*, 1941). When the initial plasma level is only 40 to 60 mg. per 100 c.cm. the plasma may not be cleared for 5 hours.

Haemoglobin liberated intravascularly may be dealt with in three different ways (Fairley, 1940). (1) It may be absorbed by the reticulo-endothelial system and converted to bilirubin, which is then discharged into the blood stream and excreted by the liver. This is probably the only mechanism brought into play when intravascular haemolysis is slight. (2) When intravascular haemolysis is of greater degree part of the haemoglobin is broken down intravascularly and haematin is liberated; this combines with serum albumin to form methaemalbumin, a pigment which may be recognized by spectroscopical examination or, when it is present in only small amounts, by Schumm's test (Fairley, 1940). The rapid intravenous injection of 14 g. or more of haemoglobin (equivalent to the rapid intravascular breakdown of 100 c.cm. of blood) will lead to the formation of sufficient methaemalbumin to produce a positive Schumm reaction, but the rapid intravascular breakdown of two to three times this amount is necessary to produce enough methaemalbumin to be recognized spectroscopically. Methaemalbumin may be detected in the plasma 5 hours after the intravenous injection of haemoglobin, and then persists for 24 hours or more (above data from Fairley, 1940). (3) Some haemoglobin is excreted by the kidney, but only when the concentration of haemoglobin in the plasma exceeds approximately 135 to 180 mg. per 100 c.cm. (Gilligan *et al.*). This concentration may be reached after the rapid intravascular breakdown of approximately 60 c.cm. of blood. Patients who are already passing albumin in the urine may excrete haemoglobin at a lower threshold (Gilligan *et al.*). Once haemoglobinuria has begun it may continue when the plasma haemoglobin level has fallen far below the initial threshold, so that haemoglobinuria may persist for 24 hours or more after transfusion.

When the urine is alkaline the majority of the haemoglobin will be present in the form of oxyhaemoglobin and the urine will be red in colour. If the urine is neutral or acid, however, methaemoglobin will be formed and the colour will be brown or black. Moreover, acid haematin will be precipitated in the kidney tubules, and small casts of this substance will be found in the urine. The latter may persist for several days after transfusion.

Haemoglobinuria is always accompanied by albuminuria (Gilligan *et al.*), and when haemoglobinuria follows an incompatible transfusion it may also be accompanied by glycosuria, as shown in animal experiments by Ottenberg and Thalheimer (1915) and as exemplified by Case 6 below.

Following intravascular haemolysis, some degree of renal failure may occur, with oliguria and a progressive increase in the blood urea concentration. This renal insufficiency is thought by some authorities to be due to blockage of the renal tubules with acid haematin, a relatively insoluble compound of haemoglobin which is formed in acid urine. This theory was originally based on the observations of Baker and Dodds (1925), who showed that in rabbits renal failure after the injection of haemoglobin solutions could be prevented by alkalinizing the urine. The theory has been challenged by De Navasquez (1940), who believes the anuria is the result of a decrease in the rate of blood flow through the glomeruli. DeGowin *et al.* (1938)

* A report to the Medical Research Council from the S.W. London Blood Supply Depot.

have stated their belief that, in man, blockage of the tubules is insufficient in many cases to account for the degree of renal failure, and they believe that damage to the tubules by some "nephrotoxic factor" may be responsible.

When renal damage occurs signs of uraemia may not be apparent until at least the second or third day after transfusion, and meanwhile the recipient may appear well. Thereafter a critical phase occurs, usually between the sixth and twelfth days (Bordley, 1931), in which the patient may become comatose and die or else start passing large quantities of urine and recover.

Extravascular Destruction.—The extravascular destruction of red cells results in the production of bilirubin. The degree of bilirubinaemia depends upon the rate of blood destruction and also upon the efficiency of the liver in removing bilirubin from the blood stream.

Signs Common to Blood Destruction, whatever the Mechanism

When, as the result of intravascular or extravascular destruction, the serum bilirubin concentration exceeds 4 mg. per 100 c.cm. (approximately) the recipient becomes jaundiced. Although the serum bilirubin concentration rises rapidly after red cell destruction, usually reaching a peak 5 to 6 hours later (Vaughan, 1942), clinical jaundice is unlikely to occur before the end of 12 hours or so, because time is required for staining of the tissues. As mentioned above, the degree of bilirubinaemia depends upon the rate of blood destruction, so that slow blood destruction is less likely to produce jaundice, other factors being equal. It may also be mentioned that the other main factor affecting the onset of jaundice—namely, the adequacy or otherwise of liver function—may be affected by the transfusion of incompatible blood. Following such a transfusion, evidence of liver damage is sometimes apparent at necropsy and may be manifested during life by the development of signs of toxic jaundice after transfusion. For instance, the van den Bergh reaction may become "direct" (see Case 5) and bile pigments may appear in the urine (see Case 6). Normally, once the destruction of the blood is complete the serum bilirubin concentration falls again, and has usually returned to the pre-transfusion level by the end of three days, and jaundice of the patient seldom lasts for more than two or three days after transfusion.

Because of the blood destruction, the increase in haemoglobin concentration expected as a result of transfusion does not occur when there is a haemolytic reaction. Evidence based on haemoglobin estimations before and after transfusion may be misleading, however, because the patient's haemoglobin concentration may be changing at the time of transfusion (due to the nature of the anaemia, for instance) or the patient's blood volume may undergo alteration. For this reason far more reliable evidence may be obtained in the majority of cases by estimating the survival of the transfused cells by the differential agglutination technique (see below).

Apart from the signs of increased blood destruction, the recipient may show signs of constitutional disturbance and complain of symptoms. For instance, a rise in temperature with or without a rigor is common and the transfusion of incompatible blood may be followed by particular symptoms, which are referred to below. It is important to realize, however, that a haemolytic transfusion reaction may occur without giving rise to any symptoms; and, conversely, it must be emphasized that in the majority of cases the occurrence of a rigor following transfusion does not indicate blood destruction.

Although it is possible to make a theoretical distinction between extravascular and intravascular destruction it is doubtful whether the distinction is a valid one. It may be that destruction is always intravascular, but that when it occurs slowly the free haemoglobin in the plasma never reaches a detectable concentration because it is rapidly absorbed by the reticulo-endothelial system, probably at the site of liberation. That this is so is suggested by the observation that in mild haemolytic anaemias no free haemoglobin can be detected in the plasma as a rule, although in an exacerbation haemoglobinaemia and haemoglobinuria may be noted. It is doubtful whether this represents a change in the mode of destruction, and it seems more likely that it merely indicates an increase in the rate of destruction.

In describing the signs characteristic of various types of haemolytic reactions it should be borne in mind, therefore, that when the signs of extravascular destruction occur the inference is probably simply that destruction has taken place at a slower rate than when the signs of intravascular destruction are noted. There are, however, other features of the various types of haemolytic reaction which are of some assistance in arriving at a correct diagnosis, and these different types will therefore be described.

Destruction of Incompatible Blood

Incompatible red cells may be either agglutinated or haemolysed by the recipient's plasma *in vitro*. Only about 30% of sera will haemolyse as well as agglutinate cells containing the corresponding agglutinin (Wiener, 1939), but it is important to realize that even when only agglutination occurs *in vitro* intravascular haemolysis may occur *in vivo*. For instance, when Rh-positive blood is transfused to a recipient whose serum contains anti-Rh agglutinins severe intravascular haemolysis may follow (see Case 6, below, and cases described by Wiener and Peters, 1940), although haemolysis never occurs *in vitro* when Rh-positive cells are incubated with sera containing anti-Rh antibodies. The chief factors affecting the mode of destruction would appear to be the titre of the incompatible agglutinins in the recipient's plasma and the rate of transfusion.

In practice intravascular haemolysis is common after incompatible transfusions, although not invariable (see Case 5, for instance). When intravascular haemolysis occurs it is often followed by some degree of renal failure.

After the transfusion of incompatible blood, clumps of agglutinated donor cells may be found in blood samples withdrawn from the recipient. These agglutinates may persist for at least 48 hours after transfusion (as in Case 4), although they may not be detectable in a sample taken immediately after transfusion (as in Case 5). It may be added that whenever there are preformed agglutinins incompatible with the donor blood—that is to say, when the anti-A or anti-B agglutinins are involved or when atypical incompatible agglutinins (usually anti-Rh) are present in the recipient's plasma at the time of transfusion (as a result of previous stimulation by pregnancies or transfusions)—the majority of the donor cells are usually eliminated within a few hours of transfusion. In such cases the agglutinins in the recipient's serum undergo characteristic changes in titre (Rø, 1937; Wiener *et al.*, 1941; Mollison and Young, 1941). There is usually an initial reduction in titre, lasting for one to two days after transfusion, which is probably due to absorption by the incompatible erythrocytes. Then there is a rapid increase, which usually reaches its peak 10 to 20 days after transfusion. Observation of these changes may present valuable evidence, particularly when the donor cannot be traced. When the recipient's plasma does not actually contain incompatible agglutinins but there is potential incompatibility, as when Rh-positive blood is transfused to an Rh-negative recipient, the rate of elimination is variable. If the recipient fails to react to the stimulus of the foreign agglutinin contained in the donor blood the latter may survive perfectly normally in the recipient. Even after a first transfusion, however, in a person not previously immunized, the recipient may respond by producing anti-Rh agglutinins (although slowly), so that the total survival of the donor cells is reduced from the normal 100 to 120 days to about 60 days (Dacie and Mollison, 1943). In such a case it is of course impossible to detect signs of increased pigment metabolism, and the only evidence of increased blood destruction is the diminished survival time of the transfused cells. In cases in which the recipient has been sensitized by previous transfusions of Rh-positive blood he may respond to a further transfusion by a rapid production of anti-Rh agglutinins and the rapid elimination of the donor cells (i.e., within five to seven days of transfusion). In such a case an increase in serum bilirubin may be detected and the response to transfusion from the point of view of increase in haemoglobin concentration is disappointing (see Wiener, 1941, and Case 1 described by Boorman, Dodd, and Mollison, 1942). Although any case in which the survival of the donor cells is diminished may be considered to be an instance of a haemolytic reaction, the term is in practice restricted to those cases in which destruction starts during transfusion. The term "inapparent haemolysis" has been applied by Wiener to cases in which destruction occurs more slowly after transfusion.

Immediately after the start of a transfusion of incompatible blood the patient may complain of severe lumbar pain (see Cases 1, 2, and 3, for instance), together with various other symptoms, such as burning of the face, tightness of the chest, etc.; some degree of collapse followed by a rigor is common. These symptoms may be delayed, however (see Cases 5 and 6), or abolished by morphine (see Case 3), or may be entirely absent (Case 4). Conversely, they may be produced by the transfusion of apparently compatible serum. One has the impression that the sequelae of the transfusion of incompatible blood are less severe now than formerly, and, if true, this may be due to the increasing tendency to administer blood at a drip rate (apart from cases of acute haemorrhage and "shock"), compared with the more rapid rate that used to be adopted.

Destruction of Stored Blood

(Data from Mollison and Young, 1942)

The rate of destruction of the erythrocytes of stored blood in the circulation of a normal recipient after transfusion depends mainly upon their previous length of storage, the preservative solution used, and the temperature of storage. The erythrocytes of blood stored with a citrate-glucose diluent (M.R.C. War Memorandum No. 1, 1940) at 5° C. for periods up to 10 days or so survive very well in the recipient's circulation, only a very small proportion being broken down during the 24 hours following transfusion and the total survival time being for all practical purposes indistinguishable from that of fresh blood. The increase in serum bilirubin concentration in the recipient's plasma after such a transfusion seldom exceeds 1 mg. per 100 c.cm. Even when the period of storage in this diluent reaches 21 days only about one-fifth of the erythrocytes are destroyed during the 24 hours after transfusion, and the increase in serum bilirubin concentration is unlikely to exceed 2 mg. per 100 c.cm. in our experience, although it should be noted that greater increases were found by Vaughan (1942). The transfusion of blood stored for 21 days in citrate-glucose mixtures is not followed by haemoglobinaemia, and, as Vaughan has shown, methaemalbuminaemia does not occur. Thus, following the transfusion of blood previously stored for 21 days or less with the preservative solution which is in general use, the only signs of increased blood destruction will be the small increase in serum bilirubin (only to be detected by the taking of a venous sample within a few hours after transfusion) and a slightly diminished survival of the donor cells; and the response should be scarcely less satisfactory even from the point of view of the increase in haemoglobin concentration than when fresh blood is used.

The erythrocytes of blood stored in a simple solution of sodium citrate, however, survive far less well in the recipient's circulation than the erythrocytes of blood stored for a similar period in a solution to which glucose has been added. For instance, the erythrocytes of blood stored for, say, 17 days in a simple sodium citrate solution are almost completely eliminated from the recipient's circulation within 24 hours of transfusion. A part of this destruction occurs instantaneously, owing to the rupture of the most fragile cells in the recipient's plasma, although the majority of the cells are not destroyed in this way. The haemoglobinaemia produced by this intravascular destruction is reinforced by the amount of free haemoglobin present in the plasma of the transfused blood, since blood stored with sodium citrate alone haemolyses far more rapidly than blood stored in solutions containing glucose (Rous and Turner, 1916). For instance, blood stored in sodium citrate is about one-tenth haemolysed by the end of 21 days, whereas blood stored for a similar period in citrate-glucose is only about 1% haemolysed. Thus the use of blood stored in a simple citrate solution for 17 days or more is very likely to produce haemoglobinuria and jaundice in the recipient. It may be mentioned that some of the prejudice that still exists against the use of stored blood dates from the early days of the present war, when glucose was not added to blood during storage.

Blood haemolyses more rapidly at room temperature than at 5° C., and it is almost certain that the power of the intact erythrocytes to survive in the recipient's circulation is adversely affected by prolonged exposure to room temperature. It is, however, only of academic interest to separate the effects of the injection of free haemoglobin from the effects of the more rapid destruction of the intact erythrocytes, and

it is simply necessary to realize that blood that is left standing about at room temperature before transfusion may give rise to the signs of blood destruction when it is transfused. It may be added that blood stored in the standard 540-c.cm. bottle does not reach room temperature for approximately five hours after being removed from the refrigerator (Oliver and Young, 1940).

Destruction of Recipient's Red Cells by High-titre Incompatible Agglutinins

Intravascular agglutination and haemolysis, followed by bilirubinaemia, have been produced experimentally by the injection of plasma containing high-titre incompatible agglutinins, but only when these agglutinins were present in exceptionally high titre (Aubert *et al.*, 1942). When group O blood the plasma of which contains very-high-titre agglutinins is transfused to recipients of other groups, and a haemolytic reaction follows, it is of course possible to demonstrate that the donor cells survive normally and that it is the recipient's cells which are destroyed (Aubert *et al.*, unpublished observations).

Haemolytic Reactions in Persons Affected with Haemolytic Anaemias

The transfusion of blood to patients affected with at least one variety of haemolytic anaemia may result in an exacerbation of the haemolytic process for a cause other than one of those already mentioned. As an illustration, in a case of nocturnal haemoglobinuria observed by Dacie and Firth (1943) the transfusion of agglutinin-free plasma resulted in the production of haemoglobinaemia, and the transfusion of a concentrated suspension of red cells brought about a very violent haemolytic reaction which was not due to the destruction of the donor cells, since, in fact, the latter survived normally (Dacie and Mollison, 1943).

Haemolytic Reactions Simulated by the Transfusion of Haemolysed Blood

Blood may become haemolysed by the action of contaminating organisms, by being overheated, or by being stored for excessive periods under unsuitable conditions. Certain bacteria will grow freely in stored blood at refrigerator temperature. These organisms usually cause haemolysis of the blood, so that, apart from the possible toxic effects of the bacteria or their breakdown products, the transfusion of infected blood may be followed by the signs of a haemolytic reaction. The consequences of the transfusion of infected blood may be very serious (as in Case 11). When blood is carelessly heated before transfusion it may become haemolysed—for instance, if it reaches a temperature of approximately 55° C. and is kept at that temperature for more than 10 minutes. Inquiry should always be made, therefore, to find out the method of heating (if any).

As mentioned above, blood may become grossly haemolysed if left for a long time at room temperature. The results of the transfusion of blood that had become haemolysed in this way are illustrated by Case 10 (see below).

Investigation of Haemolytic Reactions

At the outset it may be mentioned that the investigation of transfusion reactions suspected of being haemolytic in nature is greatly facilitated if certain samples are available at the time of investigation. These are: (1) a sample of the donor blood, and (2) a pre-transfusion sample of the recipient's blood. In order that these samples shall be available when they are wanted it is desirable that they should be obtained as a routine and kept for a period of 24 hours after every transfusion. It may prove impracticable to obtain a pre-transfusion sample of blood from the recipient before every transfusion, but it should be made an invariable rule that a sample of the transfusion fluid is kept. This is most easily secured by leaving the dregs in the bottle. The apparatus used may be washed immediately after transfusion, but the bottle itself should be recapped and put on one side, preferably in the refrigerator. In all cases the name of the donor or the serial number of the bottle of transfusion fluid should be recorded on the patient's notes. When there is a detachable label on the bottle this should be fastened to the notes.

When pre-transfusion samples of blood are taken from the recipient as a routine they may be conveniently obtained

through the needle used for transfusion if the latter is connected with the apparatus by means of an adaptor. Ideally, a few cubic centimetres of blood should be taken into a plain tube and allowed to clot, and a similar quantity should be taken into a tube containing oxalate. These samples should be kept in a refrigerator and not be discarded until the end of 24 hours after transfusion, when the patient has been seen again and been found to respond favourably to the transfusion. When the patient is seen within 24 hours after a transfusion and a haemolytic reaction is suspected, a large sterile venous sample of blood should be obtained and also a clean sample of urine. If the patient is seen later the same samples should be obtained, although many of the tests set out below will be inapplicable.

Precautions to be Observed in Obtaining the Venous Sample.—A large (20 c.cm.) sterile syringe with a needle of not less than 20 s.w.g. should be used. The syringe and needle should be absolutely dry or else must be rinsed in sterile isotonic saline. Venepuncture must be clean. After venepuncture the needle must be removed from the syringe before the blood is squirted into the sample tubes. The sample should be partitioned between two plain tubes (one sterile) and one tube containing dry oxalate.

The Sample of Urine.—It will be sufficient to obtain a clean sample of urine, unless there is any risk of contamination with blood (in a recently delivered woman, for instance); in such cases a catheter sample must be obtained.

Examination of the Samples

In an obscure case the majority of the following tests may have to be made:

(a) **Donor's Blood (from the Transfusion Bottle if Possible).**—(i) A sample should first be withdrawn from the bottle with a sterile pipette and kept in a sterile tube in case bacteriological examination is required. (ii) A mixed sample should next be centrifuged to determine the amount of free haemoglobin in the supernatant plasma. (iii) The blood should be regrouped, testing both cells and plasma; if necessary, the Rh group of the cells should be determined. (iv) The cells should be tested against the recipient's serum, using the pre-transfusion sample of the latter when available. (v) The titre of the agglutinins in the donor's plasma should be estimated if necessary.

(b) **The Recipient's Blood (Post-transfusion Sample).**—(i) The sample in the sterile tube should be submitted for bacteriological examination. (ii) The serum from the second clotted sample should be examined by naked eye for the presence of haemoglobin or excess bilirubin. If there is no obvious pinkness the serum should be examined spectroscopically. The amount of bilirubin or the icterus index should be estimated, and Schumm's test for methaemalbumin should be applied. The examination of the pre-transfusion sample, when available, is a valuable control. A small amount of serum should be reserved for serological tests. (iii) The oxalated sample should be divided: one-half should be centrifuged, so that the plasma can be examined for pigments as a check on the examination of the sample of serum; the other half should be kept for serological tests. (iv) **Serological Tests.**—(a) A suspension of the recipient's blood should be prepared by adding one drop of blood to approximately 2 c.cm. of isotonic sodium citrate or saline (to make a 2% suspension approximately). This suspension should be examined under the low power of the microscope for the presence of agglutinates. With experience it is possible to recognize the difference between weak diffuse clumping due to auto-agglutination and the presence of small scattered well-formed agglutinates lying in a field of free cells, the latter appearance being due to the presence of clumps of donor cells in the suspension. (b) The blood should be regrouped, testing both cells for agglutinogens and serum (the latter from the clotted sample) for agglutinins. If necessary, the Rh group of the cells should be determined and the serum examined for the presence of anti-Rh agglutinins. (c) The agglutinins in the serum should be titrated for comparison with the results of titrations carried out on samples obtained later and on the pre-transfusion sample (when available). As explained above, characteristic changes occur after the transfusion of incompatible blood, and if a sample of the donor's blood is not available, or if there is any question of the faulty labelling of a bottle so that the true donor is not known for certain, this investigation may be invaluable. (d) If necessary, differential agglutination tests should be carried out.

Note on Methods of Differential Agglutination.—Application of the technique of differential agglutination permits the recognition, in a sample of blood taken from the recipient, of foreign red cells derived from a previous transfusion. The donor cells can be recognized either by being agglutinated by a test serum which does not agglutinate the recipient's cells, or by their failure to be

agglutinated by a test serum which completely agglutinates the recipient's cells. For example, when the donor is Rh-positive and the recipient Rh-negative, by testing with an anti-Rh serum the donor's cells (if surviving) can be recognized as discrete clumps lying in a sea of free cells. Alternatively, when the donor belongs to group O and the recipient to group A the recipient's cells can be completely agglutinated with an anti-A serum and the free (donor) cells can then be counted in a haemocytometer. Many further possibilities of differentiation are available if anti-M and anti-N test sera are also used.

Application of the method of differential agglutination is practically essential for the solution of many transfusion problems, but reliable results can only be obtained with considerable experience. For a description of methods see Wiener (1942) and Dacie and Mollison (1943).

(c) **Sample of Recipient's Urine.**—The urine should be examined by naked eye and its colour noted; it should then be examined spectroscopically for oxyhaemoglobin and methaemoglobin. If the supernatant fluid contains no pigments a sample should be centrifuged and the deposit examined for the presence of pigment casts. The urine should also be examined for the presence of urobilin (Schlesinger's test). The usual tests for specific gravity and for albumin should be carried out and the reaction determined.

(The concluding part, with case reports and a list of references, will appear in next week's issue)

THREE CASES OF PROLONGED FAECAL FISTULA FOLLOWING OPERATION FOR APPENDICITIS

BY

P. T. CRYMBLE, F.R.C.S.

Professor of Surgery, Queen's University, Belfast

A slight temporary faecal fistula following the removal of a gangrenous or perforated appendix is not an uncommon occurrence, but in our experience it closes naturally in a few weeks. One may look for it in those cases in which the inflammation has spread on to the wall of the caecum, or in which closure of the appendix stump has been unsatisfactory. In the past 15 years three persistent and complete faecal fistulae following appendectomy operation have been encountered, the duration being one, three, and four and a half years. Once the fistula had become established no motions were passed by the normal route, there was marked incapacity for work, and the social result was appalling.

Case I

The first case was that of a child of 6 who was brought to Queen Street Children's Hospital 15 years ago. The child was wasted, anaemic, in a filthy condition, and had been confined to the house for a year. Its condition did not permit of attendance at school. There was a history of two operations—one for removal of the appendix, with drainage of an abscess; and a second one for the cure of an incisional hernia. It was after this second operation that the faecal fistula appeared.

On examination we found the ileo-caecal valve opening on the surface of the right iliac fossa and discharging squirts of fluid faeces. The entire caecum appeared to be absent, and it was decided that a local closure was impossible. After some building up, the abdomen was opened through the left rectus, and a diversion route established in the following way: terminal ileum divided, and both ends closed; transverse colon divided, and both ends closed; lateral anastomosis between the proximal end of the ileum and the distal end of the transverse colon.

As a result of this severe operation the child had a stormy period of five days, but recovered, and was left with a mucous fistula in place of a faecal fistula. The child returned home, put on weight, recovered strength, and was able to attend school. She attended occasionally at the out-patient department, but one was inclined to postpone the removal of the isolated bowel for some years. However, our hands were forced, as some six months after her diversion operation she walked into the out-patient department with the entire ascending colon prolapsed through the fistula and hanging down under her skirt. She was admitted to hospital, the fistula circumcised, and the terminal ileum, ileo-caecal junction, and ascending