Medical Memoranda

Pernicious Anaemia treated with Pyloro-duodenal Extract and Liver in Tablet Form

Parenteral liver therapy is occasionally complicated by severe reactions. These may take the form of local pain or general effects such as headache, flushing, urticaria, dyspnoea, and even collapse. They are usually overcome by injecting minute doses of liver extract—1/250 c.cm., 1/25 c.cm., and 1/2 c.cm.—at intervals, together with adrenaline. A nervous patient, upset by the reaction, may refuse further injections, and in such a case resort must be had to an oral preparation. An order made by the Minister of Health prohibits the oral use of liver extract as such, but in this preparation the liver is unextracted. It was tested on the following untreated case of pernicious anaemia.

CASE HISTORY

The patient, a woman of 76, was admitted to hospital during an air raid in a state of nervous prostration, her house having been bombed. On recovery she complained of left-sided headaches, which, she stated, had been present for the past six months, occurring in attacks lasting several days at a time. During the last few weeks she had also been subject to fainting attacks. On examination she was found to be a thin pale woman. Her memory was much affected and she had no recollection of her recent misfortune. The right pupil was fixed and irregular, reacting neither to light nor to accommodation. The media of both eyes were opaque. The conjunctivae and mucous membranes of the mouth were pale. Blood pressure was 120/80. The spleen was not palpable, but the splenic dullness was increased. The knee- and ankle-jerks were absent and the calves were more tender than normal. The other systems showed no abnormal physical signs. Blood count: red cells, 1,720,000; Hb, 47%; C.I., 1.38; leucocytes, 2,000—polymorphs 45%, eosinophils 1%, basophils 2%, lymphocytes 48%, monocytes 4%. The red cells were large and showed marked anisocytosis and some poikilocytosis. Reticulocytes amounted to 0.1%. A histamine test meal revealed complete achlorhydria. The patient was given 4 tablets of binaemon three times a day. This was equivalent to a daily dose of 6 g. of fresh liver and 3 g. of fresh pylorus.

Progress.—The effect on the blood picture is shown in the accompanying chart. The reticulocyte crisis did not start till the

eighth day, and was small, reaching only 5% maximum, but the rise in the red cell count and in haemoglobin was fairly rapid. By the end of the fourth week the red cell count had risen by 2,000,000, and by the seventh week had reached 4,000,000. This is similar to the recovery rate met with when raw liver is used but slower than that produced by parenteral liver extract, which raises the count to 4,000,000 in about a month. Normal figures were finally reached in the eighth week.

CONCLUSIONS

This preparation of liver and pyloro-duodenal extract in tablet form was found to be effective in treating pernicious anaemia. The difference in cost between this and parenteral treatment is not great. With the latter, assuming that initial doses of extract corresponding to 300 g. of liver are spread over the first 4 days and followed by weekly doses equivalent to 100 g. till the blood count is normal, as recommended by Murphy (1939), the cost of the treatment would be approximately 17s. at hospital prices, as compared with 23s. for the oral method.

My thanks are due to Dr. Allen Daley, Chief Medical Officer to the London County Council, for permission to publish this paper, and to Organon Laboratories Ltd. for supplies of binaemon tablets.

J. STEWART LAWRENCE, M.D., M.R.C.P.,

Physician, E.M.S.

REFERENCE

Murphy, W. P. (1939). *Anaemia in Practice: Pernicious Anaemia*, Philadelphia.

Glandular Fever

The article by Drs. J. Shafar and J. C. R. Weir (*Journal*, Feb. 20, p. 218) on a case of glandular fever with a meningeal reaction raises a number of points of interest at the present moment, when there appears to be an increase in the incidence of infectious mononucleosis. Those authors refer to the general characteristics of their series of some 200 cases. In particular they stress the benign nature of the condition, its marked infectivity, and the unreliability of the Paul-Bunnell reaction. They also state that in differential white cell counts of the blood the polymorphonuclear leucocytes totalled some 50%, and that the remainder was made up of small and large lymphocytes with an average of 10% of monocytes.

During the past few months we have admitted to the County I.D. Hospital here two patients whose cases are fully reported below. These cases closely resemble each other, and were regarded as examples of a clinical entity conforming to the classical description of glandular fever. It will be seen, however, that they differ widely from the cases of Drs. Shafar and Weir in certain important respects. These are: (a) The course of the illness was by no means benign. Both patients were sharply ill with intermittent pyrexia, of 14 days' duration, which reached 103.8° F. in one instance and 102.0° F. in the other. (b) Infectivity was apparently not pronounced. Both were single cases which occurred separately under Service conditions, and no other cases were reported. Haematological investigation of attendants was not carried out, but none showed any clinical signs of infection. (c) The Paul-Bunnell test in both cases was positive in high dilutions. (d) The differential white cell counts of the blood showed much higher degrees of mononucleosis, the combined percentages of lymphocytes and monocytes reaching as high as 82. (e) There was marked faucial involvement.

It is felt that these considerations help to strengthen the opinion expressed by Drs. Shafar and Weir that there may be "separate and distinct varieties of glandular fever."

CLINICAL HISTORY OF TWO RECENT CASES

Case 1.—A man aged 21 was admitted on July 29, 1942, with a history of headache, shivering, malaise, and abdominal pain of 4 days' duration. On examination he was found to have a temperature of 102.6° F. and a pulse rate of 106. There were no other positive clinical findings, but, in view of his severe headache, lumbar puncture was performed. Clear fluid was obtained, not under pressure, and laboratory investigation showed no departure from normal. A blood count taken on admission gave a total of 5,000 white cells per c.mm.—polymorphonuclear leucocytes 54%, lymphocytes 41%, and monocytes 5%. Blood culture proved negative, and a Widal reaction revealed only slight H-agglutination, which was disregarded as resulting from inoculation. Pyrexia was intermittent, and on Aug. 3 the patient complained of sore throat. He was then found to have a greyish membrane on both tonsils. This separated easily without bleeding, and a throat swab was negative for *Corynebacterium diphtheriae*. Next day non-tender enlarged glands were observed in neck, axillae, and groins, and there was a morbilliform rash on the abdomen. By this time the leucocytes consisted of: polymorphs 32.5%, and lymphocytes and monocytes together 67.5%, with a total white cell count of 11,300 per c.mm. The Paul-Bunnell test was positive with a titre of 1:896. By Aug. 5 the patient was acutely ill, with a high temperature, rapid pulse, and an extension of the tonsillar membrane and the rash. On this day the white cells numbered 26,000 per c.mm.—polymorphonuclear leucocytes 29%, lymphocytes 22%, and monocytes 49%. He remained very ill for three days and then rapidly improved, with recession of all symptoms. He was discharged well on Sept. 15.

Case 2.—A man aged 22 developed severe headache on Oct. 10, 1942, with nausea and anorexia, and noticed a transient rash, apparently of erythematous type, on both forearms. He reported sick on Oct. 19 and was admitted to sick quarters the same day. He was found to have an irregular pyrexia, reaching on occasion 102° F., and by Oct. 24 he had painless glandular swellings in neck, axillae, and groins. The patient was now admitted to hospital here, and was found to have, in addition, marked faucial congestion, with patches of loose exudate on both tonsils. His blood picture on admission was: polymorphonuclear leucocytes 18%, lymphocytes 70%, and monocytes 12%, with a total white cell count of 16,200 per c.mm. The Paul-Bunnell test was positive with a titre of 1:640. Next day he complained of severe headache, and lumbar puncture was performed. The cerebrospinal fluid was normal in all respects. On Oct. 29 the tonsillar membrane had extended, and the white cells numbered 15,200—polymorphs 19%, lymphocytes 75%, and monocytes 6%. The pyrexia subsided on Nov. 1, and the patient subsequently improved rapidly. He was discharged well on Nov. 21.

I am indebted to Dr. J. L. Horne, M.O.H., for permission to report these cases, and to Dr. C. C. McCallum, R.M.O., for help in the preparation of the clinical notes.

R. J. LUMSDEN, M.B., Ch.B., D.P.H.

County Infectious Diseases Hospital,
Strathpeffer, Ross-shire.