BRITISH MEDICAL JOURNAL

LONDON SATURDAY JUNE 5 1943

THE OPHTHALMOSCOPE IN CLINICAL MEDICINE THE INTERIOR OF THE LIVING EYE CONSIDERED AS A BIOCHEMICAL LABORATORY*

BY

A. MAITLAND RAMSAY, M.D., LL.D., F.R.S.Ed.

Until the ophthalmoscope was discovered the interior of the living eye was *terra incognita* to the physician. The ocular structures deeper than those visible in the anterior chamber were for all clinical purposes a sealed book, and our knowledge of the inner eye and its diseases had to be acquired from the observation and the interpretation of symptoms. That method of procedure is beset with many pitfalls, for it rests almost entirely upon the patient's own account of the disease, and very few of them can give a clear and connected description of the character, the site, and the duration of the symptoms. It is only when the doctor is able to analyse all the patient has told him that he can, by a few direct questions, separate the wheat from the chaff. When he does succeed in so doing valuable information is always obtainable; but the clinical value of such an examination depends almost entirely upon the surgeon's special skill and his personal experience. Nevertheless much useful information has been acquired and accumulated by the careful study of the symptomatology of disease, especially when a knowledge of medicine went hand in hand with literary culture. A striking example of what can be achieved in that way is seen in William Mackenzie's *Practical Treatise on Diseases of the Eye*, in which accurate medical observation is accompanied by such vivid clinical description that the reader almost sees the patient rising from the printed page.

The Clinician and the Laboratory

Clinical observations differ from laboratory investigations in medical practice, inasmuch as the latter are quite impersonal, whereas the former always regard the patient as an individual entity. That aspect of the position has tended to shrink into the background in recent years owing to the steadily increasing importance that is now attached to intensive laboratory research. In no circumstances, however, can the clinician be replaced by the laboratory worker. In a large number of cases both of them are required if the patient is to have the benefit of the highest medical skill; but when consultation is necessary the clinician must never allow himself to be supplanted by the laboratory worker. In the initial diagnosis of disease the clinician must always take first place. He works beside the patient, and ought to exhaust every means of examination at his disposal to reach a diagnosis by himself, and when he fails he ought always to know the kind of assistance he is requiring before he seeks help from his colleague in the laboratory. The best results are most likely to be got when both workers meet and discuss together the problem they are anxious to solve.

When clinical symptomatology is uncontrolled by objective examination, or by any form of experiment, patients who are not in the same category are apt to be grouped together. For

example, before the introduction of the ophthalmoscope patients suffering from failing sight or loss of sight were all included under the general heading "amblyopia and amaurosis." Many attempts were made to reclassify those said to be suffering from amblyopia; but as vision becomes more impaired accurate diagnosis is wellnigh impossible, and when the stage of blindness or amaurosis is reached some with grim humour defined that condition as a state of matters in which the patient sees nothing and the surgeon sees nothing.

The laboratory worker always aims at scientific precision, and concerns himself for the most part with analysis and measurement. He puts his chief reliance upon chemical and microscopical tests, and upon experiments whose results can be tabulated in figures. The clinician requires something more. His education at the bedside has qualified him for dealing with human suffering; consequently he is able to make a character study of the individual patient, and knows how to help him with words of hope and encouragement. Our forefathers paid great attention to the study of human beings both in health and in disease, and they encouraged medical students to cultivate their power of observation and to train their sense of touch; but this personal training of eyes and hands is passing into the background. In these days we must be careful not to neglect the clinical tradition of the personal art and skill of mind and hand, without which a family doctor will find the most up-to-date scientific training in the laboratory insufficient in many of the situations he is called upon to cope with in the daily round of general practice. Modern medicine has become so highly mechanized that a medical practitioner is being tempted to rely more and more on the findings of an instrument and on a report from a laboratory. We are forgetful that the human brain, when it is highly trained, is the finest of all instruments; and when we employ some mechanical apparatus to do its work, and save ourselves the trouble of thinking, we must be careful that we do not sidetrack our own intelligence.

The Laboratory of the Eye

Among the instruments on which a medical student is taught to rely in his clinical work, the ophthalmoscope is in a category by itself. The ophthalmoscope enables us to see the nervous and vascular structures of the fundus oculi and to study physiology and pathology during life. For our immediate purpose we may liken the interior of the living eye to a fully equipped biochemical laboratory, in which Nature herself performs all the experiments. With the help of the ophthalmoscope we can look into this laboratory through the window of the pupil and watch Nature at work; and as Nature's ways throughout the body are everywhere consistent, we are justified in concluding that what we can see in the eye is an index of what is taking place hidden from view in other regions of the body. The fundus oculi therefore affords a unique opportunity for the clinical study of vital processes, and is a convenient place for the clinician to meet the pathologist. In the laboratory of the eye the patient can never be forgotten. The pathologist working in his own department examines and

* An address, illustrated by lantern slides, given to the Bute Medical Society, University of St. Andrews.

furnishes a report on the specimens that have been sent to him, but he does not need to have seen the patient or to be interested in the human aspects of the case; the clinician, however, always has to deal with a patient behind the disease.

Notwithstanding his skill and the perfection of his laboratory methods a chemist's experiments are not invariably successful, and in the laboratory of the human body Nature's experiments are also followed by success and failure. When Nature's experiments are successful we are quite unconscious of the functioning of our bodily organs. Everything seems to go with the precision of clockwork. We have a supreme sense of well-being. We enjoy good health, and feel quite fit for our daily duties. When, on the other hand, Nature's experiments are unsuccessful we at once become conscious of abnormal sensations, which we recognize as symptoms of disease. In some instances an ophthalmoscopic examination enables us to detect disturbance of metabolism in its earliest stages. The physician who neglects the routine use of the ophthalmoscope in general practice misses a great deal. The original models of the instrument were difficult to manipulate, but the modern electric ophthalmoscope enables us to examine the eye as easily as we can take the temperature, auscultate the chest, or apply chemical tests to the urine. Before the medical practitioner can understand what he is looking at, however, he must be able to correlate the ophthalmoscopic appearances with the microscopic anatomy of the fundus oculi. Each of the three tunics of the eyeball contributes to produce the ophthalmoscopic picture of the eye ground. The sclerotic may be regarded as the white canvas on which the picture is painted. The blood in the capillaries of the choroid and retina is responsible for the red colour, while the transparency of the retina gives it a peculiar lustre and characteristic translucency.

The Fundus Oculi

In every organ of the body there is a wonderful adaptation of structure to function. In the eye perfect transparency is the primary requisite, and that has been secured for all the structures especially concerned with vision by an almost complete impermeability of the intra-ocular capillaries to the passage of colloidal substances. In a state of health water, gases, and crystalloids readily escape from the retinal capillaries, but colloids for the most part are retained in the blood. As a result of this retention of colloids a peculiar translucency is the outstanding characteristic of the normal fundus oculi, and at the onset of disease that translucency is the first thing to be lost. The capillaries are very sensitive to the quality of the blood. The least taint of toxæmia causes them to dilate rapidly, and, by damaging the endothelium of their walls, allows colloids to escape. Congestion of its capillaries, and its infiltration with colloids, quickly impair the normal transparency of the retina, which can be recognized clinically by lack of translucency in the ophthalmoscopic appearance of the fundus oculi. Instead of its normal characteristic brilliance the fundus looks like glass which has lost its lustre after having been breathed upon. At the outset the departure from the normal is very slight; but clinically it is one of those variants situated on the borderline separating the physiological from the pathological.

The ophthalmic surgeon occasionally sees those cases among patients suffering from eye strain sent to him by colleagues. All of them suffer from headache due to an error of refraction, and the great majority are at once relieved by the use of suitable spectacles; but a few get no benefit from optical treatment alone. These patients form a group by themselves, and all of them present the pathognomonic sign of lack of translucency in the ophthalmoscopic picture of the fundus oculi. There is no deterioration in the patients' usual acuity of vision, but some complain of attacks of misty sight accompanied by black spots floating in front of the eyes or by sparks and flickering lights dancing in the field of vision. These subjective symptoms are supposed to be due to increase of blood pressure in the central artery of the retina, and although they may be premonitory they are quite distinct from those signs accompanying an ordinary sick headache or an attack of migraine. An important concomitant sign is that although the quantity of urine passed in 24 hours is somewhat less than usual, the specific gravity is not increased in the same ratio, and when

a specimen is kept overnight in a urine-glass there is almost invariably a deposit containing numerous minute crystals of calcium oxalate; but there is no interference with the excretion of urea. It is probable that, in the case of those patients, waste products which ought to have been excreted in the urine have been retained in the blood, and have caused a mild form of toxæmia by dilating the capillaries both of the retina and of the glomeruli of the kidney. That hypothesis gave the clue to the successful treatment of those patients. More than 50 years ago a similar condition in general medicine was described by Sir Andrew Clark in his papers on renal inadequacy, and by Charles Murchison in his book on functional derangement of the liver. These authors have demonstrated, from a clinical standpoint, that in both the kidney and the liver the disorder is functional, and that healthy action can be restored by the use of eliminant remedies. When, in addition to the use of the glasses required to correct the error of refraction from which the patients we are considering suffered, a course of eliminant treatment is prescribed relief from the symptoms is speedy and permanent. The normal ratio between urine volume and the specific gravity is restored, the oxaluria disappears, the retina recovers its normal transparency, and the headache is cured.

The Fundus and the General Circulation

The circulation can be studied in the fundus oculi as an integrated whole, and as the circulation of the blood in the eye is governed by the same rules as those which regulate its distribution to every organ of the body it gives a reliable clue to the state of the general circulation. The intra-ocular circulation is peculiar inasmuch as it has to be carried on under the pressure of the intra-ocular fluids, and the smooth steady flow of blood throughout the retina depends upon an accurate balance being maintained between blood pressure on the one hand and the pressure of the intra-ocular fluids on the other. By the usual method of ophthalmoscopic examination pulsation is seen in the veins as they pass over the optic disk, but no pulsation is visible in the arteries unless there has been a breakdown in the physiological adjustment of those two pressures. For example, if there is a sudden fall in the diastolic systemic pressure, as occurs in aortic regurgitation, or an increase in intra-ocular pressure as in glaucoma, pulsation in the main branches of the retinal artery can always be seen with the ophthalmoscope. Venous pulsation therefore is physiological, but arterial pulsation is always pathological. The walls of the vessels in a state of health are so transparent that arteries are distinguished from veins by the colour of the blood. When used by the direct method the ophthalmoscope acts as a low-power microscope, and gives a magnification of 15 diameters. We are thus able to see arterioles during life which we cannot see with the unaided eye in a post-mortem specimen.

Ocular Manifestations of Disorders

In what follows I shall limit myself to a consideration of the ocular manifestations of the injurious effects of disordered metabolism upon the cardiovascular-renal system. In the act of living there is a constant manufacture of toxic products, which, if health is to remain unimpaired, must either be destroyed at once or removed from the body with the least possible delay. Our bodily life therefore seems to be very much a matter of chemistry. It is only at the level of the capillaries that exchanges in both directions take place, for the purposes of metabolism, between the blood and the tissues. By that means the physiological processes of assimilation and elimination are conducted and the nutrition of the body is adequately maintained. The capillaries are very sensitive to the presence of impurities in the blood, and high blood pressure is the usual accompaniment of toxæmia due to the absorption of metabolic waste products; but so long as the walls of the blood vessels remain healthy there is little danger of hypertension causing rupture and haemorrhage. That is a fact which should not be forgotten, because so many are said to be suffering from high blood pressure nowadays that the subject is a common topic of conversation far beyond the bounds of the medical profession. As a rule the general trend of all such conversations is unnecessarily gloomy. It is well known that in some families there is a hereditary predisposition to a state of high blood pressure in late middle life; but it is not always

remembered that although increase of blood pressure is frequently an accompaniment of advancing years it is not in itself a disease. Very considerable variations are quite compatible with ordinary life and health, and probably also with normal expectation of life. Ellis is of the opinion that in more than 90% of the patients the hypertension is benign. Allbutt describes this condition as hyperpiesia, which often exists unaccompanied by symptoms, and is sometimes discovered in the course of a routine medical examination for life insurance. Those who suffer from a benign form of increased blood pressure ought to be discouraged, therefore, from taking too much interest in their high blood pressure.

In practical medicine undue attention ought never to be directed to any single sign or symptom. It is the association of symptoms that is of value in prognosis, and as a rule the ophthalmoscope and the sphygmomanometer ought to be used together. The clinical value of the actual number registered by the latter instrument can only be properly assessed after careful digital examination of the pulse, testing the urine, and thorough interrogation of the cardiovascular system as a whole. Unless there is some sign of cardiac weakness or of cerebral disturbance the restrictions placed upon the patient's general mode of life have as a rule little, if any, influence on the hypertension; but by interfering with the enjoyment of life and curtailing usefulness they create an anxiety neurosis which is far more harmful than the increased blood pressure. It is even worse still when on the supposition that the patient is suffering from dangerous disease he is sent to a spa, where strenuous measures are taken to lower blood pressure. Such treatment may be followed by disastrous results. A sudden fall in systolic pressure is apt to be accompanied by a relative increase in the diastolic, with engorgement of blood in the capillaries. One of the first signs of the upset in the physiological ratio of the systolic-diastolic pressures is the occurrence of haemorrhage from the capillaries of the retina. When that happens the prognosis is very serious, because it is often the prelude to a complete breakdown in general health. A disturbance in the physiological ratio between the systolic and the diastolic pressures is of much worse prognosis than a far higher hypertension in which the normal ratio is maintained and both pressures run parallel. When the hypertension is malignant the prognosis is grave. In these circumstances a form of Bright's disease is almost certainly present, the downward course of which is very rapid. The condition is invariably fatal, and death with uraemic symptoms usually occurs within a year. According to Ellis, in his Croonian lectures for 1941, the pathognomonic sign of malignant hypertension is the occurrence of papilloedema in both eyes, and in some cases the ophthalmoscope also reveals retinitis and haemorrhages.

When toxæmia and high blood pressure are of long duration the ophthalmoscopic picture of the fundus oculi becomes so lustreless that it resembles red velvet, and when the toxin is of special virulence loss of transparency and oedema of the retina become so very pronounced that capillary haemorrhages are usually seen scattered irregularly all over the eye ground. For example, the toxin due to pregnancy excites a very virulent reaction, and attacks eyes and kidneys simultaneously. Acute inflammation of both the retina and the kidneys is present from the outset, and the pathological changes seen in the eye with the ophthalmoscope are an index of what is taking place in the kidney. The inflammation of the retina appears suddenly, runs a violent course, and is characterized by such intense oedema that detachment may occur. Multiple capillary haemorrhages are always present. Both eyes are affected, and sight is impaired, and it may be lost very early in the course of the disease. In patients who have had no ante-natal supervision loss of sight may be the first warning of the approach of danger. The symptoms are often so overwhelming that the case may well seem hopeless; but the experienced physician knows that complete recovery is to be expected after the child is born. No better illustration could be given of the clinical aphorism that when a disease is due to toxæmia a favourable prognosis can always be given if the toxic agent can be wholly eliminated before organic disease of the blood vessels takes place. Unless that occurs, and the toxic agent is eliminated, recovery will be far from complete, and hypertension and albuminuria will persist. In these circumstances blood pressure

is likely to remain high, and in course of time arteriosclerotic changes will appear in the retinal arteries and renal function will steadily fail.

Early Signs of Pathological High Blood Pressure

One of the earliest signs of pathological high blood pressure is constriction in the calibre of some of the branches of the retinal artery, and the degree of constriction is a reliable guide to the height of the systemic arterial hypertension. The constricted vessel usually takes a straighter course than is the case with a normal branch of the retinal artery, and when spasmodic constriction is accompanied by straightening of the course of the vessel it is as a rule evidence of organic disease. Loss of transparency of the wall of the retinal artery is another of the early results of persistent pathological hyperpiesia; consequently when spasm occurs in a branch of the retinal artery the outline of the vessel wall is clearly visible, and when spasm is complete the function of the retina supplied by the affected vessel is lost, and will not be restored until the spasm is relieved and the circulation of the blood is re-established. An analogous state of matters can occur in the circulation of the lower limbs—intermittent claudication: severe cramping pain in the muscles of the calf follows exertion, and compels the patient to rest until the arterial spasm passes off. Whenever that takes place the pain is relieved and the patient is able to walk. In both the eye and the leg an underlying constitutional state has existed long enough to produce vascular changes sufficiently serious to cause temporary obstruction of the circulation. When these vascular symptoms are persistent they are usually premonitory of the onset of arteriosclerosis, which is always pathological. Old age of itself will not cause it.

Arteriosclerosis

The earliest signs of arteriosclerosis are often seen in the fundus oculi; therefore in many instances we are able to study them in considerable detail during the patient's lifetime. When sclerosis occurs in the retinal vessels it is a fairly sure sign that it is also present in other organs of the body. Arteriosclerosis, however, is a most capricious disease. One organ may be affected dangerously while others escape.

Arteriosclerosis of the retinal blood vessels is similar to the arterio-capillary fibrosis of Gull and Sutton, and the ophthalmoscopic appearances have been admirably described by Marcus Gunn. The arteries are irregularly contracted, abnormally tortuous, and even the smallest exhibit a brightness of the central light streak which is very characteristic. The artery loses its transparency and becomes hard and rigid; consequently when it crosses a vein it not only conceals the latter from view, but also mechanically exerts so much pressure upon its walls that the flow of blood is hindered, and any sudden rise in blood pressure is apt to precipitate rupture and haemorrhage. In these circumstances the use of the ophthalmoscope is our best guide to the state of the cerebral blood vessels. There is a close relationship clinically as well as anatomically between the retinal and the cerebral blood vessels. Both of them arise from the internal carotid artery. When, therefore, arteriosclerosis develops both sets of vessels are likely to be affected simultaneously, but not necessarily to the same degree. Serious cerebral haemorrhage is more frequent than severe retinal haemorrhage. The vessels in the brain which rupture most frequently are of larger size than any in the retina. In the great majority of cases the true essence of retinal haemorrhage is capillary dysfunction rather than increased blood pressure acting upon a sclerosed artery. When, however, either retinal haemorrhage or cerebral haemorrhage occurs during sleep the patient has probably been dreaming, and the excitement of the dream has caused a rise in the general blood pressure. In these circumstances the haemorrhage is probably due to the rupture of a sclerosed artery either in the eye or in the brain, or in both simultaneously.

The Warning of Retinal Haemorrhage

Haemorrhage from the blood vessels of the retina is always evidence of disease and ought to be taken as a warning. Its causation should be investigated as thoroughly as would be the case after haemoptysis or haematemesis, but it is a mistake to take too gloomy a view of its significance. As an isolated sign it is not likely to increase the gravity of the prognosis

regarding the patient's life, but if the bleeding is extensive the delicate structure of the retina may be injured so seriously that sight is gravely endangered.

Danger to sight depends upon the size and the site of the haemorrhages. In the great majority of medical cases the capillaries are probably the starting-point of retinal haemorrhage; but, when arteriosclerosis is advanced, rupture of an arteriole will determine a copious and destructive bleeding. When blood escapes into the inner layers—i.e., the conducting fibres—of the retina the clots are broken up by the nerve fibres. As a rule they are of bright red colour, small and flame-shaped. They generally disappear by absorption in about three weeks' time, but, unless the underlying toxæmia is eliminated, bleeding of this nature, once it begins, is prone to continue, and fresh haemorrhages take the place of those which have become absorbed. In this situation haemorrhages give rise to so few subjective symptoms that the patient may be unaware that anything is seriously wrong until his eye has been examined with the ophthalmoscope. When, on the other hand, the outer bacillary layer—i.e., the functioning unit—is implicated sight is seriously threatened from the outset by damage to the rods and cones, and the probability is that vision will be permanently impaired, if it be not altogether lost. These haemorrhages are deep-seated, and do not disappear quickly. They are dark red, and more or less circular. When the macula itself is involved central vision is obliterated, and the patient is unable to read or to recognize a friend on the street. Recovery of sight is not probable if the haemorrhage is deep-seated, and in any case vision is sure to be distorted—metamorphopsia—because of disturbance in the regular mosaic of the cones in the macula.

Types of Retinitis

High blood pressure, arteriosclerosis, and haemorrhage are the pathological forerunners of retinitis: but it is the quality of the blood that determines the distinctive aetiology in an individual case. Pathologists have described many varieties of inflammation of the retina; but the three most important clinical types—viz., the arteriosclerotic, the albuminuric, and the diabetic—are easily differentiated by an examination of the urine.

1. *Arteriosclerotic Retinitis*.—The ophthalmoscopy of this disease has been carefully described by Foster Moore, who was the first to suggest that it is a clinical entity distinct from albuminuric retinitis. In this form of disease persistent high blood pressure is always accompanied by extensive sclerosis of the retinal arteries. Its chief clinical characteristic is that the urine contains no albumin and there is complete absence of oedema. The patients are usually in late middle life, and are as a rule of stout build, of a fresh ruddy complexion, and of active habits. They enjoy speaking about their good health and often boast that they never require a doctor, while all the time their constitution is being steadily, though silently, undermined by the furtive advance of arterio-capillary fibrosis. Unless he has used the ophthalmoscope the family doctor may easily be put off his guard by the absence of albumin in the urine, and he is greatly taken aback when his patient is struck down suddenly by apoplexy or by a cardiac seizure. Mahomed insisted that high blood pressure rather than albuminuria is the pathognomonic sign of certain forms of Bright's disease, and some regard arteriosclerotic retinitis as a sign of chronic nephritis in which only the merest trace of albumin is found occasionally in the urine.

The ophthalmoscopic appearances vary according to the stage reached by the disease. Its onset is characterized by a reduction in the calibre of the arteries and by a loss of transparency of their wall. As the disease advances the veins are crushed where they are crossed by an artery, and many of the arterioles, especially those in the region of the macula, become twisted like a corkscrew. In the early stages the optic disk is pink from capillary congestion, but as the disease progresses it becomes pale and atrophic from sclerosis of its vascular supply. Small punctate haemorrhages, usually increasing in size and number, are seen scattered irregularly over the eye ground, and in addition there are numerous small circular hard-looking white spots situated for the most part in the central area of the fundus. The walls of the larger arteries become more and more opaque until at length the column of blood is concealed and the vessels resemble silver or copper wire. As time passes the lumen of the artery becomes almost obliterated and is now little more than a fibrous cord, and the area of the retina dependent upon it for its blood supply is starved, with the result that in the later stages of the disease patches of atrophy appear in the eye ground, and vision becomes permanently impaired.

2. *Albuminuric retinitis* can only be diagnosed after toxæmia, high blood pressure, and the presence of albumin in the urine have

all been established. Both eyes are usually affected, but as a rule not to the same extent. The pathogenesis of the retinal changes depends upon high blood pressure and spasmodic constriction of the arteries; therefore the retinitis ought not to be regarded as a separate entity, but as an important and visible part of the general degenerative processes which are taking place in the kidneys. It is usually said that the patient is not likely to live more than two years after retinitis develops; probably that is true when applied to hospital patients as a whole, but there are many exceptions in private practice. The patient's general health does not become worse because his eyes have become affected, but the unfavourable prognosis after the discovery of the ocular signs is due to the fact that by the time the retinitis is detected the disease of the kidney is advanced and progressive. The earlier the retinitis appears, therefore, the worse the general prognosis, more especially if the disturbance in the eye coincides with the onset of uraemic symptoms.

Ophthalmoscopically the pathognomonic sign is oedema of the retina, and that is a sure indication that the reserve power of the kidney is being encroached upon. In the more acute cases white "cotton-wool" patches of soft-looking exudate are scattered over the fundus. The veins are distended and tortuous, and there are always signs of arteriosclerosis, accompanied by haemorrhages varying in both size and shape. They are scattered irregularly over the fundus, and are more numerous in the region of the macula. Lastly, one of the most characteristic features of renal retinitis is a star-shaped arrangement of minute white glistening spots round about the macula. This stellate arrangement of white dots, although very characteristic of albuminuric retinitis, is not pathognomonic of the disease, because it may occur in any form of acute neuroretinitis in which there is much oedema of the retina round about the macula.

3. *Diabetic retinitis* occurs for the most part in the elderly—more especially in those who suffer from a comparatively mild form of glycosuria complicated by high blood pressure, by an appreciable degree of arteriosclerosis, and by the more or less persistent presence of a trace of albumin in the urine. It is almost never seen in the young, who are the chief sufferers from the acute forms of diabetes. The ophthalmoscopic picture differs from that of albuminuric retinitis by the absence of oedema and the "cotton-wool" patches of exudation. There are, however, in a typical case round yellowish-white plaques, situated for the most part between the papilla and the macula. As the disease advances these patches tend to coalesce and ultimately to form a ring surrounding the whole macular region. Haemorrhage is the outstanding feature of diabetic retinitis; consequently the ophthalmic surgeon is more concerned about the sight than about the life of the patient. The degree of vascular degeneration is the most important factor in the prognosis. The vascular changes are steadily progressive, and for the most part they fall upon the veins. Even the characteristic deep-seated punctate haemorrhages which are so suggestive of diabetes are apparently derived from venous radicles, and in some desperate cases massive haemorrhages from the veins flood the vitreous humour. These copious dense haemorrhages frequently occur during sleep, and the patient awakens to find himself blind in the eye affected. In some cases both eyes are implicated.

Thrombosis of the central blood vessels of the retina is the ocular complication most to be dreaded when glycosuria is accompanied by albuminuria and when high blood pressure is persistent. Vision is lost suddenly, but several premonitory seizures usually occur before blindness becomes permanent. This condition ought always to be regarded as a serious warning of cardiovascular failure. The eye itself is also seriously compromised, because thrombosis is always accompanied by profuse retinal haemorrhage, which is often followed by the onset of secondary glaucoma. When that stage is reached the patient suffers from incurable blindness and more or less constant pain; consequently enucleation of the eyeball is the only treatment that can bring relief.

Commentary

To sum up very briefly, my endeavour has been to reach the following conclusions:

1. The interior of the living eye is a laboratory in which vital processes, both normal and abnormal, can be studied during the patient's lifetime.

2. Vascular changes such as capillary congestion, arterial degeneration, haemorrhage, and exudation can be seen with the ophthalmoscope; but the value of such observations in the prognosis of general disease cannot be properly estimated until they are correlated with such symptoms of general disease as high blood pressure, albuminuria, glycosuria, uraemia, and anaemia.

3. The debt which general medicine owes to ophthalmology cannot be fully appreciated until the physician examines the eye with the ophthalmoscope as habitually as he feels the pulse, interrogates the heart and lungs, and applies chemical and microscopical tests to the urine.