The visceral pleura of the right lung was similarly infiltrated to a maximum thickness of 1.2 cm., and the right lung was completely collapsed and fixed against the vertebrae. Four firm, broad, fleshy bands of the growth extended between the two layers of the pleura; they appeared like adhesions replaced with growth. On the cut surface of the right lung small areas of collapsed lung tissue were visible (Fig. 2), but the greater part consisted of white, somewhat

FIG. 2.—The neoplastic mass opened to show the collapsed lung containing nodules of growth, the parietal pleura greatly thickened and infiltrated, and bands of neoplastic tissue connecting the two. A, Tongue. B, Collapsed lung. C, Neoplastic bands. D, Nodules of growth. E, Parietal pleura.

soft growth. The bronchi contained muco-pus. Many bronchi were involved in the neoplasm, but this could not be seen actually arising from a bronchial wall. The hilar glands were much enlarged with white growth. The trachea and bronchi of the left lung contained muco-pus; the left hilar glands were not enlarged. Both lobes of the left lung were partly collapsed. Two small, flat, thin, round plaques were seen on the visceral pleura of the left upper lobe.

There was considerable venous engorgement in the chest and neck veins. The lumen of the oesophagus was irregularly distorted by discrete mediastinal glands enlarged by secondary neoplastic deposits. The liver showed a pale mottling; its relation to the growth has already been described. Metastases were detected in the right bronchial, all the mediastinal, the right supraclavicular, one large right axillary, and the coeliac axis lymph glands. The right adrenal gland was displaced and partly surrounded by an extension of the neoplasm beneath the diaphragm, but the capsule of the gland appeared intact, a fact confirmed microscopically. The left adrenal was normal.

FIG. 3.—Photomicrograph showing masses of polyhedral cells and a few multinucleated giant cells.

Microscopical Examination.—Sections of the right lung and pleura and lymph glands showed a poorly differentiated carcinoma (Fig. 3) composed of masses of polyhedral cells varying in size and including

multinucleated giant-cell forms in some areas. In many places where the tumour cells outlined the trabeculae of connective tissue they tended to be columnar-shaped. A large number of the cells were somewhat vacuolated; sections stained with mucicarmine gave an indefinite result. A few small areas of necrosis were present. Mitoses were not numerous.

Conclusion

The growth was an anaplastic carcinoma of the bronchus with extensive pleural involvement and metastases in the bronchial, mediastinal, right supraclavicular and axillary, and coeliac axis lymph glands.

We wish to express our thanks to Dr. W. J. Pearson for permission to publish the case and to Prof. R. S. Pilcher for performing the thoracoscopy and biopsy. We are indebted to Dr. D. M. Vaux of the E.M.S. Pathological Service for conducting the necropsy and reporting upon the fluid, the biopsy, and post-mortem material.

REFERENCES

- Beardsley, J. M. (1933). *Canad. med. Ass. J.*, **29**, 257.
Hirsch, E. F., and Ryerson, E. W. (1928). *Arch. Surg.*, Chicago, **16**, 13.
McAldowie, A. M. (1876). *Lancet*, **2**, 570.
Menne, F. R., and Anderson, M. W. (1941). *J. Amer. med. Ass.*, **117**, 2215.
Simons, E. J. (1937). *Primary Carcinoma of the Lung*, Chicago.

NASAL INTUBATION: DANGERS AND DIFFICULTIES FROM THE RHINOLOGICAL ASPECT

BY

A. R. DINGLEY, F.R.C.S.

Surgeon, Royal National Ear, Nose, and Throat Hospital

Pernasal intubation is to-day so often employed as a preliminary step in anaesthesia that it is perhaps not out of place to emphasize, from the rhinological standpoint, how necessary it is to have an adequate knowledge of the existing intranasal conditions in every case, in addition to a realization of the implications of the widely varying anatomy of the nose and of the factors limiting the safety of nasal intubation which nasal pathology can introduce. The fact that damage to the delicate structures of the nasal defensive mechanism occurs so relatively infrequently reflects great credit on the skill and care of the anaesthetist of to-day; but that harm can, and does, result cannot be denied, and serious consequences have followed the passage of a tube, either from faulty technique or from ignorance of the presence of intranasal conditions which should have debarred intubation. In other cases, less serious and often overlooked sequelae have manifested themselves later.

The nasal passages have highly important and essential physiological functions to perform: by warming, moistening, and filtering the inspired air they provide a highly specialized defensive mechanism for the protection of the respiratory tract; they ventilate the nasal sinuses, and provide drainage for them when necessary; the patency of the Eustachian tube is maintained via their channels, and the olfactory sense is dependent on their healthy state. These nasal channels, lined by highly vascular ciliated epithelium, are often tortuous, frequently obstructed by large sharp spurs, and always intolerant of abuse.

It is unwise to regard every patient as possessing a nose which is a safe channel for the passage of a tube, or all types of case as being suitable for nasal intubation. As unwelcome complications, although infrequent, can follow this method, it is well that it should be reserved only for those cases in which it is clearly indicated and in which other methods, unattended by the same potentialities, have proved inadequate. It is not contested that nasal intubation, which owes so much to the genius of Dr. Magill, is an excellent procedure in selected cases and in the hands of those who realize the need for respecting certain contraindications. The enormous number of successful cases bear witness to its efficiency, and the following observations are merely a plea for a realization of certain difficulties and dangers which are met with, and an attempt to help in surmounting them. The subject is approached purely from the rhinological aspect, and no particular knowledge of anaesthetics is professed. It is the oto-rhino-laryngologist,

and not the anaesthetist, who is confronted with these occasional complications, not only because such complications are almost invariably the concern of his department, but because the cases most likely to be affected are those in which nasal and pharyngeal operations, and tonsillectomy in particular, have been performed.

It is clearly unwise to pass a tube through a nasal passage the interior of which is a closed book to the introducer; it is not enough to ask a patient if he can breathe through the chosen airway, as that will disclose neither a sharp spur nor an infected nose. More useful help can be gained by an examination with a headlight *after* shrinking with cocaine; this may reveal a passage in which trauma is unavoidable, and which not even the most gentle persuasion could negotiate with safety: trauma is the prerequisite of all intubation troubles.

Complications

Haemorrhage.—This may be caused by undue force being applied in a narrow nose or by coming up against an impassable obstruction: the highly vascular mucous membrane is abraded, and the seeds of possible infection are sown. In a few cases haemorrhage has been very considerable, and in one tonsillectomy case the surgeon was blamed because the patient lost as much as two pints of blood.

Infection.—This is secondary to abrasion of the mucosa, although it may be caused primarily by faulty aseptic technique. Though more likely to occur in an already infected nose—in which case an aggravation of symptoms will result—it may also be found in a hitherto perfectly healthy one. In tonsillectomy or septic pharyngeal cases infection is particularly likely to occur if trauma has taken place; but the causation in these cases is different, and it is, I believe, the *withdrawal* of the tube from an infected field over the abraded area in the nose which is responsible. It was the occasional virulent sinus infection after tonsillectomy in a war hospital that first attracted my attention, and I noticed that this sequence had occurred: haemorrhage from the intubated side of the nose, followed by an infection of the antrum on the same side, in cases in which radiographs and transillumination had shown the antrum to be healthy before operation. By withdrawal of the tube, on which have accumulated organisms and septic blood-clot resulting from the operation, and depositing these on the abraded mucosa or sinus ostia, infection is readily initiated. Such sinus or nasal infections greatly increase the patient's discomfort and prolong convalescence; they rarely clear up without intranasal drainage, and may necessitate a stay in hospital, in Service cases, of 4 to 6 weeks instead of the usual 7 to 10 days. Such nasal infections may encourage secondary haemorrhage from the tonsillar bed or nasal passage or lead to acute otitis media. It is indeed difficult to explain to the tonsil patient why the antrum should require puncture, the ear an incision, or the nose packing.

Adhesions between the two surfaces of the nasal passage, septal and turbinal, may follow if both of these have been traumatized in opposed areas.

Some Contraindications

(a) If the operation and anaesthesia can be in every way efficiently, safely, and satisfactorily controlled by other methods, nasal intubation is not the preferable method rhinologically.

(b) Large sharp spurs, marked turbinal hypertrophy, and gross septal deviations should be considered contraindications, particularly in cases of tonsillectomy or septic pharyngeal states. Purulent secretions and crusts, alar and vestibular infections, and large polypi should also debar the passage of a nasal tube.

(c) Septicaemic or very toxic patients are not suitable subjects for intubation; if intubated at all, the oral route is preferable, and particular gentleness is required, as a minimum of trauma to the nasal, laryngeal, or tracheal membranes is likely to be followed by a serious spread of infection.

(d) Tubes which have slipped into the oesophagus are likely to be infected, and are better not withdrawn through the nose or reinserted into the nasal passages, particularly if bleeding during introduction has occurred.

(e) In cases with a history of recurrent sinusitis the oral route is preferable.

(f) Cases in which one side of the nose is being operated upon are better not intubated via the remaining nostril; particularly if the lesion is an infective one, the oral route is again preferable.

(g) Children undergoing removal of tonsils and adenoids should not be pernasally intubated. It is unwise to pass a good-sized tube past a large adenoid pad; and, moreover, removal of adenoids, now often undertaken as the first step in the operation, is seriously interfered with by the presence of the tube in the nasopharynx. It is not difficult to remove tonsils with a tube through the mouth if intratracheal anaesthesia is desired, and this in no way interferes with adenoid removal.

Precautionary Measures

The following suggestions, most of which are well known but will bear repetition, are made in an endeavour to surmount some of the difficulties presented:

1. Careful sterilization of the hands, all instruments necessary for intubation, sprays, and lubricants is essential; these are best placed on a sterile towel spread over the patient's chest.

2. The nasal passages should be sprayed with 5% or 10% cocaine before anaesthesia is begun.

3. Inspection of the selected airway with a good headlight is advisable.

4. If there is resistance to the passage of a tube no attempt to force a passage is permissible. If much bleeding has followed the attempt the tube should be withdrawn and antiseptic ointment (sulphanilamide, for example) inserted. The tube should now be inserted, preferably by the oral route, though in some cases the use of the other nostril may be reverted to after inspection.

5. All blood-clot and secretions should be removed from the throat and post-nasal region in tonsil cases, preferably by suction, before the tube is removed; fragments of infected blood-clot are likely to be withdrawn along with the tube and deposited in the nose.

6. If it is known that trauma has occurred during intubation it is possible to withdraw the tube via the pharynx at the end of the operation. By grasping it with forceps as it lies in the posterior pharynx it can be withdrawn bent upon itself, from larynx and nose simultaneously; by this means the infected tube does not pass over the abraded mucosa. With soft tubes this manoeuvre is simple; with more rigid ones it is difficult. It is possible to cut the tube in half in the pharynx, and withdraw it in two pieces; it is needless to add that this suggestion will not meet with universal approval in these days of rubber shortage; but, nevertheless, it is an easy way out if the need is imperative.

MAINTENANCE TREATMENT WITH DIGITALIS

BY

ALFRED S. ROGEN, M.D., F.R.F.P.S.

(From the Department of Materia Medica and Therapeutics, University of Glasgow and Stobhill Hospital, Glasgow)

It is a well-established clinical practice to recommend the continuance of a maintenance dose of digitalis for patients with congestive cardiac failure who have responded well to the drug. Gold (1941) stated that "only about 15% of cardiac patients owe to the habitual use of digitalis the fact that they are able to carry on with a reasonable degree of comfort. The reputation of the drug is not enhanced, to say the least, by its use in the remaining 85% of the cardiac population." In Stobhill Hospital there has for some years been a special cardiac clinic for the supervision of patients with cardiac involvement who, after residence in the wards, have been found suitable for out-patient treatment. It has been a general rule, when digitalis therapy has proved necessary, to ascertain the adequate maintenance dose and to continue its administration. In some instances this treatment has been carried on for over four years. Patients attend at regular intervals, and many of them take an intelligent interest in their treatment. It was thought that these patients were excellent subjects for determining whether, in view of Gold's statement, we had been unduly liberal in dispensing digitalis.