described earlier in discussing dosage. Another way to produce sprains in knee cases is by failing to redevelop the vastus internus and then making it work against the same resistance as the rectus femoris component. This in its turn is due to failure to recognize the vastus internus as a separate muscle and therefore to use the entirely separate pulley circuit appropriate to its action.

The third type of progression in exercise therapy is progression in time—that is, carrying out the exercise under the same stress for increasing periods. This is very important in rehabilitating industrial workers, because industry demands not only that a workman shall be able to exercise his muscles against certain stresses but that he shall be able to do so continuously for whatever number of hours constitutes a "shift."

Principle of Variation in Form

If exercise therapy is to be carried out for many hours a day it must be disguised in various forms to make it palatable to the patient. Medical gymnastics alone would become boring; the patient would lose interest and initiative, and therefore cease to put his best endeavours behind them. So exercise therapy must be capable of assuming different forms, though in doing so it must conform as nearly as possible to the general principles already laid down. Unfortunately this is not always the case, and in the two chief disguises—occupational therapy and recreational therapy—the art of "sugar-coating" the pill has proved so diverting that the ingredients of the pill itself have been rather overlooked. Another reason why there must be variation in form is that medical gymnastics do not lend themselves to certain conditions. For example, no patient would sit in a gymnasium flexing and extending his fingers for six hours a day, but he will do it in an occupational-therapy department.

It is beyond the scope of this paper to discuss either occupational therapy or recreational therapy, but there are a few points in regard to both that bear on principles. Occupational therapy is used, of course, in two quite different ways—as a mental diversion (with which this paper is not concerned) and as a disguised form of remedial exercise. Employed in the latter sense, it becomes more important than ever to ensure that the proper muscles and joints are used, and that the patient is not simply learning trick movements. This can often be achieved by fixing the compensatory joints in plaster—e.g., in netting by fixing the metacarpo-phalangeal joints in order to make the patient use his proximal interphalangeal joints. Forms of craft-work should be favoured in which the movements are an alternate contraction and relaxation, such as weaving; those involving impaction stresses and sustained contraction should be avoided. Progression in range and power should not be lost sight of, and with ingenuity both can be incorporated into most forms of occupational therapy. For example, looms have been designed at Berry Hill Hall for shoulder, forearm, wrist, knee, and ankle movements, each with its own mechanism for ensuring effective muscle isolation and progression in both range and power.

The chief aim of the surgeon in all this should be not to acquire a detailed knowledge of occupational therapy but to satisfy himself that the essential principles of therapeutic exercise are being satisfied. Very little need be said about recreational therapy. It is a form of general exercise in which some sort of rough progression is possible, as, for example, in progressing from simple sitting games to walking and then running games. Games can never be very specific, nor is it desirable that they should be; their chief role is to redevelop general bodily function in a psychologically stimulating atmosphere, and the principal concern of the surgeon will be to protect his patients against the possible ill effects of their own enthusiasm or that of their fellows.

Summary

The basic principles of exercise therapy have been discussed and the following five requirements laid down:

(1) Exercises must be both focal and general. (2) They must be administered with due regard to correct dosage. This is variable from case to case and at different stages in the same case. (3) They must be rhythmic in character. (4) They must be progressive in range, power, and time. (5) They must

be variable in form, the chief forms being gymnastic, occupational, and recreational.

It is considered that a knowledge of these principles is all that is required or can be expected of the surgeon or physician who has to incorporate exercise therapy into his system of treatment. The detailed application and elaboration of principles is the province of the technical assistants in the various forms of exercise therapy.

JAUNDICE FOLLOWING ADMINISTRATION OF HUMAN BLOOD PRODUCTS

BY

H. V. MORGAN, M.B., M.R.C.P.

Physician, E.M.S.; Chief Assistant to the Medical Professorial Unit,
St. Bartholomew's Hospital

AND

D. A. J. WILLIAMSON, M.B., B.S.

Surgeon Lieutenant, R.N.V.R.; Late Chief Assistant,
St. Bartholomew's Hospital

At the present time human blood and its derivatives are being widely used for transfusion purposes. It therefore seems relevant to describe a small series of cases of jaundice which have followed the intravenous administration of plasma or reconstituted dried human serum.

Summary of Literature

Findlay and MacCallum (1937, 1938; with Murgatroyd, 1939) reported the occurrence of 89 cases clinically resembling infective hepatitis among 3,100 persons inoculated against yellow fever from 36 days to 7 months previously. The vaccine consisted of yellow fever virus suspended in human serum, which was obtained at first from patients who had recovered from yellow fever, but latterly from healthy donors without a past history of jaundice. The volume of serum injected in each case never exceeded 0.275 c.cm. These observers clearly demonstrated that the disease was not a form of yellow fever, and suggested that it was due to a hepatotoxic virus in the serum. Soper and Smith (1938) and Fox *et al.* (1942) recorded similar cases in South America.

A widespread outbreak has recently occurred among members of the American Forces inoculated against yellow fever (Memorandum on Yellow Fever Jaundice, 1942). 28,585 cases were reported during the first half of 1942, 62 of which were fatal. The clinical condition, which again resembled "catarrhal jaundice" except for the occurrence of joint pains and urticaria in 20% of cases, developed 40 to 120 days after inoculation. Necropsy showed acute or subacute necrosis of the liver, sometimes associated with nodular hyperplasia. Of particular interest is the complete absence of contact cases.

MacNalty (1937) reported 37 cases of acute infective jaundice, with seven deaths from acute hepatic necrosis, occurring 16 to 100 days after the administration of convalescent measles serum. All the 37 cases were from a group of 82 to 109 persons inoculated with serum of a particular batch. Probert (1938) in describing seven of these cases of infective hepatitis mentioned the occurrence of jaundice in two uninoculated children who two months previously had been in contact with some of the cases. Findlay and MacCallum quote several observers who have reported cases of hepatitis occurring in horses after inoculation against a number of equine diseases. The only common constituent of the inocula was horse serum (Theiler, 1919; Gordon, 1935; Marsh, 1937; Slagsvold, 1938).

Clinical Material

Late in May, 1942, an atypical case of jaundice occurred at an E.M.S. hospital (Case 3). A fortnight later a patient discharged from this same hospital (Case 4) was admitted to an associated hospital with a similar clinical condition. Comparison of the two cases showed that both had received intravenous serum several weeks previously. It was therefore felt that these might be examples of the type of jaundice caused by the administration of serum. An investigation of all cases known to have been transfused with serum or plasma in the past year was begun, and the following results were obtained.

During the period Aug., 1941, to June, 1942, 56 patients at the two associated hospitals are known to have had transfusions of liquid pooled plasma or reconstituted dried human serum. The majority of these were suffering either from peripheral vascular disease (Hayward, 1942) or from disorders associated with low plasma protein concentration. Most of the patients were otherwise in good general health at the time of transfusion and none was acutely ill. Their ages ranged from 15 to 78 years, though most were between 40 and 70. Of the cases which developed jaundice all but one (Case 8) were transfused at the sector hospital. The quantity of fluid transfused varied from 300 c.cm. to 6 litres, usually given in weekly amounts of 800 c.cm. Reactions consisting of pain in the back, vomiting, rigors, and pyrexia were not infrequently noted at the time of administration, but there was no obvious relation between the severity of the reaction and the occurrence of jaundice. The serum, which was obtained from the Medical Research Council, was prepared for use in the following manner:

The bottles of dried serum and of freshly distilled water were warmed to 37° C. in a water-bath. 200 c.cm. of the distilled water was then added to each bottle, which was gently shaken until all the serum had dissolved. The serum was thus reconstituted in normal strength and was administered through a Medical Research Council transfusion outfit.

Of the 56 patients, 50 have been traced, and 9 (18%) of them have since developed jaundice. Six of these cases were under our direct observation at the time of jaundice, but three (Cases 1, 2, and 8) were questioned retrospectively.

Case Reports

Case 1.—Railway signalman aged 59. Intermittent claudication and gangrene of toes due to degenerative vascular disease. Dec. 1 to Dec. 11, 1941: Three transfusions, each of 1,080 c.cm. of plasma. Dec. 23, 1941, to Jan. 9, 1942: Amputation of toes and then of leg under nitrous-oxide-oxygen-ether anaesthesia, followed by removal of dressing with trichlorethylene analgesia (inhalation of only 1 c.cm.). The wound was dressed with sulphanilamide powder. Feb. 7: Discharged to his home, where he had no known contact with any case of jaundice. Early in March, 1942 (approximately 90 to 100 days later): His wife noticed yellowness of sclera. Simultaneous onset of lassitude and anorexia; no rigors or sweating; no abdominal discomfort or vomiting. He was constipated, passing "white" stools and dark urine. No rash, no joint pains, no bruising. His condition remained unchanged for five to six weeks, when his symptoms cleared except for the jaundice, which persisted for a further month. There were no contact cases among his family or friends.

Case 2.—Labourer aged 41. Spondylolisthesis. Feb. 16, 1942: Transfused with 800 c.cm. serum. May 9 (82 days later), malaise. May 10: Developed abdominal discomfort and jaundice with dark urine and clay-coloured stools; diagnosed by his doctor as severe "catarrhal jaundice." June 5: Able to return to work, although the jaundice and abdominal discomfort persisted for a further four or five weeks.

Case 3.—Tool-maker aged 51. Intermittent claudication due to degenerative vascular disease; indolent plaster sore. Jan. 29 to March 17, 1942: Six weekly transfusions of serum, each of 800 c.cm. Plaster sore dressed with sulphanilamide powder. May 24 (68 to 115 days later): Generalized erythema and minutely papular rash on trunk and limbs. The rash, which lasted for three days, was not irritant and did not affect the head and neck. May 25: Progressive rise of evening temperature to a maximum of 103.2° F. on May 31, when the patient was noticed to be yellow. June 1: Jaundice deeper; liver palpable three fingerbreadths below the costal margin, and tender. Spleen impalpable. Urine contained bile pigments but not bile salts. Stools pale but not clay-coloured. W.B.C., 5,200 per c.mm. Blood sulphanilamide concentration too low to estimate. Serum van den Bergh reaction: Direct, positive in 2 minutes; indirect, 12 units. No abdominal symptoms. The temperature subsided during the next six days, but the jaundice disappeared more gradually. June 14: W.B.C., 8,200 per c.mm.—polymorphs 4,675, basophils 40, eosinophils 165, lymphocytes 2,910, monocytes 410. June 19: Liver impalpable; urine bile-free. Has remained well since. With the onset of jaundice there was a marked improvement in the rate of healing of his ulcer. Agglutination of *Leptospira icterohaemorrhagiae* negative. Blood Wassermann reaction negative.

Case 4.—Engineer aged 62. Intermittent claudication due to degenerative vascular disease. April 20, 1942: Transfused with 800 c.cm. serum. June 8 (49 days later): Malaise, anorexia, and abdominal discomfort lasting for three days. June 10: Noticed to be jaundiced. June 12: Admitted to hospital. Definitely jaundiced, with enlarged tender liver two fingerbreadths below costal margin. Spleen impalpable; urine contained bile pigments; and the stools

were pale. Except for temperature of 100° F. on the first day he remained apyrexial. June 17: W.B.C., 13,400 per c.mm.—polymorphs 9,510, basophils 130, lymphocytes 2,550, monocytes 1,210. Van den Bergh reaction: Direct, positive in 20 seconds; indirect, 6 units. June 22: Passed three loose melaena stools. The depth of jaundice fluctuated but improved. July 19: Discharged with liver just palpable. Aug. 4: Still slightly jaundiced and liver still palpable. Sept. 21: Free from jaundice; liver impalpable.

Case 5.—Cabinet-maker aged 62. Intermittent claudication due to degenerative vascular disease. March 5 to 26, 1942: Four weekly transfusions of serum, each of 800 c.cm. June 16 (82 to 103 days later): Malaise and anorexia for one week. June 22: Epigastric discomfort and dark urine. June 29: Admitted to hospital apyrexial and definitely jaundiced, with a large tender liver two fingerbreadths below costal margin. Spleen impalpable; stools pale; urine contained bile pigments. Tourniquet test positive. July 1: W.B.C., 4,200 per c.mm. Van den Bergh reaction: Direct, positive in 10 seconds; indirect, 10.5 units. July 9: More deeply jaundiced. Liver enlarged to three fingerbreadths and very tender. Spleen just palpable. Depth of jaundice varied, but gradually lessened. Aug. 10: Liver two fingerbreadths and tender. Spleen still palpable. Takata-Ara weak positive. W.B.C., 5,000 per c.mm. Sept. 5: Discharged with slight jaundice and a liver which was just palpable.

Case 6.—Publican aged 57. Intermittent claudication due to thrombo-angiitis obliterans. Sept. 8 to Oct. 7, 1941: Four transfusions of serum, each of 800 c.cm. Feb. 12 to April 1, 1942: Four transfusions of serum, each of 800 c.cm. June 27 (87 to 135, or 245 to 274 days later): Became jaundiced. Symptom-free and apyrexial apart from a rise in temperature to 99° F. on one evening. Liver and spleen impalpable; urine contained bile pigments; stools remained normal in colour. W.B.C., 17,000 per c.mm.—polymorphs 8,670, eosinophils 170, basophils 85, lymphocytes 6,205, monocytes 1,870. The patient's jaundice gradually subsided in the next three weeks. Blood Wassermann reaction negative. Agglutination of *L. icterohaemorrhagiae* negative.

Case 7.—Male aged 59; messenger. Diabetes mellitus with gangrene of toes. March 11 to May 7, 1942: Six transfusions, five of 800 c.cm. of serum and one of 1,080 c.cm. of plasma. July 20 (74 to 131 days later): Epigastric discomfort and flatulence for a few days; jaundice with tender enlarged liver three fingerbreadths below costal margin; apyrexial; urine contained bile pigments; stools pale. Aug. 2: Van den Bergh reaction: Direct, positive in 30 seconds; indirect, 7.5 units. Aug. 12: Discharged from hospital feeling well, with only slight jaundice; urine free from bile.

Case 8.—Female aged 45; secretary. Calcinosis circumscripta. May 16, 1941: Transfused with 300 c.cm. of serum. July 22 (67 days later): Anorexia, malaise, feverishness, and dark urine; pale stools. Aug. 5: Jaundice confirmed by her own doctor; this persisted for about 6 weeks.

Case 9.—Traveller aged 31. Intermittent claudication due to thrombo-angiitis obliterans. June 2 to 18, 1942: Three transfusions of serum, each of 800 c.cm. Oct. 1: Lumbar sympathectomy performed under nitrous-oxide-oxygen-ether anaesthesia; followed by pyrexia up to 100.8° F. for 24 hours. Oct. 3 (107 to 123 days later): Nausea, which lasted for a few days, and gradually increasing jaundice; liver enlarged; spleen impalpable; urine contained bile pigments; stools, normal in colour, contained no occult blood; apyrexial. Oct. 5: Blood van den Bergh reaction: Direct, immediate positive; indirect, 12.5 units. W.B.C., 8,900 per c.mm.—polymorphs 6,230, eosinophils 180, lymphocytes 2,135, monocytes 355. The jaundice gradually subsided in the next three weeks. Blood Wassermann reaction negative. Agglutination of *L. icterohaemorrhagiae* negative.

Epidemiology

Of these cases all but one (Case 8) were transfused in the sector hospital. There were no cases of infective hepatitis among patients in the wards of this hospital during the significant period. Although three nurses developed infective hepatitis during this period of seven months, they were working elsewhere in the hospital, and at no time had contact with the cases described. There were no cases of jaundice among the medical and nursing staffs of the wards concerned. All the patients who returned to their homes before developing hepatitis denied contact with any case of jaundice. The cases were distributed throughout four wards, and significant contact between them while they were jaundiced occurred in only one instance. Case 6 was in Ward MG1, and not confined to bed during the whole period in which Case 3 was jaundiced. He himself developed jaundice five weeks after the onset of prodromal symptoms in Case 3. There were 32 other patients in the ward at this time; of these only one (Case 9) has developed jaundice, and he only after an interval of three and a half months.

No other cases of jaundice have come to our notice among persons in contact with our patients.

Clinical Picture

The clinical features of these cases are summarized in the accompanying table. The onset of symptoms, which was sudden, occurred 49 to 107 days after the last transfusion. Jaundice, which appeared at the onset of the illness in five

at the time, and the only common causal factor was the administration of serum or plasma. It therefore seems probable that these fluids are of aetiological significance and that this outbreak is similar to those occurring after yellow fever and measles inoculation. Our cases differ, however, in that they were given relatively large quantities (300 to 6,400 c.cm.) intravenously, compared with very small amounts (0.275 c.cm.) intramuscularly. Most of our patients, too, were known to

Summary of Cases

Case	Age and Sex	Amount of Serum or Plasma (c.cm.)	Symptomless Period in Days	Symptoms and Signs								Pathological Investigations						
				Onset of Jaundice (Day of Disease)	Malaise, Nausea, and Abdominal Discomfort	Rash	Pyrexia	Liver Enlargement in F.B.s	Spleen	Stools	Melaena	Duration in Weeks	Total W.B.C. per c.mm.	Direct van den Bergh	Indirect (units)	W.R.	Takata-Ara	Leptospira Agglutination
1	59, M.	3,240 plasma	90-100 approx.	1st	Present several weeks	—	—	—	—	“ White ”	—	10	—	—	—	—	—	—
2	41, M.	800 serum	82	2nd	Present several weeks	—	—	—	—	Clay-coloured	—	8-9	—	—	—	—	—	—
3	51, M.	4,800 serum	68-115	8th	—	+	Up to 103° F. 1st fortnight	3 F.B.s. Tender	—	Pale	—	4	5,200	+	12	—	—	—
4	62, M.	800 serum	49	3rd	Present three days	—	100° F. on 1st day	2 F.B.s. Tender	—	Clay-coloured at times	+	7½	13,400	+	6	—	—	—
5	62, M.	3,200 serum	82-103	7th	Present one week	—	—	3 F.B.s. Tender	+	Pale	—	12	4,200	+	10.5	Weak +	—	—
6	57, M.	3,200 + 3,200 serum	87-135 or 245-274	1st	—	—	99° F. on 1st day	—	—	Normal	—	3	17,000	—	—	—	—	—
7	59, M.	4,000 serum + 1,080 plasma	74-131	1st	Present a few days	—	—	3 F.B.s. Tender	—	Pale	—	3½	—	+	7.5	—	—	—
8	45, F.	300 serum	67	1st	Present one week	—	—	—	—	Pale	—	6	—	—	—	—	—	—
9	31, M.	2,400 serum	107-123	1st	Present a few days	—	Post-oper. only	3 F.B.s	—	Normal	—	3	8,900	+	12.5	—	—	—

cases, was intense in some patients, and often varied in degree from day to day. Malaise, nausea, and epigastric discomfort, which sometimes persisted for some weeks, accompanied the jaundice in seven cases and preceded its onset by a few days in three. In only one case (Case 3) was a rash observed: it preceded the onset of jaundice by eight days. None of our patients complained of joint pains or biliary colic. In all but one of the cases we examined the liver was enlarged and tender; but in only one (Case 5) was the spleen palpable. In no case was ascites demonstrated. There was a positive direct van den Bergh reaction in five cases and unduly pale stools in seven. The severity of the illness, which bore no relation to the volume of fluid transfused, varied from the symptomless jaundice of Case 6 to the severe illness with intestinal haemorrhage of Case 4. On the whole the cases were of moderate severity, and only Case 4 gave rise to anxiety. The duration of the illness varied from 3 to 12 weeks.

Diagnosis

The clinical picture, though showing considerable variation, most closely resembles that of infective hepatitis as described by Cullinan (1939); it does, however, differ in the following respects:

(1) Our patients belong to a considerably older age group than that usually affected (Newman, 1940; Norton, 1939), although Pickles (1939) states that infective hepatitis is not uncommon among older people. Fox *et al.*, in describing an outbreak of jaundice following yellow fever inoculation, report a significantly higher incidence among the older age groups. (2) The illness was, on the average, of greater severity and more protracted. (3) In only one (Case 6) was there a history of contact with a possible source of infection. (4) No cases of jaundice have been reported among persons in contact with our patients.

We were not led to expect a subacute necrosis due to any of the better-known hepatotoxic chemicals. The absence of relapses in our patients is also against this diagnosis.

The clinical picture of our cases does not resemble spirochaetosis icterohaemorrhagica: there was no albuminuria or marked polymorphonuclear leucocytosis, and the serum from three cases gave negative agglutination reactions.

Discussion

From the clinical and epidemiological evidence it seems unlikely that this is a straightforward outbreak of infective hepatitis. These were the only cases of hepatitis in the hospital

be suffering from arteriosclerosis, and might conceivably have had some liver change, rendering them more susceptible to hepatotoxic agents. In two of the cases, however, there was no evidence of vascular disease.

There are three main possibilities as to the part played by the serum in the production of jaundice:

1. Hypersensitivity Reaction.—There was no direct evidence in favour of this view: our patients did not give a history of allergy and showed no eosinophilia. The time relationship would appear to be too long for an anaphylactic type of reaction.

2. Introduction of a Hepatotoxic Agent.—In some of the recorded outbreaks there has been evidence that only certain batches of serum have been hepatotoxic. In our series comparatively few of the batch numbers are available, and in only two cases (4 and 9) do we know all the numbers concerned. In these two cases there was one common batch (120 KY), which was also administered to Case 7. Serum from this batch was given to at least two other patients, who did not develop jaundice. Batch 120 KY was derived from a pool of 50 litres of serum, a contribution of 250 donors, whom it was not possible to trace. It is probable that Case 8 did not have serum from this batch, and Case 1 was transfused only with plasma. There is therefore no evidence of an abnormality confined to one batch of serum. If a hepatotoxic agent has been introduced it must be one which is common to both dried serum and liquid pooled plasma. It might be either a virus contained in the fluid, as suggested by Findlay and MacCallum, or some toxic chemical substance such as altered plasma protein. In our series there is no evidence to favour one or other of these hypotheses.

3. Increased Liver Susceptibility.—Certain agents, such as neoparsphenamine (Stokes, Reudemann, and Lemon, 1920) and acriflavine (Murray, 1930), are believed to reduce the resistance of the liver, rendering it more susceptible to the virus of infective hepatitis. It is possible that our cases represent an outbreak of this disease, with the serum acting as a toxic agent, but this seems unlikely in view of the lack of infectivity and the long symptomless period.

Summary

Nine cases of jaundice are described which developed 7 to 16 weeks after transfusion with serum and plasma, and closely resembled the hepatitis following inoculation against yellow fever and measles.

The association of such cases of jaundice with transfusion may easily be overlooked in view of the long symptomless period which intervenes.

We wish to thank various members of the staff of St. Bartholomew's Hospital for permission to publish the cases under their care, and gratefully acknowledge the helpful criticism of Dr. A. E. Gow, Prof. Ronald V. Christie, and Dr. Hugh Barber.

REFERENCES

- Cullinan, E. R. (1939). *Proc. roy. Soc. Med.*, **32**, 933.
Findlay, G. M., and MacCallum, F. O. (1937). *Trans. roy. Soc. trop. Med. Hyg.*, **31**, 297.
— (1938). *Proc. roy. Soc. Med.*, **31**, 799.
— and Murgatroyd, F. (1939). *Trans. roy. Soc. trop. Med. Hyg.*, **32**, 575.
Fox, J. P., Manso, C., Penna, H. A., and Pará, M. (1942). *Amer. J. Hyg.*, **36**, 68.
Gordon, W. S. (1935). *Quoted Proc. roy. Soc. Med.*, **31**, 799.
Hayward, G. W. (1942). *British Medical Journal*, **1**, 285.
MacNalty, A. S. (1937). *Ann. Rep. Chief Med. Off. Min. Health*, p. 38, London.
Marsh, H. (1937). *J. Amer. vet. med. Ass.*, **91**, 88, 330.
Memorandum on Yellow Fever Jaundice (1942). *J. Amer. med. Ass.*, **119**, 1110.
Murray, D. H. (1930). *J. R.A.M.C.*, **54**, 19.
Newman, J. L. (1940). *Med. Officer*, **64**, 21.
Norton, J. A. (1939). *J. Amer. med. Ass.*, **113**, 916.
Pickles, W. N. (1939). *Epidemiology in Country Practice*, London.
Probert, S. A. (1938). *British Medical Journal*, **2**, 677.
Slagsvold, L. (1938). *Norsk vet. Tidsskrift*, **50**, 69.
Soper, F. L., and Smith, H. H. (1938). *Amer. J. trop. Med.*, **18**, 111.
Stokes, J. H., Reudemann, R., and Lemon, W. S. (1920). *Arch. intern. Med.*, **26**, 522.
Theiler, A. (1919). *Rep. vet. Res. S. Africa*, **5**, 7.

A NEW AND SIMPLE METHOD FOR SUBCUTANEOUS LIGATURE OF VARICOSE VEINS

BY

B. W. GOLDSTONE, M.B., B.Sc., F.R.C.S.Ed.
Major, R.A.M.C.

The injection of various solutions into varicose veins often causes their obliteration; but this technique is generally useless when the upper part of the great saphenous vein is varicose: the downward pressure of this long column of blood unfortunately unseals the attempt at thrombosis.

In the past decade surgical opinion has recognized this factor, and operations for high ligation of the vein have become standardized. These are usually, and rightly, combined with injection of the vein's lower reaches. This combined operation is generally very successful. Lately, several surgeons have practised a still simpler technique: the vein is compressed subcutaneously instead of by open operation. I use the word "compressed" advisedly, as will appear later from the text, although these others refer to their methods as ligation. By this technique a curved needle is thrust into the skin at the side of the vein; it passes deep to the vessel and reappears at the other side of it. The two free ends are then tied over a pad of gauze. Below this the varicosities of the lower leg are injected.

I find that there are several marked advantages in these subcutaneous methods, particularly from a military point of view. First, there are a large number of men who obstinately refuse to be "cut about." When it is explained to them that the cure (I carefully avoid the word "operation") entails absolutely no cutting, they then find that they can furnish no adequate reason against being cured. Also, the method, being ambulatory, effects a great saving in hospital beds. With an open operation, even after the patient leaves the ward, he must remain relatively immobile until the wound has healed strongly. The use of the subcutaneous operation should result in raising thousands of soldiers into a higher category.

The method hitherto used is described as subcutaneous ligation. It is certainly subcutaneous, but it is a unilateral compression, not a ligation. A true ligation should evenly compress the vein's entire circumference towards its centre. A much more effective and permanent result is likely to follow from a true ligation. The simple method described below forms a true ligation around the vein. In addition the little operation is complete at one sitting. Patients need not return to the clinic for removal of stitches. Usually I tell them to return, but it is more for my interest than in theirs.

Method

The patient stands and the highest visible part of the internal saphenous vein in the thigh is shown; but it is advisable to

keep at least 2 in. below the saphenous opening. A local anaesthetic wheal is raised on each side of the vein. A curved cutting needle is then threaded with catgut. It is plunged into one wheal, passes deep to the vein, and emerges through the skin via the opposite wheal (Fig. 1). This curved needle (Fig. 2) is now removed from the catgut and a straight needle

FIG. 1

FIG. 2

FIG. 1.—Stage I: Curved needle passes through skin, then deep to vein, to emerge through skin at opposite side. Thus the two skin punctures A and B are made.

FIG. 2.—Stage II: Curved needle has emerged. A straight needle must now be substituted on to the catgut.

substituted on to the catgut. This straight needle is now thrust back into the skin via the same minute opening through which the curved needle has recently emerged. The straight needle passes in front of the vein and emerges through the same minute hole where the curved needle made its original entry (Fig. 3).

FIG. 3

FIG. 4

FIG. 3.—Stage III: Straight needle has re-entered skin via B and is about to emerge via A. Here needle passes in front of vein.

FIG. 4.—Stage IV: Straight needle has emerged and the two original ends of the catgut are ready to be tied together with a reef-knot, forming a true ligation around the vein.

The two ends of the catgut are once more in juxtaposition (Fig. 4); they are tied fairly tightly and cut short; the knot retracts and almost disappears. The little operation is complete. It is extremely simple to do; in fact, reading about it demands more effort than actually doing it. The area receives no dressing other than a small piece of bare elastoplast. This is removed in one week, when the catgut knot will have disappeared spontaneously. There will now be a minute puncture and the surrounding skin will have a slightly traumatized appearance, but no further treatment is necessary; in a short time the skin regains its normal appearance and elasticity.

A few days after the original ligation the varicose veins below the knee should receive one or more injections as necessary, but great caution should be observed; only a small quantity of the thrombogenic fluid should be injected at first (1 c.cm. of 5% sodium morrhuate would be a good initial or test dose). The reason for this cautious and small injection is obvious: with the main internal saphenous vein ligatured the venous current is so slow that dilution or washing away of the chemical hardly occurs; from a minimal dose maximal results are obtained.

For a week after the ligation patients can move about and are quite fit for light duties. After that they rapidly become normal.

Discussion

For the past few months I have carried out this simple procedure on every suitable case that has come under my care, the results being invariably successful. I must admit that as yet I have had no large series of cases. The procedure is so simple that I have already begun to ligate the larger varicosities of the lower leg in preference to their injection, though care is exercised in avoiding flexures.