I should be the last to deny that all of these things can cause death when they are severe enough. But it must be strongly asserted that the findings in human beings are often disappointingly small—disappointing, that is, to the mind which is prepared only for anatomical findings. Yet it is not these things, nor the petechial haemorrhages of classical times, which tell us most; it is rather their variability or their absence. To find nothing would be in line with the most informative of experimental evidence. This raises the question as to how it can come about that an initial lesion severe enough to cause neural paralysis can be survived though death occurs hours or days afterwards. There is no question about the fact that this can and does happen; and it is certain, it is sure. Only one explanation suggests itself—that an injury sets going a self-propagating event that leads sooner or later to death. It would be a mistake to draw too much on analogy, and any explanation must be regarded as both speculative and tentative. But death from burns, from fractures, and from haemorrhages will serve as shorthand notes indicating at least a parallelism. We can only speculate at the moment on what the answer will be in the case of cerebral contusion. It is the fate of all of us to watch patients with cerebral lesions dying from something characterized by tachycardia and cardiac irregularity, from upsets of thermal regulation, from metabolic disturbance which we have not yet found a means to study properly, from something that bears the hall-mark of a disease process as much as, say, pneumonia. The old beliefs in the omnipotence of the bulbar centres has been corrected by the discovery of the vital centres re-representing them at and above the tentorial level. Conceivably it may become possible to demonstrate alteration in oxidation-reduction mechanisms in this area which will afford an index of cellular pathology. It seems legitimate to suppose that the physical effects of injury lead to affection of at least one enzyme system, the lack of which can for a time be repaired by the altered energies of others, which also in turn give out. Thus, step by step, dissolution comes about. Only in such a way can temporary survival with final extinction be given a cause. And it will be recognized that the level at which all of these happenings occur is in the more primitive but still no doubt extremely complex and largely integrated neural mechanisms considerably below cortical level.

Conclusion

Although the beliefs expressed above have minimized the importance of the cortex in the picture of concussion, it is not my belief that cortical disturbances are absent. Electrical records clearly show that they are frequent; while some patients, by confusion, confabulation, and lack of insight (sometimes for many days after return to consciousness), demonstrate that the whole brain integration continues to be disturbed. It is not possible at the moment to do more than speculate on the causes and mechanisms of this dysfunction. The real point that I have wished to examine is the nature of unconsciousness, and I have endeavoured to show that there is a body of evidence attesting to its production in a relatively focal manner, and that the focus is not, as has been so generally believed, in the cerebral cortex.

BIBLIOGRAPHY

- Breslau, J. (1914). *Arch. klin. Chir.*, **103**, 478.
 Cairns, H., Oldfield, R. C., Pennybacker, J. B., and Whitteridge, D. (1941). *Brain*, **64**, 273.
 Denny-Brown, D., and Russell, R. (1941). *Ibid.*, **64**, 93.
 Fischer, H. (1870). *Samm. klin. Vortr.*, No. 27, 1st ser., 119.
 Fulton, J. F., and Bailey, P. (1929). *J. nerv. ment. Dis.*, **69**, 1.
 Globus, J. H. (1940). *Arch. Neurol. Psychiat.*, Chicago, **43**, 125.
 Greenfield, J. G. (1938). *Proc. roy. Soc. Med.*, **32**, 43.
 Harrison, F. (1940). *The Hypothalamus*, **20**, 635, Assoc. Res. Nerv. and Ment. Dis.
 Jefferson, G. (1938). *Arch. Neurol. Psychiat.*, Chicago, **40**, 857.
 — (1942). *Glasg. med. J.*, **138**, 77.
 Kleitman, N. (1939). *Sleep and Wakefulness*, Chicago.
 Knauer, A., and Enderlen, E. (1922). *J. Psych. Neurol.*, **29**, 1.
 Kocher, T. (1901). *Hirnerkrankung*, etc., Vienna.
 Miller, G. (1927). *Arch. Surg.*, Chicago, **14**, 891.
 Penfield, W., and Erickson, T. (1941). *Epilepsy and Cerebral Localization*, p. 143, Springfield, Ill.
 Ranson, S., and Magoun, H. (1939). *Ergebn. Physiol.*, **41**, 56.
 Symonds, C. P. (1940). Concussion and Contusion of the Brain, in Brock's *Injuries of the Skull, Brain and Spinal Cord*, Baltimore.
 Trotter, W. (1932). *Choyce's System of Surgery*, **3**, 363, London.
 Von Economo, C. (1926). *Handb. norm. path. Physiol.*, **17**, 591.
 — (1930). *J. nerv. ment. Dis.*, **71**, 249.
 White, J. C. (1940). *The Hypothalamus*, **20**, 855, Assoc. Res. Nerv. and Ment. Dis.
 Williams, D., and Denny-Brown, D. (1941). *Brain*, **64**, 223.
 Witkowski, L. (1877). *Virchows Arch.*, **69**, 498.

NUTRITIONAL DEFICIENCY IN THE PATHO-GENESIS OF DISEASE*

BY

JOHN YUDKIN, M.A., M.D., Ph.D., F.I.C.

Lieut. R.A.M.C.; formerly Sir Halley Stewart Research Fellow at Dunn Nutritional Laboratory, Medical Research Council and University of Cambridge

More and more has the role of vitamin deficiency claimed our attention in recent years, and a variety of diseases and syndromes have been treated with a vitamin or a combination of vitamins. Frequently the reasons for doing this are obscure and the results of the treatment equivocal; and the relation between the condition and vitamin deficiency often remains difficult to assess.

Nutritional deficiency may be responsible for the production either of a recognized disease, of hitherto unknown aetiology, or of obscure signs and symptoms in an individual. For example, claims are sometimes made that vitamin deficiency plays a part in the production of renal lithiasis (Higgins, 1935) and certain muscular dystrophies (Bicknell, 1940). Again, a patient is seen with an unusual form of keratosis of the skin, or with oedema which does not easily respond to ordinary therapy, or with a sore tongue with few other signs or symptoms, or with an unusual type of subcutaneous haemorrhage: to what extent are vitamin A, vitamin B₁, riboflavin, and vitamin C respectively implicated in the production of these disturbances in the individual?

In this paper I shall consider certain general principles which might act as a guide in the evaluation of the part played by vitamins in the causation of disease. Just as we use Koch's postulates in determining the role of specific bacteria in causing disease, so we may lay down the following criteria for assessing the role of nutritional deficiency in causing disease: (1) It should be possible to demonstrate the existence of a deficiency of the nutrient; (2) deficiency of the nutrient should result in the production of the disease; (3) the disease should be cured by replacement of the deficient nutrient. To satisfy each of these criteria,* evidence may be collected in several ways:

Diagnosis of Nutritional Deficiency

Criterion 1.—Existence of deficiency

A. General evidence of deficiency

- (i) Evidence of inadequate diet
- (ii) Evidence of defective absorption
- (iii) Evidence of increased utilization

B. Specific evidence of deficiency

- (i) Laboratory tests for specific deficiency
- (ii) Clinical tests for specific deficiency

Criterion 2.—Deficiency as the cause of the condition

- A. Evidence from animals
- B. Evidence from human beings

Criterion 3.—Cure of condition by correction of the deficiency Therapeutic test

Criterion 1: Existence of Deficiency

The first criterion is that there should be evidence of nutritional deficiency in the condition. This evidence may be either general or specific.

A. General Evidence of Deficiency

There may be evidence of a deficiency in any part of the pathway between the nutrient and the tissues.

(i) Evidence of Inadequate Diet

A dietary history is an essential first step in diagnosis. The following conditions are especially apt to lead to an inadequate diet.

(a) *Low Income Levels.*—On the whole, foods which are good sources of vitamins are also more expensive; cheaper foods supply mainly calories (Orr, 1936).

(b) *War.*—Wartime measures, such as the introduction of the national loaf and the vitaminization of margarine, together with rationing, the price control of foods, and the distribution of cheap or free milk and fruit juices, have decreased the disparity between the diets of various classes. Nevertheless, there are many who still

* Based on a paper read at a meeting of the Nutrition Society (Scottish Branch) at Gleneagles, Aug. 26, 1942.

† Certain exceptions to the fulfilment of these criteria are discussed below.

cannot afford to buy the whole of their meat or bacon ration or to spend much on vegetables. Moreover, since war leads to restricted imports, to a decreased man-power for food production, and to the destruction of crops and other foodstuffs, it may be a potent cause in the production of deficiency disease.

(c) *Season*.—An example of the part played by this factor is the low level of vitamin C in foods in the spring and early summer. It is therefore at this time that the effects of its deficiency are most likely to be seen (Harris, 1942).

(d) *Geography*.—Differences in food production and dietary habits lead to the classical examples of geographic endemic deficiency disease such as the association of beriberi with the consumption of polished rice and pellagra with the consumption of maize.

(e) *Special Diets*.—Possibly the commonest cause of adult scurvy in Britain is the consumption of a too rigid "gastric" diet by patients with peptic ulcer, without adequate precautions for assuring the supply of vitamin C (Harris *et al.*, 1936). Also, beriberi has been reported in food cranks who have lived on a diet containing excessive amounts of highly purified carbohydrates. It is possible that the well-known occurrences of pellagra in mental hospitals are partly due to the perverted food habits of the patients.

(f) *Institutions*.—Outbreaks of deficiency diseases are often reported from institutions; nor is this surprising in prisons and asylums, where the allowance for food may not be sufficient to supply even the necessary calories. But the unique nature of institutional feeding may result in an inadequately balanced diet even if economic considerations do not restrict the choice of food. In the first place, the food and its preparation, for a large number of people, are in the hands of a few individuals, who may easily, through ignorance or lack of care, be responsible for a diet of poor quality. Secondly, the preparation of food on a large scale introduces new problems in connexion with the preservation of its nutritive quality; it requires special care, for example, to preserve the vitamin C (King *et al.*, 1942).

(ii) Evidence of Defective Absorption

(a) *Water-soluble Vitamins*.—Many affections of the gastrointestinal tract appear to decrease the absorption of vitamins of the B group. Hyperemesis gravidarum (Strauss and McDonald, 1933) and alcoholic gastritis (Minot *et al.*, 1933) are often associated with deficiency of vitamin B₁; in the latter condition there is probably a second factor in the diminished intake of the vitamin in the diet. Gastrectomy, colitis, and intestinal obstruction often lead to a complex deficiency of the vitamins of the B₂ group (B.M.J., 1942).

(b) *Fat-soluble Vitamins*.—The absorption of these substances is bound up with the absorption of fat. In obstructive jaundice, in which the absorption of fat is decreased, there is diminished absorption of vitamins A (Brees and McCoord, 1940) and K; it is known that the difficulty in controlling haemorrhage in obstructive jaundice is due to this "conditioned" deficiency of vitamin K (Warner *et al.*, 1938a, 1938b). The effect of large doses of liquid paraffin in hindering the absorption of carotene is another example of induced defective absorption of fat-soluble vitamins (Rowntree, 1931).

(iii) Evidence of Increased Utilization

(a) *Pregnant and Lactating Women*.—In some degree pregnancy and lactation increase the demand for all the nutrients, including the vitamins. Osteomalacia and beriberi are examples of deficiency disease which may be precipitated by these increased physiological needs.

(b) *Heavy Manual Workers*.—It is likely that the utilization of many nutritional factors is greater with increased physical exertion. Many Europeans consume diets which probably contain no more vitamin B₁ than those consumed by many Chinese; the high incidence of beriberi among the latter is probably associated with the very heavy work which they do.

(c) *Fever*.—There is apparently an increased destruction of vitamin C and possibly other vitamins during fever (Abbasy *et al.*, 1936, 1937; Harris *et al.*, 1937). Usually the primary condition, such as tuberculosis or rheumatic fever, overshadows deficiency of the vitamin, but this deficiency may play a part in delaying recovery or in the persistence of certain signs.

B. Specific Evidence of Deficiency

This evidence may be obtained either by laboratory tests or by clinical tests.

(i) Laboratory Tests for Specific Deficiency

(a) *Vitamin Concentration in the Blood*.—The determination of the blood concentration of vitamin A, B₁, or C has often been used to assess the nutritional level in respect of these factors. The results of such investigations, however, appear to depend to a large extent on the immediate past diet, and give less indication of the more permanent nutritional state. A very short time on a diet high in vitamin A, for example, will lead to high blood levels although the subject still shows signs of deficiency (S. Yudkin, 1941); the converse is also true. To some extent this has been recognized with

vitamin C: estimations on the whole blood or plasma have been found to be less reliable as indices of vitamin C nutrition than estimations on the white-cell layer (Heinemann, 1938, 1941).

(b) *Vitamin Excretion in the Urine*.—With the water-soluble vitamins, the amount excreted in the urine, and especially the rapidity with which large doses are excreted, have been useful methods for detecting slight deficiency (Harris and Ray, 1935). Such tests have been employed mainly with vitamin C, but they are also proving of value in the diagnosis of deficiency of vitamin B₁ (Banerji and Harris, 1939; Wang and Yudkin, 1940) and nicotinic acid (Kodicek and Wang, 1941).

(c) *Functional Tests*.—Slight deficiency of a vitamin often causes physiological or biochemical changes, which may be detected by the use of special techniques. Impaired dark-adaptation, increased capillary fragility, or a rise in blood phosphatase has been used to demonstrate mild deficiency of vitamins A, C, and D. A rise in pyruvic acid in the blood or urine, particularly after the ingestion of glucose, has been suggested as a means of recognizing early deficiency of vitamin B₁ (Banerji and Harris, 1939). Unfortunately these methods, although of great value, have been used very uncritically by some workers, and the published results have often been conflicting. The main reason for this is that tests such as those on dark-adaptation and capillary fragility appear superficially very simple to employ, but in reality require most careful application and rigid control (Hecht and Schlaer, 1938; J. Yudkin, 1941; *Lancet*, 1942; Yudkin and Ferguson, 1943).

(ii) Clinical Tests for Specific Deficiency

Clinical examination may reveal signs which are similar to those seen in the known deficiency disease, or they may be found in association with such signs.

(a) *Signs Resemble Those in Recognized Deficiency Disease*.—The realization that the neuritis of alcoholism and that of pregnancy toxæmia were due to deficiency of vitamin B₁ came, partly at any rate, from the recognition of the similarity between these and the signs of mild "dry" beriberi. But this type of evidence must be used carefully: it has led, for example, to the quite illogical use of vitamin B₁ in neuritis of many sorts, including sciatica (Roch and Sciclounoff, 1938).

(b) *Signs Associated With Those of Recognized Disease. In Individuals*.—In individuals with signs of a deficiency disease the presence of other signs whose pathogenesis is obscure suggests that these are also due to the same deficiency. Thus obscure neurological signs found in Cingalese prisoners appear to be due to deficiency of vitamin A, since other signs of A deficiency—e.g., phrynodermia—were also present; moreover, beriberi is uncommon in Ceylon, so that the neurological signs were unlikely to be caused by deficiency of vitamin B₁ (Nicholls, 1935). In view, however, of the frequency with which multiple deficiency occurs, caution must be used in interpreting such findings. Until recently, for example, it was thought that some of the changes in the skin and tongue in pellagra were due to the lack of the pellagra-preventing factor. Only after the latter was isolated as nicotinic acid was it found that these signs were due to a superimposed deficiency of riboflavin (Young, 1937-8; Sydenstricker, 1941). Incidentally, this type of evidence can be used in the opposite way: if certain obscure signs are rarely found in association with signs of a deficiency disease it is unlikely that they are due to this deficiency. Thus it was at one time suggested that disseminated sclerosis and subacute combined degeneration of the cord were due to deficiency of vitamin A, but the fact that other signs of this deficiency are rarely found in these conditions makes it unlikely. *In a Population*.—If a deficiency disease is common in a certain population the same deficiency may often be the cause of other less well defined conditions. Much of the so-called toxæmia of pregnancy in Hong Kong, where beriberi is common, has been shown to be due to deficiency of vitamin B₁ (Nixon, 1942).

Criterion 2: Deficiency as Cause of Condition

Our second criterion—that the signs should be reproducible by a deficiency of the suspected factor—may be applied either in experimental animals or in human beings.

A. Evidence from Animals on Experimental Deficient Diet

Analogy from animal experiments has proved very useful in recognition of signs of human deficiency. The production of haemorrhage due to lack of vitamin K (Dam and Schönheyder, 1934) and vascularization of the cornea due to lack of riboflavin (Wolbach, 1937; Bessey and Wolbach, 1939) are examples of this. On the other hand, such analogy must not be carried too far, as has been done with vitamin E and the production of neuromuscular disease (Bicknell, 1940) or vitamin A and the production of neurological disease (Mellanby, 1934).

B. Evidence from Human Beings on Experimental Deficient Diet

As might be expected, there are many difficulties in conducting such experiments on anything like an adequate number of subjects,

and it is not surprising, therefore, that few have been reported. It would be expected that the results would be very convincing, but in fact the conclusions reached are often conflicting. Thus some workers have claimed that a dietary deficiency of vitamin A produces a deterioration in dark-adaptation within a few days (Hecht and Mandelbaum, 1940), while others have found no change after five or six months (Steffens *et al.*, 1939). It is possible that these divergent results are due to such factors as differences in individual requirement, differences in the stores of the vitamin before the experiment is begun, and differences in the composition of the deficient diet—that is, the possible concurrent existence of other deficiencies.

Criterion 3: Cure of Condition by Correction of Deficiency Therapeutic Test

Wherever possible, the effect of correction of the supposed deficiency should be observed. Here again, however, it is unfortunately true that many reports in the literature of the cure of symptoms and signs by the administration of a vitamin are vitiated by the failure to observe certain general principles. The commonest source of error is the use of inadequate controls, and especially the failure to allow for spontaneous remission. On the other hand, even if deficiency really exists, negative results may follow the administration of the missing factor. Often this is due to inadequate dosing or to defective absorption; in such cases large doses, if necessary given parenterally, must be tried. Failure to cure may also be due to the fact that the pathological process has progressed too far; it is not always possible, for example, completely to cure long-standing peripheral neuritis of beriberi by the administration of vitamin B₁.

A less well recognized reason for the failure to achieve improvement by the correction of an existing deficiency is the simultaneous existence of other deficiencies. It is often overlooked that a diet is deficient in several factors, that the administration of only one of these may produce no improvement, and that this does not imply that the diet must necessarily have been adequate in this factor. For example, Peterhead children on diets undoubtedly deficient in vitamins A and D as well as many other nutrients were no better than the controls when given these substances (Sutherland, 1934). It was accepted in this experiment that deficiency of one or more other nutrients was the limiting factor in preventing improvement. Similar results were recently obtained in experiments on Cambridge school children (Yudkin, 1943). It is probable that the lack of effect of supplements of vitamin C in the oft-quoted experiment with South African miners (Fox *et al.*, 1940) was due to the similar existence of multiple deficiency; restoration of an adequate supply of this one substance without increasing the supply of the other nutrients was itself not sufficient to affect the health of the workers, apart from decreasing the number of cases of frank scurvy.

Lastly, when a positive effect is obtained by the use of a vitamin supplement it is not always legitimate to assume that this has proved the existence of deficiency. The fact that vitamin C is diuretic and hence relieves the oedema of cardiac failure (Evans, 1938) does not of course imply that the condition is due to deficiency of the vitamin. Again, if, as has been suggested, large doses of vitamin D have a favourable effect on psoriasis (Brunsting, 1938) it is not to be inferred that psoriasis is a deficiency disease. It is important, in other words, to distinguish between replacement therapy and pharmacological therapy.

General Comments

Diagnosis by rule of thumb, without consideration of biological realities, is to be deprecated no less with deficiency disease than it is with any other condition. We may mention here some of the more important considerations which must be borne in mind.

Multiple Deficiency.—Human dietaries are such that deficiency of one nutrient only is unlikely to occur. This means that Criterion 1 may indicate deficiency of a nutritive factor which is not the cause of the condition under investigation. It would appear that the signs which will occur in a multiple deficiency will depend partly on the relative extent of the various deficiencies and partly on the capacity of the organism to store the different dietary factors. On a diet completely deficient in the vitamin B complex, for example, rats will almost certainly die of deficiency of vitamin B₁ before developing signs of other deficiencies.

Individual Variation in Requirement.—It cannot be expected that all individuals have exactly the same requirements for the various nutritional components or that they would show the same signs when depleted of one or more of these components. It is possible that there are also racial as well as individual differences, as illustrated by the reported behaviour of

Rumanian and Russian prisoners in a German camp during the last war. It is stated (Bigwood, 1942) that all were on the same diet, yet the former developed pellagra and the latter scurvy.

Other Constituents of the Diet.—It has been shown that signs of deficiency of riboflavin can be considerably aggravated if the deficient subjects are dosed with other components of the B complex (Sebrell and Butler, 1938; Sydenstricker *et al.*, 1940), and there is similar experimental evidence in dogs (Morgan, 1941). This provides another reason for considering the whole of the diet in attempting the diagnosis of deficiency disease.

Specificity of Tests for Deficiency.—No single test can be held to prove deficiency. There are, for example, other causes of poor dark-adaptation than deficiency of vitamin A; other factors producing increased capillary fragility than deficiency of vitamin C. It is possible also to have a high level of vitamin A in the blood coexisting with definite functional deficiency of this vitamin.

Finally, it should be stressed that, even if deficiency has been proved, it may be the effect rather than the cause of the condition. An extreme example is deficiency of vitamin C in tuberculosis, in which it is clear that no one would suppose that the deficiency causes the infection. On the other hand, it should be remembered that, even when it is not the ultimate cause of a definite pathological state, deficiency may be a predisposing cause. It is likely, for example, that deficiency of either vitamin C or nicotinic acid renders the individual more readily susceptible to ulcerative stomatitis.

I am greatly indebted to my brother, Flight Lieut. Simon Yudkin, for much helpful discussion and criticism during the preparation of this paper. To Dr. Leslie Harris I owe the encouragement and stimulus which aroused and maintained my interest in nutritional problems.

REFERENCES

- Abbasy, M. A., Hill, N. G., and Harris, L. J. (1936). *Lancet*, **2**, 1413.
 — Harris, L. J., and Hill, N. G. (1937). *Ibid.*, **2**, 177.
 Banerji, G. G., and Harris, L. J. (1939). *Biochem. J.*, **33**, 1346.
 Bessey, O. A., and Wolbach, S. B. (1939). *J. exp. Med.*, **69**, 1.
 Bicknell, F. (1940). *Lancet*, **1**, 10.
 Bigwood, E. J. (1942). *Proc. Nutr. Soc.* (in the press).
 Brees, B. B., and McCoord, A. B. (1940). *J. Pediatr.*, **16**, 139.
British Medical Journal (1942). **1**, 77.
 Brunsting, L. A. (1938). *Proc. Mayo Clin.*, **13**, 280.
 Dam, H., and Schönheyder, F. (1934). *Biochem. J.*, **28**, 1355.
 Evans, W. (1938). *Lancet*, **1**, 308.
 Fox, F. W., Dangerfield, L. F., Gottlieb, S. F., and Jokl, E. (1940). *British Medical Journal*, **2**, 143.
 Harris, L. J. (1942). *Lancet*, **1**, 642.
 — Abbasy, M. A., and Yudkin, J. (1936). *Ibid.*, **1**, 1488.
 — Passmore, R., and Pagel, W. (1937). *Ibid.*, **2**, 183.
 — and Ray, S. (1935). *Ibid.*, **1**, 71.
 Hecht, S., and Mandelbaum, J. (1940). *Amer. J. Physiol.*, **130**, 651.
 — and Schlaer, S. (1938). *J. opt. Soc. America*, **28**, 269.
 Heinemann, M. (1938). *J. clin. Invest.*, **17**, 751.
 — (1941). *Ibid.*, **20**, 39.
 Higgins, C. C. (1935). *J. Amer. med. Ass.*, **104**, 1296.
 King, E. J., *et al.* (1942). *Lancet*, **1**, 207.
 Kodicek, E., and Wang, Y. L. (1941). *Nature*, **148**, 23.
Lancet (1942). **2**, 699.
 Mellanby, E. (1934). *Nutrition and Disease*, Edinburgh.
 Minot, G. R., Strauss, M. B., and Cobb, S. (1933). *New Engl. J. Med.*, **208**, 1244.
 Morgan, A. F. (1941). *Science*, **93**, 261.
 Nicholls, L. (1935). *Indian med. Gaz.*, **70**, 550.
 Nixon, W. C. W. (1942). *Proc. Nutr. Soc.* (in the press).
 Orr, J. B. (1936). *Food, Health and Income*, London.
 Roch, M., and Sciclounoff, F. (1938). *Schweiz. med. Wschr.*, **68**, 1343.
 Rowntree, J. I. (1931). *J. Nutr.*, **3**, 345.
 Sebrell, W. H., and Butler, R. E. (1938). *Publ. Hlth. Rep.*, Wash., **53**, 2282.
 Steffens, L. F., Bair, H. L., and Sheard, C. (1939). *Proc. Mayo Clin.*, **14**, 698.
 Strauss, M. B., and McDonald, W. J. (1933). *J. Amer. med. Ass.*, **100**, 1320.
 Sutherland, R. (1934). *British Medical Journal*, **1**, 791.
 Sydenstricker, V. P. (1941). *Ann. intern. Med.*, **14**, 1499.
 — Sebrell, W. H., Cleckley, H. M., and Kruse, H. D. (1940). *J. Amer. med. Ass.*, **114**, 2437.
 Wang, Y. L., and Yudkin, J. (1940). *Biochem. J.*, **34**, 343.
 Warner, E. D., Brinkhous, K. M., and Smith, H. P. (1938a). *Amer. J. med. Sci.*, **196**, 50.
 — (1938b). *Proc. Soc. exp. Biol.*, N.Y., **37**, 628.
 Wolbach, S. B. (1937). *J. Amer. med. Ass.*, **108**, 7.
 Young, T. C. M. (1937–8). *Indian J. med. Res.*, **15**, 453.
 Yudkin, J. (1941). *British Medical Journal*, **2**, 747.
 — (1943). *Lancet*, **1**, 755.
 — and Ferguson, A. (1943). *British Medical Journal*, **1**, 633.
 Yudkin, S. (1941). *Lancet*, **2**, 787.

A. V. Hardy (*Publ. Hlth. Rep.*, Wash., 1943, **58**, 833) used sulphadiazine in the treatment of 19 chronic typhoid carriers, 4 convalescent carriers (including one treated as a case), 21 clinical cases, and 1 clinical relapse, with the following results. Quantitative cultural tests showed that sulphadiazine had a marked bacteriostatic effect on *Eberthella typhosa* in the enteric tract, but that the chronic carrier state was not terminated by this treatment.