compete for recognition as generalizations of the facts of observation, and these "facts" are not always clearly distinguishable from assumptions. Psychology is in a ferment, and we must not plunge the student into the fermenting-vat. The exponents of this field of medicine must first put it in order and lend it coherence before psychological medicine can hope—or ask—to take its proper place in medical education. This is not to say that the present place taken by it in the curriculum is adequate, but to suggest that a wise infusion of medical education by the principles of psychological medicine demands something more fundamental than the compilation of a long list of titles of lectures and appointments to be crammed into an already overloaded course.

Essentials of Reform

Similar considerations must determine and limit the requirements of other special branches of medicine. Each must be subordinated to the balance and moderation we wish to see in a new scheme of medical education. In short, the essentials of reform in medical training may be summarized as follows:

(1) A carefully thought out pruning of the amount of factual knowledge we seek to impart, ruthlessly undertaken in order that

(2) the student may be enabled to develop a more thoughtful and critical attitude, and a deeper understanding of the causes and manifestations of disease.

(3) An explicit recognition of the necessity of a leaven of theoretical medicine, side by side with its vocational aspects, and the provision of a category of teachers who shall develop it.

To these ends every branch of medicine must be prepared to make some sacrifices, so that we shall turn out thoughtful and discriminating graduates, capable of handling the unending flow of new experiences that must come to them in the practice of their profession, and capable also of independent judgment. On this sound stock we can always graft a higher order of knowledge in any desired branch of medicine.

The considerations put forward in this article, and thus summarized, will be familiar to all readers of Sir Thomas Lewis's volume of addresses entitled *Research in Medicine*. In these the fundamental principles of medical education, and the qualities and outlook of the teachers of medicine, theoretical and vocational, are discussed more amply and with more authority than I can command. No one who bears any responsibility for the reform of medical education can rightly ignore the principles laid down and expounded in these addresses.

Conclusion

There are, of course, other aspects of the total problem of medical education not dealt with in this summary expression of a point of view: for example, the type of general education best fitted for the future medical student; the status and categories of teachers we shall need; the better organization of clinical teaching, and the necessity for a drastic reform of our present chaotic and purely vocational examination system. In respect of the last-named, the gross disorganization of the precious last year of the student's clinical training by the quarterly instalments of the Conjoint Finals would defeat any system of teaching.

In short, the problem is a most complex one, and can be solved only when we have formulated the general principles of a reform of the curriculum and of our methods of teaching. The immediate and urgent issue here raised is the necessity for avoiding another patchwork and compromise revision of medical training and the swamping of the student by the addition to his burdens of fresh requirements from new branches of medicine, or from old branches that have taken new names and adopted new slogans. It would be folly to assume that this danger does not exist. The increasing and inevitable specialization in medicine has called into existence sectional interests nearly all striving for what they regard as adequate recognition for themselves, and it is the growth and momentum of these interests that may well threaten the drawing up of a new and rational plan of medical education.

In 1932, in the columns of this *Journal*, Dr. Charles Wilson, as he then was, wrote an article, entitled "The Student in Irons," in which in clear and admirable terms he laid bare

the defects of the content and methods of our teaching of the medical student. Therein he presented, more lucidly than I can hope to do, the aims that should inspire us in any reform. A brief citation from that paper may be allowed.

"It is nowhere denied that we ask impossible tasks of students, who are bewildered by hours of listening, and that we stamp out the habit of reflection by a ceaseless drill. And this is done in the name of education. . . . So strange an anomaly may set us thinking how we are come to such a pass. The blunt truth would appear to be that the handling and revision of the curriculum are based on a misconception of what we can and ought to attempt in a medical school. The one purpose of the student's years is, it seems, not to train and test habits of thought, but to collect and store a set of facts, as squirrels hoard the nuts on which they hibernate. . . . It may be conceded that such a policy was reasonable when medicine was a slowly progressing science that did not greatly alter in a man's lifetime. To-day it leaves him bankrupt when he has scarcely begun. The little he knows is soon antiquated, and he cannot keep up with a rapidly changing science, for we have so contrived to educate him that he cannot educate himself. . . . We trust neither his industry nor his capacity. He must be driven into different pens for every hour of the day, along with a drifting flock that presses now this way and now that. And this, if I may be allowed to change a metaphor, in the hope that he may one day recognize a disease because he has seen it before."

Surely, this analysis goes far more deeply into the problem than do the two reports we have considered, and shows a recognition of the total requirements of a medical education that is absent from them.

If now we are to content ourselves with a so-called reform that attempts no more than to fit still more into the medical course on the pretext that medicine is enlarging its scope, we shall have betrayed the medical student of the future, and shall have failed to do what is now possible to build a science of medicine with its own body of first principles and its own grammar.

PETHIDINE AS AN OBSTETRIC ANALGESIC A REPORT ON 150 CASES

BY

BEDELIA GALLEN, M.B., Ch.B., D.P.H.

Resident Obstetric Medical Officer, St. Mary Abbots Hospital,
London

AND

FREDERICK PRESCOTT, M.Sc., Ph.D., M.R.C.S.

Wellcome Research Institution, London

Pharmacology of Pethidine

Pethidine (also known as dolantin, dolantal, and demerol) is the name given by the General Medical Council to the hydrochloride of the ethyl ester of 1-methyl-4-phenyl-piperidine 4-carboxylic acid, which was synthesized in 1939 by Eisleb and Schaumann in a search for atropine derivatives. The chemical relationship of pethidine to atropine is seen in the following formulae:

Unlike morphine it is a relatively simple compound, although on close inspection of the formula some remote chemical similarity can be discerned between pethidine and morphine; thus both possess a piperidine ring.

The early clinical and pharmacological work on pethidine was done on the Continent, although most of the publications since 1940 are inaccessible owing to the war. It is claimed that clinically it is an outstanding analgesic, midway in action between morphine and codeine. Its analgesic action has been studied by Climenko, Homburger, and Barlow (1943), using the Hardy, Wolff, and Goodell (1940) technique, which consists in measuring the pain threshold of the skin to varying intensities of thermal radiation from a lamp. It was shown that 100 mg. by mouth raised the pain threshold by 50% for as long as six hours. By this method Batterman and Himmelsbach (1943) have shown that 50 mg. of pethidine intramuscularly is twice as potent as 22 mg. of codeine, and that 100 mg. parenterally approximates in analgesic potency to 1/6 gr. of morphine.

It is stated that pethidine is unique in possessing a morphine-like analgesic action, as well as being effective as a spasmolytic and sedative. Its pharmacology has been extensively investigated by Duguid and Heathcote (1940), Schaumann (1940), Gruber *et al.* (1941), and Barlow (1943). The spasmolytic effect has been attributed to a direct depressant action on smooth muscle resembling that of papaverine, and to an atropine-like action on the parasympathetic nerve endings. It is more effective than papaverine in depressing smooth muscle, but less active than atropine. It relieves visceral pain not only by raising the pain threshold but by relaxing smooth muscle. Intubation studies by Batterman (1943) have demonstrated the spasmolytic effect of pethidine on the human stomach, pylorus, and small and large intestines. Animal experiments have yielded conflicting results, depending on the species used, on the technique employed, and on whether the organ was isolated or *in situ*. The spasmolytic effect of pethidine on the human ureter has been demonstrated by Climenko and Berg (1943).

The sedative effect of pethidine, which is more pronounced when given parenterally, is definite but not as marked as that of morphine. Rovenstein and Batterman (1943), who have used pethidine as a post-operative sedative, consider that the effect of 100 mg. hypodermically is equal to that produced by 1/4 gr. of morphine. After large doses sleep and drowsiness may occur, particularly after parenteral administration, but this is of short duration and less than the period of analgesia, and the patient can be easily aroused (Batterman, 1943). The effect wears off in two hours, although if the drug is given at night or to relieve pain sleep may be prolonged. Unlike morphine, pethidine does not produce a state of amnesia.

Effect on the Uterus.—Studies on the effect of pethidine on uterine muscle are conflicting. According to Dreyer (1943) it relaxes the intact and excised uterus; but this is denied by Gruber *et al.* (1941), who claim that in the dog it stimulates the intact and excised organ. This opinion is shared by Yonkman *et al.* (1943). Solomons and Widdess (1943) also state that it stimulates the guinea-pig and rabbit uterus *in vitro*. Using the balloon technique of Chassar Moir, they report that pethidine causes a slight increase in the tone and frequency of contraction of the human uterus *in vivo*. Care must be taken in interpreting clinical results with the balloon technique, as the introduction of a foreign body such as a balloon into the uterine cavity may produce reflex contractions. Woodbury (1943), using this technique, found that pethidine had no effect on uterine contractions in the human pregnant uterus. The effect of any drug on the uterus, animal or human, depends on the blood supply and stage of proliferation of the organ, which in turn depend on the oestrogen and progesterone tide. Thorp (personal communication) states that pethidine enhances the contraction of the rabbit uterus treated with pituitary extract in the oestrogenic phase, but relaxes it after contraction with adrenaline; in the progesterone phase pethidine has no effect. According to Thorp pethidine has no effect on the muscle tone of the uterus of the pregnant cat. In the following study it was evident that pethidine exerted a definite antispasmodic action on the cervix during labour.

Clinical Studies in Labour

The study was made because the unique properties of pethidine that have been reported—its combined analgesic, antispasmodic, and sedative actions—suggested that it might be a

very useful drug for producing obstetric analgesia. Also the fact that it is less of a respiratory depressant than morphine (Batterman, 1943) should render it safer for the baby than the latter, the dangers of which are well known. The reported absence of mental depression, urinary retention, and constipation following the administration of pethidine (Batterman, 1943) also appears to be a definite advantage, as these symptoms are commonly produced by morphine. Lastly, the relatively high margin of safety compared with morphine permits repeated dosage. The possible future shortage of morphine also makes the clinical evaluation of the newer analgesics imperative.

Since this study was begun two publications on pethidine as an obstetric analgesic have appeared in America, where the drug is known as demerol. Gilbert and Dixon (1943) report that in doses of 100 mg. to 400 mg. it noticeably shortens the time of labour in primiparae. The average dose was 300 mg. Adequate analgesia without amnesia was obtained by 72% of the patients; in combination with a barbiturate satisfactory amnesia was experienced by 61%. No adverse effects were observed on mother or baby. Roby and Schumann (1943) gave pethidine intramuscularly in doses of 100 mg. in combination with scopolamine to women in labour as soon as pains began. Satisfactory amnesia was produced in 85% of the cases; 14% of the babies needed some form of resuscitation.

Selection of Patients; Administration; Dosage

Clinical Material.—The series treated with pethidine reported here consisted of 150 cases. Detailed records were kept on the first 100 cases only. Of these, 53% were placid and not apprehensive; 47% were nervous and apprehensive. One of the latter group had a spastic cervix which dilated satisfactorily under the influence of pethidine, but required forceps delivery. Another had a rigid cervix, which failed to dilate and necessitated lower-segment Caesarean section. The 100 cases recorded comprised 70 primigravidae and 30 multigravidae, and all were apparently normal on medical and obstetric examination.

Method of Administration.—Oral administration, which was tried at first, was found to be unsatisfactory on account of the frequency of vomiting and the slowness of action. Roby and Schumann also found the oral route unsatisfactory. It was observed that it took from 20 to 60 minutes for the maximum analgesic effect to be produced by mouth. Pethidine was therefore given either intramuscularly or intravenously, or by the combined routes. Of the 70 primigravidae, 33 received the drug solely by intramuscular injection, while 37 were given both an intramuscular and an intravenous injection. Of the 30 multigravidae, 20 were given the drug intramuscularly, and 10 received it intravenously with or without an accompanying intramuscular dose.

Dosage and Times of Administration.—Two dosage schemes were tried:

(a) A total dose of 200 mg. of pethidine was given intramuscularly either alone or in combination with the stock mixture used in the labour ward (see Table I) or else with scopolamine gr. 1/150. The pethidine was given in two doses of 100 mg., one an hour after the other. The second injection was given either alone or with the stock mixture or scopolamine. In cases in which the first stage of labour was prolonged for more than 8 to 10 hours after the first injection of pethidine further injections were given, up to a total of 400 mg. in 24 hours. Gilbert and Dixon (1943) gave up to a maximum of 650 mg. in the same period. It was exceptional in our series to give more than 200 mg. The average time taken for the production of analgesia after intramuscular administration was 15 minutes, and the duration of action 3 to 4 hours.

(b) In order to obtain a more rapid effect some cases received pethidine intravenously. A dose of 100 mg. was given initially. This acted in 5 to 10 minutes. An hour later another 100 mg. was given intramuscularly, either alone or in combination with scopolamine gr. 1/150. The duration of action was 3 to 4 hours. If the first stage of labour was prolonged for more than 8 to 10 hours after the initial dose further injections of 100 mg. were given intramuscularly up to a total of 400 mg. in 24 hours.

Vomiting and other unpleasant side-effects may occur if pethidine is given rapidly by the intravenous route. This was minimized by asking the patient to close her eyes and by taking five to ten minutes over the injection. Ideally, the first injection of pethidine is best given when the cervix is

two fingers dilated and the patient is having regular contractions. Although pethidine does not depress respiration to such an extent as morphine, it would appear that it is not wise to give it within 2½ hours of the expected delivery of the child. This figure compares favourably with that of four hours for morphine. Many of the patients receiving pethidine actually slept during labour.

Evaluation of Analgesic Effect

The degree of analgesia produced in the first 100 cases compared with controls receiving the stock mixture used in the labour ward (chloral hydrate gr. 30, potassium bromide gr. 30, tinct. opii m. 10) was as follows:

Degree of Analgesia	Pethidine Cases	Control Cases
Complete	10	0
Satisfactory	50	10
Fair	34	90
Nil	5	0
Amnesia	1	0
	100	100

These figures refer to the first stage of labour only, as in most cases Minnitt's gas-and-oxygen apparatus was used to complete the second stage of labour.

In only one case in this series—a young primigravida—was amnesia produced. Initially she was given morphine gr. 1/6 and scopolamine gr. 1/150, followed by 100 mg. of pethidine intramuscularly, which was repeated twice, and a further dose of scopolamine gr. 1/150. Intermittent chloroform anaesthesia was administered at the end of the second stage. It is of interest to record that the duration of labour from the time of administration of the morphine-scopolamine was 5½ hours. Roby and Schumann (1943) obtained satisfactory amnesia in 85% of their cases. A dose of 100 mg. of pethidine was given intramuscularly, followed by scopolamine 1/100 gr. A further injection of scopolamine gr. 1/200 was given an hour later, and in the case of primiparae repeated every one to five hours.

Effect on Duration of Labour

Table I gives the average duration of labour, in hours, of the various groups, the interval between the initial dose of pethidine and the birth of the child, the interval between the final dose of pethidine and the birth of the child, and the average duration of the third stage in minutes.

TABLE I

Medication	Parity †	No. of Patients	Route		Average Duration of Labour (hours)	Average Time between Initial Dose and Birth (hours)	Average Time between Final Dose and Birth (hours)	Average Duration of Third Stage (min.)
			i.m.	i.v.				
Pethidine with or without stock mixture*	P	59	26	33	20-70	7-21	5-34	20
	M	23	15	8	16-47	5-09	3-18	20
Pethidine and scopolamine	M	11	7	4	15-95	9-66	6-76	14-5
	P	7	5	2	16-89	5-90	5-30	15-5
Patients receiving stock mixture only	P	70			18-81			18-9
	M	30			10-74			17-8

* The stock mixture contained chloral hydrate gr. 30, pot. bromide gr. 30, tinct. opii m. 10, aq. chloroformi ad 1 oz.

† P=Primiparae; M=Multiparae.

On examining this table it would appear that labour was prolonged in those patients receiving pethidine, in comparison with controls not receiving the drug. This apparent prolongation of labour was in our opinion due to the high proportion (47%) of apprehensive and nervous women in the selected cases. The actual duration of labour, however, from the time of administration of the pethidine in both primigravidae and multigravidae falls well within, if not below, average time limits. Gilbert and Dixon (1943) report that in their series pethidine noticeably shortened the duration of labour in primiparae, and Roby and Schumann (1943) state that the drug does not prolong labour.

Instrumental Delivery

The use of pethidine did not appear to increase the incidence of instrumental delivery. In this series forceps were used in 10% of all cases, but there was always a good obstetric reason

for this. In only one case was there weakening of uterine contractions necessitating the application of forceps. The incidence of instrumental delivery in the control group not receiving pethidine was 13%. The reasons for the use of forceps in the two groups were as follows:

Reason for Forceps	Pethidine Cases	Control Cases
Contracted outlet	5	2
Deep transverse arrest	1	4
Extreme nervousness and rigid perineum	1	0
Extreme nervousness and lack of co-operation	2	2
Weak uterine contractions and rigid perineum	1	1
Maternal distress	0	2
Foetal distress	0	1
Threatened uterine inertia	0	1
	10	13

Obstetric Complications

The use of pethidine did not increase the incidence of obstetric complications, of which there were three. Two were cases of retained placenta and one a rigid cervix. In one instrumental delivery the placenta was retained, but was successfully expressed by Credé's method. In no other case in the series was there any weakening of the contractions, and hence there has been no tendency to post-partum haemorrhage. On the contrary, it was observed that in a case in which pethidine had been used placenta and membranes were much more liable to come away complete from the uterus. In the second case of retained placenta the latter was adherent to the uterus, necessitating manual removal, but in this instance there was a history of a retained placenta in a previous delivery. In the case of the rigid cervix, in spite of a labour prolonged for 48 hours, and the use of morphine, scopolamine, and pethidine, the cervix failed to dilate, and lower-segment Caesarean section had to be performed in the interest of mother and child.

It would appear that pethidine exerts a definite antispasmodic effect on the cervix, particularly in the clinical entity spasmodic cervix, in the treatment of which it is superior to morphine.

In the control group obstetric complications occurred in 2%. Both were cases of post-partum haemorrhage from a relaxed uterus due to weak contractions.

Reactions in the Mother

The commonest toxic manifestation in the mother was vomiting, which occurred in 28% of the cases. As previously stated, oral administration was discontinued because of vomiting and delayed action. It was also observed that a transitory rise in blood pressure of some 10 to 20 mm. systolic and/or diastolic was fairly common within 15 minutes of administration. This rise of blood pressure did not last for more than 30 to 60 minutes. No association between vomiting and rise of blood pressure was established. On account of the possible rise in blood pressure pethidine was not given to cases complicated by toxæmia or hypertension. Other untoward symptoms often observed, particularly after intravenous injection, were dizziness, tingling of the limbs, vertigo, and dryness of the throat. In no case was excitement observed. The frequency of vomiting following intravenous injection was effectively reduced by injecting the pethidine very slowly over a period of five to ten minutes, while requesting the patient to keep her eyes closed.

Condition of Babies

Of the 100 babies born to mothers receiving pethidine 91 were apparently normal and active at birth, and 9 were slow—i.e., required resuscitation measures. These figures are similar to those of Roby and Schumann (1943). There were two deaths, due respectively to a tentorial tear and a prolapsed cord. The neonatal morbidity for the first three days after birth was nil. Seven of the babies were cyanosed and 7 suffered from respiratory depression. Slight impairment of pulsation was noted in 9 babies and poor muscle tone in 8. In no case did any of the babies fail to respond to routine methods of resuscitation for the newborn, crying well within at most 20 to 30 minutes of birth and responding to stimulation.*

* In the control group given the chloral-bromide-opium mixture there were 2 slow babies. One was limp, but responded quickly to treatment. The other was blue, and went into white asphyxia; it recovered with treatment.

In respect of the 9 slow babies Table II shows the dose of pethidine and the time between its administration and the birth of the child.

TABLE II

Parity	Degree of Dilatation of Cervix on Initial Dose	Initial Dose	Time of Initial Dose before Birth	Final Dose	Time of Final Dose before Birth
1	4 fingers	100 mg. i.m.	2 hrs. 10 min.	100 mg. i.m.	30 min.
1	3 "	100 mg. i.m.	55 min.	—	—
1	3 "	100 mg. i.m.	2 hrs. 25 min.	100 mg. i.m.	20 min.
1	Anterior lip	100 mg. i.v.	20 min.	—	—
0	"	100 mg. i.v.	2 hrs.	—	—
0	3 "fingers"	100 mg. i.v.	2 hrs. 35 min.	100 mg. i.m.	1 hr. 5 min.
0	4 "	100 mg. i.v.	2 hrs. 35 min.	100 mg. i.m.	1 hr. 10 min.
0	3 "	100 mg. i.m.	2 hrs. 25 min.	100 mg. i.m.	1 hr. 25 min.
0	4 "	100 mg. i.m.	3 hrs. 15 min.	100 mg. i.m.	1 hr. 15 min.
		+ scopolamine gr. 1/150		+ scopolamine gr. 1/150	

In all these cases the progress of labour was much quicker than was anticipated clinically. From experience with them it was decided that it is undesirable to administer pethidine within 2½ hours of the expected delivery of the child. The average time between the final dose and the end of labour in Gilbert and Dixon's cases was 2 hours 22 minutes in primiparae.

Summary

In the dosage employed pethidine is an effective obstetric analgesic. It does not, however, produce amnesia unless given with adequate doses of other drugs—e.g., scopolamine and barbiturates.

Only 5% of the patients failed to obtain any relief from pethidine. Analgesia was complete or satisfactory in 60%.

Two dosage schemes were found to give satisfactory results: (a) Initial dose 100 mg. intramuscularly, repeated one hour later with a chloral-bromide-opium mixture or with scopolamine; (b) 100 mg. pethidine intravenously, followed by 100 mg. intramuscularly an hour later, either alone or with scopolamine gr. 1/150. In either technique further intramuscular injections of 100 mg. can be given up to a total of 400 mg. in 24 hours. The first injection is best given when the cervix is two fingers dilated and the patient is having regular contractions.

Pethidine and scopolamine form an effective combination for producing obstetric analgesia.

Given intravenously pethidine produces analgesia in 5 to 10 minutes; intramuscularly in 15 minutes. The duration of action is 3 to 4 hours. Intravenous administration produces a more rapid and powerful initial action, but intramuscular administration is highly satisfactory and equally effective. Intravenous administration is contraindicated in labour complicated by toxæmia or hypertension.

Pethidine has a definite antispasmodic action on the cervix.

Statistically, in comparison with controls pethidine appeared to prolong labour. This was probably due to the selection of the cases. The figures compare well with average figures for the duration of labour.

No increase in the incidence of instrumental delivery or obstetric complications was associated with the use of pethidine.

Reactions in the mother included vomiting, a temporary rise of blood pressure, dizziness, tingling of the limbs, and dryness of the throat. These were transient, and, although unpleasant, gave no cause for alarm.

Of the babies 91% were apparently normal and active at birth; 9% were slow, and required resuscitation.

There were no deaths among the babies that could be attributed to pethidine. The neonatal (three days) morbidity was nil.

This investigation was carried out in the obstetric unit of St. Mary Abbots (L.C.C.) Hospital, Kensington, from Jan., 1943, onwards. Thanks are due to the consultant obstetrician, Mr. W. C. W. Nixon, for his helpful criticism of the investigation and the manuscript.

REFERENCES

- Barlow, O. W. (1943). To be published. Quoted by Batterman, loc. cit.
 Batterman, R. C. (1943). *Arch. intern. Med.*, **71**, 345.
 — and Himmelsbach, C. K. (1943). *J. Amer. med. Ass.*, **122**, 222.
 Climenko, D. R., and Berg, H. (1943). *J. Urol.*, **49**, 255.
 — Homburger, E., and Barlow, O. W. (1943). To be published. Quoted by Batterman, loc. cit.
 Dreyer, N. (1943). Quoted by Batterman and Himmelsbach, loc. cit.
 Duguid, A. M. E., and Heathcote, R. St. A. (1940). *Quart. J. Pharm. and Pharmacol.*, **13**, 318.
 Eisleb, O., and Schaumann, O. (1939). *Dtsch. med. Wschr.*, **65**, 967.
 Gilbert, G., and Dixon, A. B. (1943). *Amer. J. Obstet. Gynec.*, **45**, 320.
 Gruber, C. M., Hart, E. R., and Gruber, C. M., jun. (1941). *J. Pharmacol.*, **73**, 319.
 Hardy, J. D., Wolff, H. G., and Goodell, H. (1940). *J. clin. Invest.*, **19**, 649.
 Roby, C., and Schumann, W. R. (1943). *Amer. J. Obstet. Gynec.*, **45**, 318.
 Rovenstein, E. A., and Batterman, R. C. (1943). *Anesthesiology*, **4**, 126.
 Schaumann, O. (1940). *Arch. exp. Path. Pharmacol.*, **196**, 109.
 Solomons, E., and Widdess, J. D. H. (1943). *Lancet*, **1**, 617; *Irish J. med. Sci.*, **206**, 637.
 Woodbury (1943). Quoted by Batterman and Himmelsbach, loc. cit.
 Yonkman, F. F., et al. (1943). *Proc. cent. Soc. clin. Res.*, **15**, 89.

OBSTETRIC ANALGESIA WITH PETHIDINE

BY

WALTER SPITZER, M.D. Prague, M.R.C.S., L.R.C.P.

Senior Resident Obstetrician, Kingston County Hospital, Surrey

Pethidine hydrochloride has been studied in our maternity department since Oct., 1941, in regard to its analgesic and spasmolytic effects on normal labour. It is a pain reliever of fairly recent introduction, the hydrochloride of the ethyl ester of 1-methyl-4-phenylpiperidine-4-carboxylic acid, discovered by Eisleb and Schaumann in their search for an ideal analgesic with spasmolytic atropine-like properties. The drug is known on the Continent as "dolantin" and in the United States as "demerol," and was officially named pethidine by the G.M.C. a year ago.

The pharmacological virtues and faults of pethidine have been carefully surveyed by Rothschild (1941) of Basle. It possesses a powerful analgesic action like that of morphine; but is stimulating rather than depressing to the central nervous system. Because of its depressant effect on smooth muscle tissue its spasmolytic properties have been compared to those of atropine and papaverine. Several authors have claimed that it can antagonize the effects of acetylcholine and histamine on the intestine (Eisleb and Schaumann, 1939; Duguid and Heathcote, 1940; Schaumann, 1940). Gruber *et al.* (1941) could not acknowledge the spasmolytic action. It is relatively non-toxic, and has little effect on the pulse rate or the pupil. It lowers the blood pressure and depresses the respiration only in large doses.

Only a few clinical reports on pethidine are available in this country. Prof. Christie (1943) undertook an investigation on behalf of the War Wounds Committee of the Medical Research Council. Pethidine proved to be of value as an analgesic when given by the mouth to 335 patients of six different hospitals. Of the patients with "severe pain" 55% obtained complete relief and 20% partial relief; of those with "moderate pain" 61% were completely and 12% partially relieved. Toxic symptoms were uncommon and never serious. Fitzgerald and McArdle (1943) described the usefulness of pethidine in neurological cases.

The Continental literature on pethidine in a variety of conditions is very encouraging; and as regards obstetrics, Benthin (1940, 1942), Fuchs (1941), and Sonnek (1941) have described excellent results in the relief of pain in labour. Detailed literature is not accessible in this country.

It seemed to me to be advisable to investigate another method of inducing analgesia in the first stage of labour which would cover three important points—namely, (1) reduce pain without having a depressing effect on the respiration and circulation of mother or baby; (2) avoid psychic depression; (3) avoid prolongation of labour. The choice of obstetric analgesics in any particular case has often to be made according to existing conditions. That is why an ideal method has not yet been agreed upon and legions of analgesic and anaesthetic substances in a variety of methods of administration are used all over the world. The exhaustive studies of obstetricians in this country have shown that "twilight sleep" is, for the baby, a dangerous first-stage treatment. All general anaesthetics weaken uterine contractions and retraction. The barbiturates depress the prothrombin levels in mother and child by causing damage to the liver, in which prothrombin is formed (Browne, 1942; Fitzgerald and Webster, 1942). Barbiturates are sometimes responsible for delay, with deleterious effects on the third stage of labour.

In domestic practice chloral hydrate and bromide and/or morphine are still most commonly used. While the first two drugs are safe and fairly satisfactory, the use of morphine for analgesia is unsafe for both mother and baby. It is encouraging that Eardley Holland (1943) has reminded the profession again of the dangerous effects of morphine as regards the respiration of the foetus. Especially in the present era, in which there is an increase in the number of premature labours, the liability to primary apnoea and atelectasis or pneumonia after use of morphine or the popular derivatives has to be taken into account. In short, all narcotics depress the respiratory centre of the foetus far more than that of the mother, and occasionally