LUNG ABSCESS IN RELATION TO THE INFLUENZA EPIDEMIC

BY

ELI DAVIS, M.D., M.R.C.P., M.Sc.

Deputy Medical Superintendent and Physician, St. Andrew's Hospital (L.C.C.)

In the fourth week of Nov., 1943, the influenza wave impacted upon St. Andrew's Hospital. Some forty days later influenza admissions dropped abruptly. During this period 5 cases of lung abscess (3 fatal) were seen among 17 patients aged 12 to 60 with influenzal pneumonia. They are described below. Of the 12 other patients, 2 died, and one of the latter cases (on which a necropsy was done) is also described. Results of sputum cultures and white cell counts are given in the Table.

Summary of 6 Cases of Influenzal Pneumonia, 5 with Lung Abscess

Case No. and Outcome	Bacteria in Sputum	Site of Abscess
I: Recovery	Staph. aureus (pure culture)	Right upper lobe
II: Death	Staph. aureus; "H. influenzae"; haem. strep.	Left "lower" lobe; left eye
III: Recovery	Staph. aureus; H. influenzae; pneumococcus	Right upper lobe
IV: Death	Staph. aureus; H. influenzae; pneumococcus	Right middle lobe; right brain
V: Death	Staph. aureus; pneumococcus	Right middle lobe; right brain
VI: Death	Haem. strep.; H. influenzae	None

At the height of the disease the white cell counts ranged from 11,000 to 18,300 per c.mm., with polymorphonuclears around 80%.

Case I

A housewife aged 38 was admitted with pneumonia of the right upper lobe, following influenza two weeks earlier. There was no response to full doses of sulphathiazole, and within 48 hours of admission radiological evidence of abscess of the right upper lobe was seen. She coughed up 2 oz. of slightly blood-stained purulent sputum daily, mostly after postural drainage. After four weeks of acute illness in hospital, with a temperature swinging between 99° and 102° and a pulse rate of 100 to 128 a minute, she began to recover steadily. Her serum contained no antibody for psittacosis antigen (Prof. S. P. Bedson) and no cold agglutinins.

Case II

The illness of this patient, a labourer aged 40, began with dizziness and collapse five days before admission. There followed pain in the chest, cough, and scanty sputum but no haemoptysis. When seen he was panting and cyanosed; temperature 101.4°, pulse rate 92, respirations 32 a minute. He was regarded as a case of influenzal bronchopneumonia, with the right lung more involved than the left. There was heavy albuminuria. He died in 48 hours, having had 18 g. of sulphapyridine, oxygen, and nikethamide. At necropsy the right lung showed confluent bronchopneumonia, with numerous abscesses 0.5 cm. in diameter in the anterior half of the upper lobe, collapse at the base of the lower lobe, and purulent bronchitis. There was right serofibrinous pleurisy, with 15 oz. of pleural fluid. The left lung showed oedema and congestion only.

Case III

In this case, that of a lorry driver aged 33, influenza began one week before admission and was followed by rigors, cough, and scanty rusty sputum. On admission he had left basal crepitations. Temperature 100.8°, pulse 112, respirations 28. Severe left uveitis developed in two days, and blisters on his fingers 24 hours later. The left eye became more and more inflamed until there was a panophthalmitis. Sulphapyridine was given in full doses from admission and sulphacetamide was applied to the eye, but without benefit. The left eye was removed on the 12th day of the disease by Mr. M. Shaw. A radiograph on the 15th day showed an abscess of the left lower lobe. Temperature and pulse settled on the 24th day, and there was steady improvement thereafter, with natural healing of the abscess.

Case IV

A labourer aged 57 had suffered from chronic bronchitis for years. He collapsed one week before admission, and cough with yellow sputum developed. When admitted he showed marked cyanosis and dyspnoea, general bronchitis and right bronchopneumonia; temperature 99.4°, pulse 132, respirations 32. He seemed to respond

to sulphapyridine, the fever subsiding quickly and the temperature remaining down while on sulphapyridine (25 g. given) and for a further 60 hours. The temperature then rose again and ranged from 98° to 103°. On the 23rd day of the disease the sputum became bloody and treacly, and abscess of the right upper lobe was diagnosed and confirmed by x rays. He coughed up 14 oz. of sputum daily, mostly in the morning, after postural drainage. His temperature settled (spontaneously) on the 43rd day of the disease, but his general condition deteriorated, and he died afebrile on the 46th day. Necropsy was refused.

Case V

A female munition worker aged 20 was taken ill 5 days before admission with a rigor, tightness behind the lower sternum, and slight cough with scanty sputum. On examination she presented a toxic aspect, and had a few crepitations over the right anterior axilla; temperature 103°, pulse 120, respirations 24. The temperature dropped abruptly after 6 g. of sulphapyridine, but rose again, gradually at first, despite continuing with the drug. By the time 25 g. was given the temperature had reached 104°. Sulphapyridine was stopped. Fever continued swinging from 100° to 104° for the next ten days. Rales became numerous at the right base. The patient insisted that she "felt fine." A drachm or so of nummular sputum was coughed up daily, and was haemorrhagic only once. On the 21st day of the disease tinkling rales were heard over the right mid-zone, and lung abscess was considered. Rales were also heard over the left base. Sulphamezathine therapy was begun; but by the 23rd day deterioration was pronounced, with drowsiness, contracted pupils, and bilateral papilloedema. The cerebrospinal fluid was turbid with masses of polymorphonuclears, but no organisms were grown from it. Sodium sulphadiazine (5 g.) was given intravenously, and was repeated next morning, in addition to oral sulphamezathine. She died on the 25th day of the disease.

Chest Radiographs.—On the 9th day of the disease x-ray films of the lungs showed no abnormality; on the 15th day there was a right hilar shadow (later proved to be due to lung abscess), with surrounding pneumonia. Blood culture was negative.

Necropsy.—There was muco-purulent bronchitis. The right middle lobe showed grey consolidation. Anterior to this grey area was a superficial acute abscess 5 by 4 by 4 cm., extending nearly to the midline behind the sternum. There was a spread of pus to the mediastinum above the base of the heart, with two small secondary abscesses here, near enlarged lymphatic glands. Intense congestion, collapse, and recent bronchopneumonia were observed in the right and left lower lobes, and recent fibrinous pericarditis. The spleen was moderately enlarged, congested, and soft. In the brain were two superficial acute abscesses, 2.5 cm. in diameter, containing green pus—one in the right temporo-parietal region, and one in the right occipital region, with purulent basal meningitis. The middle-ears were clean. The pus from the brain abscess grew a pure culture of haemolytic streptococci, although the organisms from sputum culture were *Staph. aureus* and pneumococci.

Case VI

A housewife aged 56 gave a history of two attacks of influenza in the previous month. The second attack began two weeks before admission and was followed by cough and chest pain. On examination there were signs of left basal consolidation and right bronchopneumonia. The temperature (99.6°) soon subsided with sulphapyridine therapy, and by the time 22 g. had been given had been normal 48 hours. The drug was stopped, but the temperature rose again, and continued to swing between 99° and 101°. She coughed up 1 oz. of blood-stained sputum daily. The lung signs persisted, and she deteriorated, dying on the 19th day in hospital. She had been given sulphamezathine for the last 36 hours of life.

Necropsy.—There was a diffuse, coarsely granular consolidation of all lobes of both lungs and dense diffuse fibrosis of the bases. The appearances suggested unresolved pneumonia on a previously fibrotic lung, and this was confirmed histologically. There were recent bilateral fibrous pleural adhesions. The spleen was twice the normal size, oedematous, congested, and soft.

Discussion

Influenzal pneumonia was diagnosed on clinical grounds. The disease began in most of the patients with the features characteristic of the influenza prevalent in the neighbourhood—fever, malaise, generalized aches and pains, and cough with scanty sputum. When pneumonia developed the temperature was not initially as high as in lobar pneumonia, but fever was more prolonged. Physical signs were not clear-cut. White cell counts were essentially the same as in primary lobar pneumonia. The sulphonamides had no obvious effect in at least 12 of the 17 cases. The remaining 5 cases cleared up well concurrently with the administration of sulphonamides.

It is conceivable that complications and mortality would have been greater but for the sulphonamides, but I do not think so.

The 17 patients considered in this paper were aged 12 to 60. Pneumonia in infants did not differ essentially from the cases I saw in the previous year. Many chronic bronchitis over the age of 60 who would almost certainly have died during the winter from bronchopneumonia and congestive heart failure had death precipitated by influenza. Of the 17 cases, 12 recovered. Recurrent epistaxes were prominent in one boy, another patient had haemorrhagic vesicles of the inner cheeks, and one had a small pleural effusion. Pneumonia was bilateral in 4 patients. Before the epidemic proper began, 2 cases of influenzal pneumonia were seen—a fatal case in August on which a necropsy was done, and one in mid-October which recovered.

It is of interest to compare this experience of influenzal pneumonia with 5 deaths and 5 cases of lung abscess (and *Staph. aureus* the chief organism concerned) with recent experience of lobar pneumonia. From Oct. 1, 1942, to Dec. 31, 1943, I saw 107 consecutive cases of primary lobar pneumonia, the ages ranging from 5 to 65. There were 2 deaths, and 2 cases of lung abscess with recovery. From Jan. to Oct., 1943, I saw 13 cases of primary atypical pneumonia (one with rising titre of psittacosis antibody—Prof. Bedson), but none since. It is tempting to inquire if there is any relation between the recent frequency of primary atypical pneumonia and the influenza epidemic.

Concurrently with the influenza epidemic, primary lobar pneumonia became much more frequent here; 26 patients aged 5 to 65 were seen in 40 days, with one death. It may be that a number of these cases were really influenzal pneumonias, but there was not enough clinical evidence for so regarding them.

Summary

In the recent influenza epidemic, among 17 cases of influenzal pneumonia—ages 12 to 60—there were 5 deaths. There were 5 cases of lung abscess (3 fatal), and one patient who recovered had a metastatic panophthalmitis. *Staph. aureus* was an important invader. Sulphonamides were not beneficial. In 107 cases of primary lobar pneumonia—aged 5 to 65—there were 2 deaths, and 2 cases of lung abscess with recovery.

I thank Mr. J. R. M. Whigham for his encouragement; and Dr. C. Hilliard, radiologist; Drs. A. B. Rosher, A. B. Bratton, and C. C. Bryson, pathologists; and Drs. R. Gross-Heisler and S. Gee of the medical team, for their help.

PROGNOSTIC VALUE OF LABORATORY INVESTIGATIONS IN TYPHOID FEVER

BY

S. S. BHATNAGAR, M.D., Ph.D., F.R.C.P.

Lieut.-Colonel, Indian Army Medical Corps

(From a Combined Indian General Hospital, India Command)

Laboratory investigations in typhoid fever are usually employed for the confirmation of clinical diagnosis, either directly by the isolation of the infecting organism from blood, urine, or faeces, or indirectly by the estimation of antibody production, chiefly Vi and O agglutinins, as proof presumptive of the presence of *Bact. typhosum* in the human body. In military hospitals in India blood cultures in the early stages, agglutinations throughout the course of infection, and collection of evidence for the excretion of micro-organisms, even during convalescence, comprise the routine procedure for such cases. The analysis of collective results so obtained, in conjunction with clinical observations, indicated that the various combinations of signs and symptoms met with in this disease were associated with distinct variations in laboratory findings, the study of which was of considerable prognostic significance. Not only was it thus possible to gain an insight into the ups and downs of the struggle between host and parasite, but specific antityphoid serum therapy, practised in a large number of cases (to be presented in a subsequent communication), could also be placed on a more satisfactory basis.

The object of this paper is to describe and discuss the prognostic inferences drawn from a study of 280 cases of typhoid fever in which laboratory investigations were carried out at short intervals throughout the course of infection. Reference is made to certain aspects of the diagnosis and prophylaxis of typhoid that are directly related to the observations recorded here. For reasons of space tables and charts are not included.

The whole group of cases has been subdivided into six different clinical and bacteriological identities. Although some overlapping of symptoms, rather than of signs, was noticeable between the various subgroups, it was not extensive enough to confuse the main issues.

Clinical and Bacteriological Types of Typhoid Cases

1. *Simple Type*.—These cases presented the textbook description of the disease, with temperature which followed the course of three well-known stadia—*incrementi*, *acmes*, and *decrementi*. The total duration of pyrexia varied from 10 to 25 days, depending upon the inoculated state of the individual. In a person who had had multiple injections of T.A.B. vaccine—prepared in a reliable institution with appropriate strains of *Bact. typhosum*—all the stages were completed quickly, and the patient looked so well towards the end of the second week that, in the absence of laboratory diagnosis, doubts could be expressed as to the nature of the infection. Such cases were usually met with among military personnel with more than six years of colour service. In the uninoculated pyrexia was prolonged until the fourth week. In both types of individuals, however, the entire course of the disease was free from complications. Bacteriologically, blood culture was usually negative, since bacteraemia in such cases was of short duration. Agglutination tests, on the other hand, were most satisfactory, and of the type commonly designated as “Widal-positive”; the O titre was about 200 and the Vi agglutinins were as described in a previous communication (Bhatnagar, 1938). It was easy to isolate the infecting organism from urine or faeces, or both, during the second, third, or fourth week. The strain isolated was agglutinable by both O and Vi sera, and its average lethal dose (a.l.d.), for white mice weighing from 20 to 22 g., was more than 100 millions—the V-W type of organism described by Kauffmann (1935).

2. *Ambulatory Type*.—These cases represented the other extreme of the prognostic picture in that they were invariably fatal. Before the patient was obliged to seek admission to a hospital or take to bed properly, he had been definitely ill for over a week. The case below illustrates the various points:

A robust, healthy man, 28 years of age and 14 stone in weight, was feeling seedy for about 10 days before admission to hospital. During this period he took aspirin and quinine regularly to enable him to carry out his duties. He was ultimately obliged to stay in bed. On admission enteric was suspected, and in spite of careful nursing and efficient palliative treatment his condition deteriorated. On the fifth day he complained of pain in the lower abdomen. Clinical examination revealed perforative peritonitis. His general condition was, however, so bad that any surgical intervention was ruled out; he died a few hours later with a rectal temperature of 110° F. Necropsy revealed typical typhoid ulcers in the small intestine.

Laboratory investigations in such cases were most disappointing: they presented a completely negative picture—so much so that the reliability of clinical diagnosis was questioned on more than one occasion. No Vi agglutinins were detected at any stage of the disease, although in some cases the O antibody titre could be considered to be suggestive of typhoid infection. Diagnosis could be confirmed post mortem only by direct culture from inflamed Peyer's patches, spleen, gall-bladder, and bone marrow. In the few cases in which this facility was available it was particularly observed that the growth of *Bact. typhosum* on first isolation was scanty and the strain of only average virulence—a.l.d. more than 100 millions, and the organism agglutinable by both O and Vi sera.

3. *Toxic Type*.—These cases were usually seen either among the civilian population or among military personnel—such as drafts arriving from over-seas—who had not had the benefit of multiple T.A.B. inoculations. The patient was clinically toxic before the end of the first week of illness; the skin