scopic antero-posterior and lateral views and then, if necessary, oblique views. Posterior fractures involving the anterior clinoid process may be elucidated by taking an axial view of the optic foramen where the outer boundary is formed by the anterior clinoid process. Cases 9 and 10 were of the type described as a "block" fracture. The central naso-frontal block of bone was detached by two lateral fissures through the ethmoids in the one case and through the temporal and frontal bones in the other.

Spinal Puncture.—I believe that all patients with a head injury should have a spinal puncture. In the case of these anterior fossa fractures, examination of the spinal fluid may give warning of meningitis, and also it will check the presence of blood if there has been any doubtful nose-bleeding.

Meningitis.—Meningitis itself may confirm the diagnosis. Case 6 showed no fracture in the original films, but the history of five recurrent attacks of pneumococcal meningitis was accepted as confirmation of a dural tear, and a stereoscopic film revealed a hairline crack in the frontal sinus. Case 3 developed mild meningitis—headache, neck rigidity, pyrexia, and 10 cells per c.mm.—on the 25th day after injury. This was controlled in a few days by sulphadiazine therapy. On the 46th day he complained of a "running cold in the head," and was observed to have a profuse C.S.F. rhinorrhoea. It may be mentioned, by the way, that, though frank meningitis is diagnosed clinically by delirium, pyrexia, neck rigidity, and spinal fluid containing several thousand cells per c.mm., I cannot remember a case in which an organism could be recovered from the spinal fluid after sulphadiazine therapy had been instituted.

Treatment

All of these 10 cases were seen some time after their original injury, when they were co-operative and could be investigated fully. There should be no urgency in operating on a recently injured head case until the patient has recovered from the cerebral trauma; and, in fact, judged by this series, the diagnosis of a torn dura would not be made for several days or weeks.

Preliminary Treatment.—All cases received prophylactic sulphadiazine—preferably by mouth, but intravenously if necessary. The dosage was 6 g. daily, but if meningitis was present this was increased to 12 g. daily and an initial 6 g. given intravenously. Blood transfusion was rarely necessary, as the operation does not produce shock and was usually completed in two hours.

Anaesthesia.—This was essentially local infiltration of 0.5% novocain solution after a preliminary hyoscine and morphine injection, which was supplemented by intravenous pentothal drip when necessary. Our procedure was to set up an intravenous saline drip apparatus at the start, and then either blood or pentothal could be rapidly substituted.

Operative Technique.—A coronal incision within the hair line was used in most cases, turning the forehead flap down to expose both supraorbital ridges. In one case only a linear midline scar already present was incised to turn a unilateral flap, but the cosmetic effect of the coronal flap is well worth achieving at the expense of a bigger incision. The bone flap was unilateral or bilateral, dependent on our confidence in the localization of the dural injury. In either case it was hinged on the temporal muscle, and was made with four burr-holes for the unilateral flap and six burr-holes if bilateral. The lower midline burr-hole was placed as low as possible after studying the size of the frontal sinus. Five cases had bilateral bone flaps, and in only one case was there any bleeding from the longitudinal sinus. The dura was stripped up off the orbital plate both medially towards the cribriform plate and posteriorly towards the sphenoidal crest. Any fracture was easily seen, and the dural tear or escape of C.S.F. was looked for on the opposing dural surface. The opposite side was explored similarly. The more difficult case was the fracture of the crista galli or cribriform plate, where, to get at the dural tear adequately, the dura had to be stripped out of the cleft beside the crista galli—a very difficult procedure. The dura could then be stripped back easily to the sella turcica region if necessary. This complete elevation on both sides was necessary in only two cases (Case 6 and Case 10). It allowed the dural holes to be pursestringed with fine silk, and gave a very good exposure, but is best avoided if any smell sensation is still present. Our procedure was then to powder the dura with sulphathiazole, apply a fascial graft over the area like a postage stamp, and seal an edge with diathermy. The crista galli was nibbled away before applying a midline graft, which would be distorted by its presence. Fascia lata was employed in six cases in which a good solid graft was indicated. Temporal muscle fascia was used in three cases, but it is thin and no great area can be covered. A muscle-graft was used in Case 5, where there was a minute and dubious hole. The graft was applied extradurally except in Case 3, in which the dura was opened widely to permit removal of the adherent anterior lobe. In this case it was much easier to apply the graft intradurally. It is reasonable to suppose that some other cases could be grafted more simply by working intradurally and avoiding a wide elevation of the dura. But the dural tears are small, and are best located by finding

the fracture and then seeing the hole or the escaping C.S.F. in that region. I do not think these small dural tears could be located as well intradurally, and also there is an advantage in avoiding trauma from direct handling of the cerebral lobes.

All bleeding was stopped by diathermy to the dural veins and wax to the bone. It is much more important to have the dural bleeding completely stopped in these low-pressure cases than in a tumour operation, in which the raised pressure will stop any extradural oozing. The whole dura was then powdered and the bone-flap and scalp sutured in place in the usual way. Bone chips were employed to fill up the frontal burr-holes.

After-treatment.—No special complication arose. Both eyes were usually closed by frontal oedema for one or two days. Sulphadiazine therapy was continued for five days after operation with the routine dosage of 6 g. daily. These patients had little pain, and some wished to be allowed up on the third day, though this was not permitted until the tenth day. When revisiting the unit on Nov. 12 all patients had been discharged and were understood to be well.

Those patients with complete anosmia did not appear to mind their disability. They could not detect any testing materials in the mouth or nose, though taste was unimpaired. One patient (Case 7) had a constant subjective sensation which he described as a "disinfectant" smell.

Conclusions

The following are the conclusions arrived at: (1) That the dangers associated with C.S.F. rhinorrhoea need stressing to medical officers. (2) That unless the possibility of dural tearing is kept in mind, the rhinorrhoea, anosmia, and radiological evidence of fracture may be easily overlooked. (3) That every head injury with C.S.F. rhinorrhoea, however slight, should undergo operative repair of the dura rather than risk the vagaries of natural repair. (4) That with the technique used the operation is neither formidable nor dangerous; there were no deaths and no complications in this series.

I wish to thank the War Office for permission to publish this paper, and also my colleagues Lieut.-Col. R. R. Lumsden, E.N.T. Specialist, Major W. H. Milligan, Radiologist, Major J. E. Caughey, Neurologist, and Major H. B. Stallard, Ophthalmologist, for their helpful collaboration in the care of these patients.

REFERENCE

Calvert, C. A. (1942). *J. Laryng. Otol.*, 57, 499.

USE OF CRUDE PENICILLIUM FILTRATE FOR LOCAL TREATMENT

BY

J. M. ALSTON, M.B., M.R.C.P.Ed.

(Archway Hospital, London)

The therapeutic value of partially purified penicillin has been established in this country (Florey and Florey, 1943; Fleming, 1943; Report to War Office, 1943; Christie and Garrod, and others, 1944) and in the United States (Keefer *et al.*, 1943), but the use of untreated filtrate of cultures of *Penicillium notatum* has been very little noticed, although successful results have been obtained with gauze impregnated with the living mould (Hobson and Galloway, 1944).

This communication gives the results of the local use of the untreated filtrate in 24 patients. The strain of *P. notatum* was kindly given by Prof. Fleming, and in growing it for this purpose the medium has been changed from time to time, but has always been a modification of the Czapek-Dox formula. At present it is made according to that of Challinor and MacNaughtan (1943) without the yeast extract which they recommend and with the addition of calcium carbonate. The following constituents are dissolved in a litre of distilled water:

Sodium nitrate (NaNO ₃)		3.0	grammes
Potassium chloride (KCl)		0.5	"
Magnesium sulphate (MgSO ₄ , 7H ₂ O)		0.5	"
Ferrous sulphate (FeSO ₄ , 7H ₂ O)		0.01	"
Potassium dihydrogen phosphate (KH ₂ PO ₄)		6.5	"
Disodium hydrogen phosphate (Na ₂ HPO ₄ , 12H ₂ O)		33.5	"
Glucose (pure B.D.H. anhydrous or "Climax" brand, Corn Products Ltd.)		40.0	"
Distilled water to 1,000 c.cm.			

Sterilize in free steam for 1½ hours in 200-c.cm. amounts in 1-litre conical flasks. Add 5 c.cm. of 10% solution of sterile calcium carbonate to every 200 c.cm. of medium before inoculating. The final pH is about 7.

The medium should form a depth of not more than 1.5 to 2 cm. in order to expose to air a large surface relative to the volume. Incubation is at a temperature of 24° C. for 8 to 10 days. When the flasks have not been shaken the mould forms a thick firm pellicle and the culture medium below is almost as clear as before incubation. The fluid medium, after growth, is passed through a Seitz filter, and is tested to make sure that the reaction is nearly neutral and that less than 0.1% glucose is left.

The assay of the product is made by comparing its action on the Oxford strain of staphylococcus with the action of a standard specimen of calcium salt of penicillin. The method mostly used is the established one of measuring the zone of inhibition of the staphylococcus around a piece of porcelain tube (containing the penicillin) set on a heavily inoculated plate of agar medium, and, occasionally for check, the assay is made by inhibition of the growth of the coccus in broth produced by serial dilutions of the standard preparation and the unknown filtrate (M.R.C. War Memorandum No. 12). The filtrates used for treatment varied in strength from 4 to 10 units per c.cm.; they do not lose strength when kept at 4° C. for seven days nor when heated to 65° C. for 2½ hours. More details of the methods used for production and assay are described in a paper by R. M. Sandercott, which is expected to appear in the *Bulletin of the Institute of Medical Laboratory Technology* in July of this year.

Specimens of pus, etc., from patients for whom penicillin treatment was desired have in most cases been cultured bacteriologically, and the most important pathogens (usually staphylococcus or haemolytic streptococcus) tested for sensitivity to penicillin.

The filtrate is applied to superficial lesions by gauze soaked in it, by a cream (made of lanette wax 4%, liquid paraffin 25%, paraffin wax 3%, and filtrate 68%) or by portions of the recently grown mould after the fluid medium had been drained away from it. Cavities in bone which are accessible to the surface are packed lightly with gauze soaked in the filtrate and sealed (if surrounding surfaces are suitable) with sheets of "cellophane" fixed by elastoplast. Deep sinuses in muscle and subcutaneous tissues are treated with indwelling catheters, which are carefully shortened as the sinuses heal from the bottom. Cavities and sinuses are washed out with saline before fresh filtrate is left in them. Treatments are carried out at first twice a day and usually, after a few days, once daily and then less frequently. In dealing with large wounds with sinuses the number of dressings is reduced to a minimum.

In general, when the treatment is successful it produces the same results as solutions of the calcium salt of the drug used locally. These are reduction in suppuration, increase of healthy granulation tissue and epithelium, decrease of pain in the lesion under treatment, and a rapid improvement in general well-being. Patients state that the filtrate stings to a moderate or, with a few batches, a severe degree when it is applied to a raw surface, but as a greater compensation it quickly reduces the tense throbbing pain and the burning sensation of the inflamed tissues from which many of them have suffered almost continuously for months or years. Experience has also confirmed that the drug is not a substitute for surgery but a very powerful ally to it, and in a complicated case repeated surgical measures may be needed to bring the benefits which they themselves are known to produce and to allow the penicillin the direct access to bacteria which it requires. In some cases of acute or chronic osteomyelitis, sequestrectomy may be needed after the filtrate has reduced the toxæmia and the inflammation of the soft tissues, but the benefit to the patient is still undoubted. Brief reports are given of all patients treated so far, with the exception of some very minor infections and some patients in whom treatment has been too recently begun for comment. The names of the surgeons who were in charge of the patients and who kindly supplied the notes are given in parentheses.

Case Histories

Case 1.*—Male, 28 years. Chronic osteomyelitis of sacrum. Multiple sinuses. Injured in Germany, April, 1942. Admitted Feb. 22, 1944, with multiple intercommunicating sinuses. No

material change until instillation of penicillium filtrate started on March 13 and continued daily for 3 weeks and subsequently at shorter intervals. Marked improvement in general condition; diminution of discharge and of haemolytic streptococci. Hip mobility restored. Total penicillin, 1,500 units. (Mr. E. T. Bailey.)

Case 2.*—Male, 25 years. Compound fracture of right femur and shell wound. Wounded Jan. 5, 1943. Admitted to Northern Hospital July 8, 1943. Plaster was applied in Tobruk four months previously. No material improvement after 3 months' sulphonamide treatment, blood transfusions, etc., until on Jan. 14, 1944, penicillium filtrate was instilled once daily for 8 days. Effect was progressive reduction in amount of discharge and of haemolytic streptococci, followed by gradual healing of 10-inch thigh wound and establishment of bony union. Total penicillin, 1,000 units. (Mr. E. T. Bailey.)

Case 3.*—Male, 17 years. Staphylococcal osteomyelitis of tibia and abscess of thigh. Admitted June 30, 1943, for possible tuberculosis of right hip. Developed abscess of thigh Aug. 10 and acute osteomyelitis in upper tibial epiphysis six weeks later. Tibial cavity drained and daily instillation of penicillium filtrate begun Oct. 18, continued for one week, and repeated for another week a fortnight later. Progressive and rapid decrease of discharge and of *Staph. aureus* and haemolytic streptococcus; closure of cavity and restoration of normal x-ray appearance. Effect on abscess of thigh similarly treated: almost immediate response of previously intractable inflammation, with healing in four weeks. Total penicillin, 800 units. (Mr. E. T. Bailey.)

Case 4.—Through-and-through war wound above knee; recent sequestrectomy. 500 units instilled during week resulted in almost total healing in a month, comparing most favourably with cases without penicillin. (Mr. E. T. Bailey.)

Case 5.—Male, 8 years. Subacute osteomyelitis of tibia for 2 months. 300 units during 10 days caused rapid reduction of profuse pus and the granulation of 2 sinuses, with much improvement of general health. The penicillium filtrate treatment was started immediately after second operation. (Dr. D. Kaye.)

Case 6.—Male, 21 years. Profuse suppuration for 3 weeks of finger with compound fracture of phalanx. 100 units during a week produced in that time a clean wound with healthy granulations. (Mr. R. A. V. Lewys-Lloyd.)

Case 7.—Male, elderly. Chronic osteomyelitis of tibia. Two sinuses were treated during 8 to 10 days with 140 units; one was practically healed and one quite healed and dry. In comparison, a third sinus was unchanged while treated with acriflavine in the same period; this sinus is now responding to penicillin. (Mr. R. A. V. Lewys-Lloyd.)

Case 8.—Female, 13 years. Sinus discharging for months following perinephritic abscess and nephrectomy. 500 units during 3 weeks caused cessation of discharge, much reduction in size of sinus, and general improvement in health. (Dr. G. M. Palser.)

Case 9.—Male, adult. Deep infected wound of palm 2½ in. by 1 in., of two months' duration; healing ceased. 30 units and mycelium during a week caused healthy granulation all over and rapid epithelization. (Mr. E. E. Bourke.)

Case 10.—Female, 46 years. Two syphilitic ulcers of leg, stationary for 5 years despite constant treatment. 300 units during a month greatly reduced swelling and pain of leg and reduced ulcers to less than half the area. Treatment is being continued. (Miss G. M. Sandes.)

Case 11.—Female, 45 years. X-ray burn and ulcer of big toe, lasting 6 years, with very slow improvement under treatment. Rapid improvement started by 2 weeks' treatment with calcium penicillin and continued to almost complete healing with 500 units of filtrate given during 5 weeks. (Miss G. M. Sandes.)

Cases 12, 13, and 14.—Three female children with gonococcal vulvovaginitis for 3½, 9, and 14 months, resistant to sulphadiazine (2) or sulphapyridine. 40 units each during 4 days resulted in cure, observed for four months. Filtrate retained in vagina by elevation of pelvis for half an hour after daily instillation. (Miss G. M. Sandes.)

Case 15.—Female, 32 years. Abscess in superficial tissues of abdominal wall associated with widespread surrounding erythema and rheumatic pains, lasting for 6 weeks after appendicectomy. During 12 days 240 units produced cessation of suppuration and progressive healing of wound. (Miss G. M. Sandes.)

Case 16.—Female, adult. Operation wound for frontal sinusitis, indolent for two months, with inflammation of surrounding skin. 100 units followed by healthy granulation of wound and subsidence of skin infection. (Mr. Bedford Russell.)

* Patients 1, 2, and 3 were demonstrated by Mr. Bailey and me at a meeting of the Clinical Section, R.S.M., on April 14, 1944.

Case 17.—Female, 38 years. Abscess and sinuses of abdominal wall for 12 months. 600 units during 7 weeks, with sulphapyridine powder during part of time, almost ended profuse suppuration, closed sinuses and gape of wound, and improved general condition. (Mr. I. Price.)

Case 18.—Female, adult. Large breast abscess with much swelling of surrounding breast; incised and one pint of pus removed. 200 units of penicillium filtrate instilled twice a day during five days reduced discharge to a little sero-pus and brought surrounding tissues to normal. (Dr. G. M. Palser.)

Case 19.—Female, 65 years. Cellulitis and suppuration in resutured abdominal operation wound. Local treatment by sulphacetamide (albicid) and by eusol produced very slow progress during two months; 150 units during 5 days ended suppuration, reduced sinuses and gape of wound, and produced healing edge all round. (Mr. W. T. Kenny.)

Case 20.—Female, 17 years. Two abscesses in right breast. The larger, after incision, treated with 300 units during 3 weeks, healed well in spite of a near-by pocket requiring another incision. This illustrated need for close contact of penicillin with the suppurating lesion, and the result was considered only moderately successful. (Dr. G. M. Palser.)

Case 21.—Male, adult. Osteomyelitis of tibia greatly improved in healthy granulation by 170 units; unopened abscess produced septicaemia, successfully treated by sulphathiazole. Rapid healing of sinuses and abscess then resumed. Septicaemia recurred. Systemic administration of penicillin needed. (Mr. E. T. Bailey.)

Case 22.—Female, 40 years. Acute breast abscess. After incision, 200 units in 9 days, followed by healing not more rapid than normal. (Dr. G. M. Palser.)

Case 23.—Female, 10 months. Subacute suppurative cellulitis of temporal region and brain abscess (probably due to *B. necrophorus*). Profuse suppuration in muscles of temporal and pterygoid regions subsided to oedema after 80 units during one week, but pus had rapidly decreased previously without penicillin. Necrosis of mastoid cells found at operation, and child died same day. At necropsy a well-encapsulated abscess in temporal lobe of brain without apparent connexion with surface; no suppuration in tissues outside skull. Effect of filtrate doubtful. (Mr. McCrea.)

Case 24.—Male, elderly. Chronic osteomyelitis and sinus of thigh. Very little improvement with 280 units of filtrate in a week followed by 250 units of calcium salt, although *Staph. aureus* isolated was quite sensitive to penicillin. (Mr. E. T. Bailey.)

Comment

It is considered that in the first 19 of these patients the filtrate has been successful, sometimes to a quite unexpected degree; in the other 5 success was partial, doubtful, or absent. It is claimed, however, that this series shows that the effects expected of penicillin can be obtained by the unconcentrated filtrate when it can be applied locally to tissues infected with bacteria sensitive to it.

Since the filtrate can be readily produced and assayed in a bacteriological laboratory which has a well-trained staff, it can be used for local application to acute and chronic abscesses, sinuses, and ulcers as a substitute for the partially purified substance until the latter is available. Since the filtrate retains its potency at 4° C. for at least a week, it is possible to provide a steady supply by the production of one or two batches weekly.

A specimen was dried in several portions on April 25, 1944, in Major J. E. McCartney's laboratory by Mr. A. G. Rayner by the method which he recently described (A. G. Rayner, 1943), and when the dried substance is dissolved in sterile water some portions have retained full potency for 8 days, but others are found to have lost it entirely immediately after drying. This irregularity is being investigated.

It has been found in this series, as it has been by other observers, that, in the course of treatment of external lesions by penicillin, *B. pyocyaneus* becomes prominent and may hinder healing after the penicillin-sensitive bacteria have been reduced or banished. I have confirmed, by sensitivity tests similar to those used for assay of penicillin, that *B. pyocyaneus* (4 strains) is quite sensitive to saturated solutions (4%) of boric acid; and Dr. G. M. Palser has found in three patients that the boric acid solution reduced the suppuration due to *B. pyocyaneus* and permitted continuation of healing, although the organism recurred when boric acid was stopped before healing was complete. This seems worth further trial.

Summary

Crude penicillium filtrate containing 4 to 10 units of penicillin per c.cm. may be used successfully for treating many acute and chronic inflammations by local application, in place of the partially purified calcium salt of penicillin: the filtrate can be dependably produced and assayed in a bacteriological laboratory in useful quantities.

I wish to express thanks to the surgeons for their co-operation, to Drs. A. Martin and C. H. Carrington and others of Imperial Chemical (Pharmaceuticals) Ltd., for imparting much of their experience; to Dr. J. H. Waelsch and Mr. E. Martin for helpful suggestions; and especially to Mr. R. M. Sandercott for his enthusiasm in developing and carrying out the technical work.

REFERENCES

- Challinor, S. W., and MacNaughtan, J. (1943). *J. Path. Bact.*, **55**, 441.
Christie, R. V., and Garrod, L. P.; and others (1944). *British Medical Journal*, **1**, 513-528.
Fleming, A. (1943). *Lancet*, **2**, 434.
Florey, M. E., and Florey, H. W. (1943). *Ibid.*, **1**, 387.
Hobson, A. J., and Galloway, L. D. (1944). *Ibid.*, **1**, 164, 230.
Keefer, C. S., and others (1943). *J. Amer. med. Ass.*, **122**, 1217.
M.R.C. War Memorandum No. 12 (1944). *Use of Penicillin in War Wounds*, p. 12. H.M.S.O., London.
Rayner, A. G. (1943). *J. Path. Bact.*, **55**, 373.
Report to War Office and M.R.C. on Use of Penicillin in War Wounds, 1943.

AUTOLYSED YEAST IN THE TREATMENT OF NUTRITIONAL MACROCYTIC ANAEMIA

BY

G. R. SIPPE, M.B., B.S.

(From the Bacteriological Laboratory, Reduit, Mauritius)

Nutritional macrocytic anaemia has been reported from India, China, Africa, and Macedonia, and there is strong evidence that it is a deficiency disease due to the absence of one or more essential vitamin-like factors in the diet. It will therefore tend to occur in communities where, for economic or other reasons, foodstuffs containing adequate supplies of these factors do not normally form part of the diet. Thus in the poverty-stricken areas of Macedonia, Fairley *et al.* (1938) found the condition to be common among refugee peasants, and Wills (1934) stated that it was especially rife in Southern India, where rice is the staple food. In Mauritius it is particularly common among those members of the community who, for religious reasons, do not include beef flesh in their diet. The condition is found in both sexes, and is most prevalent during the second and third decades. It is especially apt to appear during the latter half of pregnancy, and it is probable that in cases of malarial splenomegaly the disease is aggravated by excessive destruction of normal and abnormal erythrocytes by the hypertrophied reticulo-endothelial system (Fairley *et al.*, 1938; Hughes and Shrivastava, 1931).

Although it is agreed that nutritional macrocytic anaemia is a deficiency disease, the part played by the factor or factors whose absence is responsible for the condition is obscure. Autolysed yeast (marmite) (Wills, 1934) and crude liver extracts both produce a maximal reticulocyte response, and Wills found that the factor or factors were water-soluble, heat-stable in an acid medium, and partially resistant to autoclaving in an alkaline medium; and Formijne (1940) has shown that potent substances can be obtained by extraction of beef flesh with 70 to 80% alcohol, but that the potency is lost in concentrations of 90% alcohol.

Wills *et al.* (1937), in the course of experiments on rhesus monkeys suffering from an artificially induced anaemia which was considered to be the counterpart of nutritional macrocytic anaemia in man, found anahaemin to be ineffective. This was surprising in view of its efficacy in cases of Addison's pernicious anaemia. Trials with anahaemin in nutritional macrocytic anaemia in man have given variable results. Wills and Evans (1938) considered it to be ineffective, and Napier *et al.* (1938) stated that some cases responded well while others failed to show any response. Fairley (1940) found very large doses to be effective in an Indian suffering from the condition.

Wills *et al.* (1937), in further experiments with campolon, which was fractionated into soluble and insoluble portions by saturation with ammonium sulphate, found that the insoluble