CHILDHOOD INFECTION AND ITS RELATION TO ADOLESCENT AND ADULT PULMONARY TUBERCULOSIS

A RECORD OF THE WORK OF THE BROMPTON HOSPITAL
RESEARCH DEPARTMENT DURING THE LAST 14 YEARS

PREPARED BY

A. MARGARET C. MACPHERSON, M.D., M.R.C.P.
Halley Stewart Fellow, Research Department, Brompton Hospital

[Drs. G. E. Beaumont, Dorothy Cooper (née Dow), Maurice Davidson, J. L. Livingstone, W. Ernest Lloyd, A. Margaret C. Macpherson, R. C. Wingfield, and F. H. Young have from time to time been responsible for or associated with the various investigations of the Brompton Hospital Research Department during the past fourteen years. They may be regarded, therefore, as signatories of this paper, but cannot hold themselves accountable, each and severally, for all the views expressed and for all the conclusions reached in this collected record.]

On March 21, 1928, the Medical Committee of the Brompton Hospital decided to promote and encourage research at the hospital by (1) the organization of a research department, and (2) the institution of a research fund. The Research Department accordingly came into being. Apart from the other activities covered by its constitution, it was felt that the problem forming the title of this paper was one which should be studied. A subcommittee of the staff (whose personnel has changed from time to time) was therefore appointed; suitable assistance was obtained with the help of the Research Fund, and the investigation was started. To the generous contributors to the research fund our thanks are due, but in particular to the Sir Halley Stewart Trust, all of whose contributions have been devoted to this special piece of work. The investigation is not yet finished; it may be years, if ever, before it is, but it is felt that enough has already been done to justify (1) a summary of the results from the various reports published to date; (2) an indication of the lines on which the investigation is proceeding at the present time.

Childhood tuberculosis was the subject studied for the most part when the department was first opened, and we believe that our findings have to some extent helped to build up the picture of this condition as it exists to-day. This aspect of tuberculosis is still prominent in the work of the department, and observation of the children continues. The number of children seen up to the present is 2,482. At the same time, during the fourteen years attention was constantly focused on the disturbing problem of tuberculosis in young adults—a problem of such importance at the present day, and promising to become of such urgency in the near future, that it calls for immediate action. Recognizing this, and the need for more definite observations on the subject, we set ourselves to investigate certain of its aspects. These investigations have brought to light some interesting facts and have led to our present study of treatment of symptomless pulmonary tuberculosis in young adults. Although as yet we cannot provide any definite results of treatment, we consider the matter to be of sufficient immediate interest to justify the publication of the way in which we are tackling the question.

The first task undertaken by the Research Department was the elucidation of some of the problems of childhood tuberculosis which at that time were still obscure. No appreciable amount of work on this subject had been carried out in this country. Among the established facts about childhood tuberculosis were the figures of the mortality rates, which showed a great fall during the present century. It was decided to analyse the figures for this period, and this formed the substance of Report I.

Report I (1930): "Tuberculosis Mortality in Children" (Lloyd and Dow, 1932a)

Analysis of the mortality rates from tuberculosis in children for the period 1898 to 1927 in England and Wales brought out the following interesting facts:

1. The decline in the mortality rate was greatest for children under 5 years old.
2. The rate of fall due to abdominal tuberculosis was greater than the rate of fall due to other forms, and in 1927 meningitis was by far the most fatal form of tuberculosis.
3. Although the mortality rate due to tuberculosis was greater in infancy than in later childhood, when taken in relation to the mortality rate from all causes tuberculosis is a much less frequent cause of death in young children than in those approaching adult life.
4. Before 1914 the 0-5 years age group had the highest tuberculosis mortality rate; since 1914 the tuberculosis mortality rate has been consistently higher in young adult life and middle age than in the 0-5 years group.
5. With the decrease in the mortality rate due to tuberculosis in the 0-5 years group the mortality peak in adult life changed from the 40-50 years group to the 20-25 years group.
6. The importance of exposure to tuberculous infection was demonstrated clearly by the fact that among the families attending Brompton Hospital there was a definitely increased death liability from tuberculosis in children aged 0-5 years who were brought up in contact with open cases of tuberculosis.

This report, while emphasizing the improved position of tuberculosis in childhood, underlined the continuing and relatively increasing menace of young adult tuberculosis. This was a problem that was continually demanding recognition and suggested further investigations, which were begun at this time but were reported at a later date.

In the meantime, the connexion between exposure to tuberculous infection and mortality due to the disease in infancy was an important finding which could with benefit be investigated still further and which might lead to solutions of practical value. If deaths from tuberculosis in childhood were related to exposure to infection, what was the relation between contact history and tuberculous morbidity in the living child? In other countries extensive tuberculin-testing had been carried out in the childhood populations. The results of these tests showed that, although the actual figures varied considerably in different countries and localities, the number of positive reactions to tuberculin was high in children exposed to infection when compared with those with no known contact with infection. It was thought that a similar investigation carried out in this country might be of value, and this was the next piece of work undertaken by the department.

Report II (1931): "The Incidence of Tuberculous Infection and its Relation to Contagion in Children under 15" (Lloyd and Dow, 1931, 1932b)

During 1930 and 1931 1,220 children attending the out-patient department at the Brompton Hospital were subjected to a tuberculin test. A comparison was first made between the two methods then in vogue in Europe, Scandinavia, and the U.S.A.—namely, the Pirquet test and the Mantoux intradermal test. The results showed that the Mantoux was the more accurate, and that the Pirquet was about equivalent to the Mantoux using 1 in 10,000 dilution of old tuberculin.

The results among this group of the population corresponded in the aggregate with those of observers in other countries, and demonstrated certain important facts. Taking the children as a

FIG. 1.—Percentage of children who reacted to the Mantoux test, arranged according to age and sex groups.

whole, there was a steady increase in tuberculous infection from birth to 15 years of age (see Fig. 1). If, however, the children who had a history of exposure to an open case of tuberculosis were separated from those with no known contact the results were very different. It was found that the number of positive reactors was almost doubled in the contact group taken as a whole compared with the non-contact group. The difference was still more marked in younger children, the number of positive reactors being five times greater in children under 5 years old in the contact group than in the non-contact group (see Fig. 2).

FIG. 2.—Percentage of contact and non-contact children who reacted to the Mantoux test, arranged in age groups.

The significance of direct contact with the tubercle bacillus was still further stressed by the fact that the results of tuberculin-testing in a group of children with no history of contact scarcely differed from the results obtained in a group of children with a history of contact with a closed case of tuberculosis (Fig. 2).

Although these facts had been established in other countries the results in this report were among the earliest published in England to show the importance of the connexion between contact with open cases of tuberculosis and tuberculous morbidity in children. The findings underlined the importance of contact history; the previous report had shown its importance in relation to tuberculosis mortality, and this later investigation its importance in relation to tuberculous morbidity in the living. The tuberculin test had provided us with valuable evidence of tuberculous infection in the living, and apparently healthy, child: was there any means of demonstrating, and possibly arresting, the tuberculous lesion resulting from this infection?

The low rate of mortality in childhood compared with the higher rate of morbidity gave little hope of help from clinical findings, but x-ray investigation might bring to light valuable information. Such investigations in the U.S.A. were creating much interest, and it was decided to make a systematic examination of tuberculin tests and x-ray findings in the children in the out-patient department at the Brompton Hospital.

Report III (1932): "A Comparative Study of X-ray and Mantoux Reactions in Children" (Lloyd and Dow, 1932c, 1932d)

During 1930 and 1931 500 children under 15 years of age were examined clinically and radiologically and had a tuberculin test. In order to assess the value of the radiographs without prejudice, these were examined and reported on without previous reference to the Mantoux reaction. This investigation made manifest the difficulties of interpretation of radiographs in children, and the conclusion was reached:

"that in attempting to localize the lesions in children who are shown to be tuberculous by a positive Mantoux reaction, the chest radiograph is not so helpful as might have been supposed. . . . While the chest radiograph is of great clinical value . . . its interpretation in the 'Mantoux-positive' child needs the greatest care."

Although the interpretation of these radiographs was difficult, it was evident that typical lesions of the adult type of pulmonary tuberculosis were comparatively uncommon in children under 15 years of age.

In order to keep in touch with this large group of interesting children a "follow-up" department had been by this time established, and it was possible during the ensuing years to

augment the findings of these last reports. There are now 989 children who have attended the follow-up clinic for a period of from 2 to 13 years; of these 756 are "contact" children, and the rest continue to attend either as controls or because they have evidence of tuberculous lesions. While the children are still in contact with cases of tuberculosis they come for examination at least once in every three months. After contact with infection has ceased and there is no evidence of any active tuberculous lesion the attendances are extended to once in six months.

Report IV (1933): "Reinvestigation of Mantoux-positive Children" (Lloyd and Macpherson, 1933a, 1933b)

After an interval of 1½ to 2 years 700 of the children included in Report II were subjected again to the Mantoux test. It was found that the great majority (96%) of the positive reactors in 1930 remained positive in 1½ to 2 years' time. There was, however, a change during this period in the degree of sensitivity: those who at the first testing reacted only to the stronger doses tended to become more sensitive. The figures also showed that a high degree of sensitivity was reached at an early age in contact children, whereas in the majority of non-contact children this degree of sensitivity was not reached at any age during childhood.

Throughout this and the subsequent investigations very few cases of active pulmonary tuberculosis were met with, and, in fact, obvious clinical tuberculosis among the children was conspicuously low throughout our observations.

In Report III the difficulties encountered in the interpretation of chest radiographs in children were described, and these, together with many diagnostic problems which made their appearance while observing the children in the "follow-up" department, showed the necessity for a more detailed study of the course of childhood tuberculosis. Between 1930 and 1939 considerable material had become available in the department, and by means of serial radiographs over a period of years it was possible to solve some of the problems which had presented themselves at the beginning and to build up a picture of the primary tuberculous lesion in childhood.

Report VII (1939): "Primary Tuberculosis of the Lungs in Childhood" (Macpherson, 1939a, 1939b)

This report contained a description of the primary tuberculous lesion of the lung and its subsequent course as it appeared among the children in the follow-up department. It was noted that the normal course of the primary lesion was usually uneventful and was unattended by clinical symptoms or signs, and in most cases was evident only in the x-ray film. Attention was drawn to the fact that the process of healing occupied more than one year and that throughout this period the focus was a potential source of dissemination. It was stressed that during this time it was advisable to facilitate healing by procuring a suitable environment for the child and to keep a close watch on the progress of the lesion.

Another characteristic of the primary complex was the relative unimportance of the lesion in the lung tissue compared with the lesions in the lymph nodes. It was these infected lymphatic glands which were more likely than the lung lesions to produce complications and to give rise to clinical evidence of disease. The relation between the glandular involvement and subsequent atelectasis and bronchiectasis was discussed. In only one case—that of a girl infected between the ages of 13 and 15 years—was the primary lesion seen to spread and to give rise to fibro-caseous disease of the adult type. The pulmonary lesions otherwise tended to heal by fibrosis or calcification.

So far the work carried out on childhood tuberculosis had demonstrated clearly that:

1. Primary infection by tubercle bacilli usually is uneventful clinically and the lesion formed heals satisfactorily in most cases.
2. Certain immediate and later complications may, however, arise, as a rule secondarily to the involvement of lymphatic glands in the primary complex.
3. Tuberculous meningitis has been, and still is, a serious outcome of tuberculous infection.
4. Tuberculous morbidity and mortality in childhood are markedly influenced by exposure to infection.
5. The "adult type" of tuberculosis does not present a serious problem in childhood.
6. Relatively to pulmonary tuberculosis in childhood, and even to tuberculous meningitis in childhood, pulmonary tuber-

culosis in adolescence and early adult life is a greater menace to the population.

The mortality rate from pulmonary tuberculosis among young adults showed little evidence of decreasing, and the matter continued to cause great concern. Here, then, was a subject bristling with problems for further elucidation and asking for a direct and immediate attack. Obviously it would be of immense value to establish a definite aetiological connexion between childhood infection and tuberculous disease in later life, but such an undertaking would require much material and occupy many years before there was any hope of providing a full and definite answer. However, the large numbers of observations we had been able to make in the follow-up department supplied us with some interesting evidence which to a certain extent indicated the relation between childhood infection and adolescent pulmonary tuberculosis. As some of those under observation passed from childhood to adolescence, adult types of lesions began to appear, usually between the ages of 15 and 19.

These lesions were as a rule localized to the upper third of the lung. In the majority of cases they were not accompanied by any clinical disturbances or by loss of weight, and in some instances there was not even an increase in the sedimentation rate. If left untreated the lesion showed a "pendulum" action, alternating between a spread of the lesion and healing, but nearly always with a tendency to spread and seldom healing spontaneously. The period of time between the first appearance of the lesion in the x-ray film and the first symptoms varied between three and five years.

These observations led us to a closer study of pulmonary tuberculosis in the young adult, and the next report dealt with the problem of contact history in pulmonary tuberculosis of young adults.

Report VI (1937): "Pulmonary Tuberculosis in Young Adults: The Significance of Contact History" (Lloyd and Macpherson, 1936, 1937)

Report VI contained an analysis of the family histories of 1,000 patients at the Brompton Hospital who had developed pulmonary tuberculosis between the ages of 15 and 25. The fact that there was a history of contact with tuberculosis before the onset of the disease in 40% of the cases was significant. Among those with a family history of tuberculosis, contact with a tuberculous brother or sister was found to be as frequent as contact with a tuberculous parent. Four out of five of the patients who gave a history of contact developed the disease within five years of the exposure to the last known contact with infection. This report again stressed the importance of contact history, and the conclusions also suggested that reinfection in adolescence or early adult life might play an important part in the onset of young adult lesions.

Another aspect was constantly making itself apparent: the advanced stage of tuberculosis reached before treatment was begun or was even sought by the young adult. To quote from the introduction to Report V: "Too many patients do not seek treatment until their disease is well entrenched. This undoubted fact is the reason why our tuberculosis service spends most of its energies, time, and money in endeavouring to patch up hopelessly damaged individuals by preventing further activity and inroad of the disease."

Those entrusted with the care and treatment of these young adults were repeatedly putting forth pleas for earlier diagnosis. But the fact remained that examination immediately the classical symptoms of early pulmonary tuberculosis were detected often showed considerable lesions in the x-ray films.

A hypothesis was formed "that perhaps the extensive lesions of adult disease are really in many cases deposited during adolescence with the accompaniment of slight or, at any rate, unexplained constitutional disturbance: that these lesions are carried unrecognized, perhaps extending slowly and without symptoms, until they make their presence known in adult life, either by reason of their spread past a definite point or because some environmental cause has allowed of their reactivation"; and our hitherto unpublished observations from the follow-up department on the symptomless lesions of adolescence (see above) tended to strengthen this hypothesis.

The testing of the hypothesis involved the radiological examination of a number of healthy adolescents.

Report V (1936): "Latent Adolescent Pulmonary Tuberculosis" (Wingfield and Macpherson, 1936a, 1936b)

During 1934 and 1935 such an investigation was undertaken; 2,381 young people between 14 and 21 years old, healthy and at work, were examined and the first record of mass radiography of a sample of the English population was published in 1936.

The results of this investigation showed that there was a definite incidence of pulmonary tuberculosis in 0.65% and a possible incidence in 1.08% of our sample of the adolescent population. Comparison between these figures and the calculated incidence of pulmonary tuberculosis in the population (0.4 to 0.75%) gave support to the hypothesis set forth above. Here, then, was a means of detecting at a really early stage adolescent lesions which later would give rise to clinical disease, and a means of providing the clinicians with material at a stage in which treatment should be easy compared with well-established lesions which they cannot hope to cure but only to arrest.

It was not overstating the case when it was remarked at this stage in Report V: "The significance and implications of this are tremendous." The report ended with a plea for further and more extensive investigations.

This was in 1936. It has taken much education and propaganda, directed during the ensuing years to the medical profession and the lay public alike, to bring about the development and application of the suggestion put forward at that time that mass radiography is a necessary weapon for the detection of pulmonary tuberculosis at a stage in which treatment, from the patient's point of view, could be of real value and, from the point of view of the State, far more economical than heretofore. It has been encouraging to find that recent examinations by mass radiography with more accurate apparatus and in more satisfactory conditions have produced results very similar to ours.

Since this report on latent adolescent primary tuberculosis, and following on the observations, to which reference has already been made, on the development of these lesions among those attending the follow-up department, attention has been constantly focused on the problem of treatment of those adolescents without symptoms but with radiological evidence of the disease. Recognizing that this would become a major problem should mass radiography be adopted, it was obviously necessary to examine the possibilities of adequate treatment from the point of view of healing the lesion and from the economic standpoint.

A healthy youth, working and playing hard, with no symptoms, is suddenly singled out from his companions and told that he has evidence of pulmonary tuberculosis. The first and very real difficulty is to persuade the boy or girl that further investigations are necessary. Fortunately, as a result of recent propaganda on the value of radiological examinations, this is becoming easier, but it is often more difficult to persuade the patient's doctor to take the matter seriously. These difficulties overcome, the patient is usually admitted into hospital for investigation. It is not uncommon that, apart from the radiographs, every finding is normal; there may be no abnormal physical signs, no pyrexia, a normal blood sedimentation rate, and normal gain in weight.

According to the hypothesis propounded in Report V, these adolescents are the very people who eventually will develop manifest disease. It is therefore folly to continue observation without treatment in the hope that the case in point may be the exception which proves the rule. This course is often desperately advocated by the boy or girl and the parents, and unfortunately is too often adopted by the doctor in charge. In some cases on our records the patient won the battle; treatment was postponed until symptoms appeared, with the result that only a "patching-up" of the individual was possible, while the value of the early x-ray examination had been entirely lost. Healing, surely, and not just temporary arrest, should be possible, and should be the aim when these early lesions are under consideration.

Some form of treatment, therefore, is called for, and at this stage the economic factor comes into prominence and must influence the course followed. At the present time such cases, when detected, are handled in one of the four following ways: (1) they are either ignored or are merely observed for a period; (2) they are kept under careful observation, with improvement

in their home environment; (3) they are sent to a sanatorium; (4) they are, in only a few instances, treated by artificial pneumothorax.

Treatment of Symptomless Adolescent Pulmonary Tuberculosis

For those who can afford the time and treatment it is probable that with rest at home, followed by gradually increasing activity and then by years of sheltered life, healing may take place. It is, however, only a favoured few who can afford this; for the majority a more rapid return to work is necessary. Moreover, it is unusual to find any person who, while apparently in good health, will submit to this treatment and at the same time retain a healthy outlook on his "illness," also the capacity to return to a normal life at the end of it.

For the majority, therefore, some other course is advised. Sanatorium treatment is the usual one adopted for these patients. Owing to the healthy condition of the patient on admission to the sanatorium and the constant urge to get home and back to work, it is usually impossible to keep him or her at the institution long enough to hope to effect a cure. There may be a temporary improvement in a few cases, but in the majority there is no material change in the x-ray picture, and the expected spread of the disease comes sooner or later after a variable interval.

It seems that artificial pneumothorax provides a more satisfactory prospect; we believe that this is the correct treatment for these cases, and it is the one which has been adopted. These symptomless patients should be ideal subjects for such treatment. As they have no constitutional disturbance it does not seem that they require prolonged bodily rest at hospital or sanatorium. Some measure of local rest to the lung is provided by a pneumothorax, and it is not necessary to attempt complete collapse of the lung: a shallow pneumothorax sufficient to control and immobilize the diseased lung, partially or completely, is what is required, and it is suggested that the pneumothorax be maintained for not longer than two or three years, by which time, on the analogy of the tuberculous joints, it is hoped the lesion will have healed. A shallow pneumothorax for a slight lesion of the lung is unlikely to give rise to complications, and causes no discomfort or subjective incapacity.

Once the collapse is established daily supervision is not necessary, and, provided that the home conditions are good, the patient need only stay in hospital for a short period. Where the home is unsatisfactory, or the patient's general condition is below normal, a period in a sanatorium may be necessary. A number of our patients have returned to work or school without any ill effects a short time after the induction of the pneumothorax. If the infiltration is bilateral the position is unaltered and the treatment is the same. A shallow bilateral artificial pneumothorax is tolerated as easily as a unilateral one, and in the case with a unilateral artificial pneumothorax there should be no hesitation in inducing a bilateral artificial pneumothorax should a tuberculous deposit appear in the other lung.

Regular supervision is ensured by the weekly attendance for refills; and this is important, because without such a tie healthy young people will naturally tend to become less and less frequent in their attendance over a period of years.

Here, then, is a form of treatment which offers a reasonable prospect of cure without prolonged exile from normal life and work. We would emphasize that, as yet, it is impossible to assess its value, but we suggest it as a reasonable alternative to other methods in use which, in our experience, are unsatisfactory in their final results or are impracticable for the majority of patients.

REFERENCES

- Lloyd, W. E., and Dow, Dorothy, J. (1931). *British Medical Journal*, 2, 183.
 — (1932a). *Bromp. Hosp. Rep.*, 1, 20.
 — (1932b). *Ibid.*, p. 57.
 — (1932c). *British Medical Journal*, 1, 701.
 — (1932d). *Bromp. Hosp. Rep.*, 1, 61.
 — and Macpherson, A. Margaret C. (1933a). *British Medical Journal*, 1, 818.
 — (1933b). *Bromp. Hosp. Rep.*, 2, 99.
 — (1936). *British Medical Journal*, 2, 1131.
 — (1937). *Bromp. Hosp. Rep.*, 6, 31.
 Macpherson, A. Margaret C. (1939a). *Brit. J. Tuberc.*, 33, 79.
 — (1939b). *Bromp. Hosp. Rep.*, 8, 30.
 Wingfield, R. C., and Macpherson, A. Margaret C. (1936a). *British Medical Journal*, 1, 741.
 — (1936b). *Bromp. Hosp. Rep.*, 5, 10.

PERIPHERAL ARTERIAL EMBOLISM

BY

H. AGAR, F.R.C.S.

Major, R.A.M.C.; Surgical Specialist

The surgical treatment of embolic occlusion of a peripheral artery has been developing over a long time. In 1907 both Sampson Handley and Moynihan attempted to relieve the condition by operation. In this country little attention was paid to their efforts, but the Scandinavian surgeons, led by Einar Key, developed a satisfactory operative technique, and in 1925 Jefferson performed the first successful embolectomy in England.

The incidence of embolic occlusion it is impossible to assess, but, although the condition is not common, I believe many cases are missed. Griffiths (1938) reports 8 cases in a three-year period in Manchester; and in St. James's Hospital, Leeds, from 1936 to 1939, with approximately 28,000 admissions, there were 7 cases in 5 patients, all of which came under my care.

The embolus may arise from the pulmonary veins, a vegetation on the mitral valve, a mural thrombus of the left auricle, or an atheromatous plaque of the aorta; or from a systemic vein in the presence of a patent foramen ovale. The cardiac origin is by far the commonest, and is often associated with auricular fibrillation. It was the focus in all my cases. Emboli usually become impacted at the bifurcation of an artery, for it is here that the lumen is most suddenly reduced in size. Of 382 Swedish cases (quoted by Riddell, 1937) the embolus was located in the common femoral in 54%, in the common iliac in 17%, in the popliteal in 11%, in the upper limb in 12%, and in the aorta in 4.5%. The few cases operated on in England suggest about the same percentages, and in my series four of the emboli were impacted at the bifurcation of the common femoral artery, two in the popliteal, and one at the bifurcation of the aorta.

Diagnosis

Diagnosis is usually easy, and depends on pain, pallor, paralysis, and loss of arterial pulsation in a limb. It is reported that immediately the artery is occluded the patient complains of sudden severe pain, often located over the site of the embolus. This description is inaccurate, for although pain may be severe it may also be absent. None of my patients complained of severe pain and only four of slight pain. In the case of occlusion of the bifurcation of the aorta the operative findings indicated that most probably the blockage had been sudden and complete; for there was no secondary thrombus, and yet the patient did not complain of real pain but only of abdominal discomfort. Deitch (1936) speaks of "agonizing colicky pain," and Riddell of "intense agonizing pain." The intensity or absence of the pain therefore must not influence the diagnosis.

All my patients did complain of numbness and loss of use in the affected limb. In five in whom the incident occurred while they were in hospital undergoing treatment for auricular fibrillation it was the first complaint to be made to the ward sister. The circulation was probably impaired for a few minutes before nerve symptoms appeared. In a case of occlusion at the bifurcation of the common carotid Griffiths (1938) reports sudden contralateral hemiplegia. A general pallor with some blotchy cyanosis was observed in all the affected limbs and was present within a short time of the first complaint by the patient. The limbs felt colder than the normal, and the area of anaesthesia was of the stocking type but not reaching as high as the site of impaction.

There is always loss of arterial pulsation beyond the blockage, but this may be misleading in the case of obstruction at the bifurcation of the common femoral artery. The common femoral artery is 1½ in. long, and, to the unwary, palpable pulsation below Poupart's ligament may give the impression that the femoral arteries are patent and that the blockage is in the popliteal. I found that if a sphygmomanometer cuff was applied round the mid-thigh no oscillations in the mercury column were visible at any pressure in cases of femoral blockage, but that they were easily seen in cases of popliteal